

Embedding a Pair of Graphs in a Surface, and the Width of 4-dimensional Prisms

Francisco Santos · Tamon Stephen · Hugh Thomas

Received: 11 April 2011 / Revised: 20 May 2011 / Accepted: 25 May 2011 /
Published online: 21 June 2011
© Springer Science+Business Media, LLC 2011

Abstract A *prismatoid* is a polytope with all its vertices contained in two parallel facets, called its *bases*. Its *width* is the number of steps needed to go from one base to the other in the dual graph. The first author recently showed that the existence of counter-examples to the Hirsch conjecture is equivalent to that of d -prismatoids of width larger than d , and constructed such prismatoids in dimension five. Here we show that the same is impossible in dimension four. This is proved by looking at the pair of graph embeddings on a 2-sphere that arise from the normal fans of the two bases of Q .

Keywords Polytope · Polytope diameter · Hirsch conjecture · Graph embedding · Prismatoid

1 Prismatoids and Pairs of Maps

The first author recently constructed counter-examples to the Hirsch conjecture [2] via the combination of the following two results:

F. Santos
Departamento de Matemáticas, Estadística y Computación, Universidad de Cantabria,
Av. de los Castros 48, 39005 Santander, Spain
e-mail: francisco.santos@unican.es

T. Stephen (✉)
Department of Mathematics, Simon Fraser University, 8888 University Drive, Burnaby,
British Columbia V5A 1S6, Canada
e-mail: tamon@sfu.ca

H. Thomas
Department of Mathematics and Statistics, University of New Brunswick, Fredericton,
New Brunswick E3B 5A3, Canada
e-mail: hugh@math.unb.ca

- If a d -dimensional prismatoid Q has width larger than d then it is possible to construct from it a polytope P that violates the Hirsch conjecture.
- There exist 5-prismatoids of width larger than 5.

Here, a *prismatoid* is a polytope that has all its vertices lying in two parallel facets Q^+ and Q^- (called its *bases*), and its *width* is the dual graph distance between those two facets. That is, the width of Q is the minimum number of steps needed to go from Q^+ to Q^- , where a step consists in moving from one facet of Q to an adjacent facet via a ridge.

The dimension of the non-Hirsch polytope P in the construction is much higher than that of Q (it equals $|\text{vertices}(Q)| - |\dim(Q)|$) but still the question arose whether the dimension $d = 5$ for Q is the smallest possible. An easy argument shows $d = 3$ cannot work, and $d = 4$ was left open in [2]. In this paper we answer this question in the negative:

Theorem 1.1 *The width of a 4-dimensional prismatoid is at most four.*

The proof of this theorem uses the following reduction, also from [2]. Let $Q \subset \mathbb{R}^d$ be a prismatoid with bases Q^+ and Q^- . Consider the bases as simultaneously embedded into \mathbb{R}^{d-1} and let G^+ and G^- be the intersections of their normal fans with the unit sphere S^{d-2} . G^+ and G^- are geodesic cell decompositions of S^{d-2} , what we call *geodesic maps*. Let H be their common refinement: cells of H are all the intersections of a cell of G^+ and a cell of G^- . Then:

- All facets of Q other than the two bases appear as vertices of H .
- The facets adjacent to Q^+ (respectively to Q^-) appear in H as the vertices of G^+ (respectively, of G^-).
- Adjacent facets of Q appear as vertices connected by an edge of H .

We call *width* of the pair of geodesic maps (G^+, G^-) the minimum graph distance along H from a vertex of G^+ to a vertex of G^- . The following lemma is clear from the above discussion:

Lemma 1.2 *The width of a prismatoid $Q \subset \mathbb{R}^d$ equals two plus the width of its corresponding pair of maps (G^+, G^-) in S^{d-2} .*

In particular, the width of a 4-prismatoid is two plus the width of a pair of maps in the 2-sphere. In the 2-sphere, knowing the (embedded) graph of a map is enough to recover the whole map, so we rather speak of a *pair of graphs* embedded in it. If the two graphs intersect other than in the interior of edges then a small perturbation cannot decrease their width, so in what follows we assume that this does not happen. Moreover, since the maps we are interested in are geodesic, an edge of one cannot intersect an edge of the other more than once, and they do so transversally; that is, the four branches that come out from an intersection point alternate between G^+ and G^- when we look at them cyclically. Putting together these conditions, we introduce the following notion, which looks natural in the context of geometric graph theory:

Definition 1.3 Let S be a closed surface and let G^+ and G^- be two graphs embedded in it. We say that (G^+, G^-) is a *normal pair of graphs* if G^+ and G^- intersect only




Fig. 1 Two graphs embedded in the torus (left and center) and the normal map they form (right), of width five

in the interior of edges, each edge of one intersects each edge of the other at most once, and all the intersections are transversal.

We also consider a somewhat stronger condition:

Definition 1.4 We say that (G^+, G^-) form a *strongly normal pair of graphs* if they are a normal pair of graphs and in addition the intersection of an edge of G^+ with the interior of a face of G^- has at most one component (and also with the roles of G^+ and G^- interchanged).

This condition is also satisfied by any pair of geodesic maps, since it amounts to saying that an edge of G^+ cannot leave the interior of a geodesic cell of G^- and then re-enter it, which follows from convexity.

We call *width* of the normal pair the minimum distance from a vertex of G^+ to one of G^- along the common refinement of the two graphs. Put differently, the width is one plus the minimum number of points of $G^+ \cap G^-$ that you need to go through in order to get from a vertex of G^+ to one of G^- . We assume that G^+ and G^- intersect at least once. (Otherwise the width can be considered infinite, but the question we address in this paper is meaningless.)

As an example, the two graphs in Fig. 1 form a normal pair of width five in the torus, and the same idea shows the existence of normal pairs of arbitrarily large width in any closed surface of non-positive Euler characteristic. This example is taken from [1], where it is used to show the existence of prisms of dimension five of arbitrarily large width. Our main result, which implies Theorem 1.1, is that such pairs do not exist in the 2-sphere:

Theorem 1.5 *Every normal pair of graphs in the sphere or the projective plane has width two.*

We prove this theorem in Sect. 3, using an argument based on the simple-connectedness of the sphere. In Sect. 4, we use the notion of Euler characteristic to provide an independent proof of the theorem, under the assumption that the pair of graphs is strongly normal (which is still sufficient to deduce Theorem 1.1).




Fig. 2 A normal pair of graphs (*left*), its terminal part (*center*) and its deep part (*right*)

2 Terminology and Initial Observations

Start with an arbitrary normal pair of graphs (G^+, G^-) in an arbitrary closed surface S . We introduce the following nomenclature: we call G^+ and G^- the *positive* and *negative* graphs, and let H be the common graph induced by the embedding. In H we have three types of vertices: *positive* (vertices of G^+), *negative* (vertices of G^-), and *crossings*. We also have four types of edges in H : each edge is *positive* or *negative* depending on whether it is contained in an edge of G^+ or one of G^- , and it is *deep* or *terminal* depending on whether its two end-points are crossings or at least one of them is not. See Fig. 2. Observe that edges ending in no crossings (that is, edges of G^+ or G^- that do not cross the other graph of the pair) do not contribute to the width, so we may as well contract them and assume our pair of graphs does not have any. In order to be allowed to use contractions, we permit our graphs not to be simple.

We also show that there is no loss of generality in assuming that S is orientable.

Lemma 2.1 *If (G^+, G^-) is a normal pair in a non-orientable surface S , then the pull-back of (G^+, G^-) via the double cover $S' \rightarrow S$ from an orientable surface S' yields a normal pair $(G^{+'}, G^{-'})$ in S' with the same width.*

Proof Every path from G^+ to G^- in S lifts to a path (actually to two paths) from $G^{+'}$ to $G^{-'}$ in S' , of the same length. Reciprocally, every path in S' projects to S . \square

3 Proof via Simple-Connectedness

As in the previous section, start with an arbitrary normal pair of graphs (G^+, G^-) in an arbitrary closed surface S . This proof concentrates on the *deep* subgraph of H , which we denote H_0 . This is the subgraph consisting of deep edges or, equivalently, the subgraph induced by the crossing points. Observe that if H_0 has no edges then the width of the pair (G^+, G^-) is trivially two: every terminal positive edge is adjacent to a terminal negative edge, and those two edges from a path from G^+ to G^- of length two. More generally:

Lemma 3.1 *The width of the pair (G^+, G^-) exceeds two if, and only if, every vertex of H_0 is incident to either two deep positive edges or two deep negative edges.*

Proof The width equals two if, and only if, there is a crossing incident to both a terminal positive and a terminal negative edge. If this does not occur then every crossing is incident to two deep edges of one sign (and to zero, one or two of the other). \square

By Lemma 2.1, there is no loss of generality in assuming that S is orientable. We therefore assume that S is orientable, and give an orientation to it. In particular, this gives a meaning to the expressions “left” and “right” of a directed edge.

Definition 3.2 A path C in H_0 is *well-formed* if when traversing C we always turn right when passing from a negative to a positive edge and turn left when passing from a positive to a negative edge. (The path is allowed to “go straight” at some of the vertices, meaning that it keeps walking along the same edge of G^+ or G^- .) A *well-formed loop* is a cycle with the same property except (perhaps) at its base vertex.

Observe that the definition is independent of the direction in which we traverse the path or cycle, since reversing the direction exchanges left and right turns but it also exchanges passings from positive to negative and passings from negative to positive. The following two lemmas prove Theorem 1.5.

Lemma 3.3 *If a normal pair of graphs in a closed surface has width larger than two then its deep part contains a well-formed loop.*

Proof Starting at any deep edge e we can build a well-formed path as follows: go straight on edges of the same sign as e until you cannot go further. When this happens, Lemma 3.1 guarantees that you are at a vertex incident to two edges in H_0 of the other sign, one to the left and one to the right. Choose the one that makes the path be well-formed and continue.


Since this process can always be continued, at some point the path will cross with itself. The first time this happens, you declare the repeated vertex to be the base for the loop obtained, which is well-formed by construction. (Observe that the original edge e may not belong to the loop.) \square

Lemma 3.4 *A well-formed loop in a normal pair of graphs in a closed surface cannot bound a disc.*

Proof Let L be a well-formed loop with base vertex v and suppose, to get a contradiction, that it bounds a disc D . We are going to show how to construct another well-formed loop in D , which necessarily should bound a smaller disc $D' \subset D$. Since the same applies to D' , we can construct an infinite sequence of loops bounding smaller and smaller discs, a contradiction.

To construct L' observe that, by normality of the pair of maps, the loop L has at least two turns, apart from the possible turn at the base point v . Since left and right turns alternate, at least one of the turns is to the left when we traverse L in the clockwise direction (clockwise is understood with respect to D and to the given orientation in it). Let u be the vertex where this happens. Put differently, u is a “concave” or “reflex” vertex in the boundary of D .

Fig. 3 The situation in the proof of Lemma 3.4



By Lemma 3.1, apart from the positive and negative edges incident to u in L , H_0 has at least one more deep edge e coming out from u . Concavity of D at u implies that e goes toward the interior of D . We use that edge e to start a well-formed path, as in the proof of Lemma 3.3, and stop when the path intersects either itself or the loop L . If the path intersects itself then we have found a well-formed loop contained in D . If it intersects L , then let w be the point where this happens and let C be the well-formed path that we have obtained from u to w through the interior of D . Let C_1 and C_2 be the two paths in which L is divided by the vertices u and w (see Fig. 3). Our claim is that C together with one of C_1 or C_2 is a well-formed loop.

Indeed, assume first that $v = w$. Then we start our loop from v using the interior path C and when we arrive at u one of the turns to C_1 or C_2 is well-formed, which closes the loop. If v is different from both u and w , assume that v lies in C_1 . The key idea is that at both u and w at least one of the turns, from C to C_1 or from C to C_2 , is valid for a well-formed path. Then:

1. If both turns from C to C_1 are valid, then C and C_1 give a well-formed loop based at v .
2. If the turn from C to C_1 is invalid at, say, u , then C and C_2 give a well-formed loop based at w , because the turn from C to C_2 is valid at u .

□

Proof of Theorem 1.5 Lemmas 3.3 and 3.4 imply that a normal pair of graphs of width larger than two has a loop that does not bound a disc, which cannot happen in the 2-sphere. For the projective plane we use the reduction of Lemma 2.1. □

4 Proof via Euler Characteristic

In this section, we provide an alternative proof of Theorem 1.5, under the stronger hypothesis that the pair (G^+, G^-) are strongly normal.

As discussed in Sect. 2, to prove Theorem 1.5, it suffices to assume that (G^+, G^-) are a pair of graphs embedded in a sphere S^2 , such that each edge of G^+ crosses some edge of G^- , and vice versa. Note that the reductions to this case preserve the property that the pair of graphs is strongly normal. We again study the graph H induced by the embedding. In this case, we focus on the complex of 0-, 1- and 2-dimensional cells (vertices, edges and faces) defined by the embedding.

We assume that H is connected, since it suffices to consider some component of H . The connectedness of H implies that the faces defined by H are contractible.

We will calculate the Euler characteristic of the sphere S^2 , using the decomposition induced by H . Since the faces of the decomposition are contractible, this is just the number of vertices of H , minus the number of edges, plus the number of faces. Our strategy will be to bound this sum by a contribution for each face. Assuming that the width of (G^+, G^-) is greater than two, we will show that the contribution for each face is non-positive, and thus that the Euler characteristic of S^2 is non-positive. Since the Euler characteristic of S^2 is positive, this is a contradiction.

We now discuss how to bound the Euler characteristic by a contribution for each face. Let F be a face of H . Consider (as a first step) the following sum:

$$\begin{aligned} c(F) = & \frac{1}{4} (\text{the number of crossing vertices incident to } F) \\ & - \frac{1}{2} (\text{the number of edges of } F) \\ & + 1. \end{aligned}$$

Note that, in this definition, we count edges and vertices with multiplicity. That is to say, we count them going around the boundary of the face, so that, if we revisit an edge or vertex, we count it again.

We have the following:

Lemma 4.1 *Let (G^+, G^-) be a normal pair of graphs such that H is connected. Then: $\chi(S^2) = |V(G^+)| + |V(G^-)| + \sum_F c(F)$.*

Proof Each of the crossing vertices contributes $+1$ to the Euler characteristic which is shared among the four faces that meet at it, and each edge contributes -1 to the Euler characteristic, which is shared between the two faces it lies on. \square

Since we want to bound $\chi(S^2)$ by a sum over the faces, we must decide, for each signed vertex, where to include it in the sum.

Suppose v is a positive vertex on face F , and that the two crossing vertices of F adjacent to v on F are x and y . We say that v *counts against* F if x and y are not incident to the same edge of G^- . We make the similar definitions for negative vertices.

Lemma 4.2 *If (G^+, G^-) is a strongly normal pair of graphs, then every signed vertex v counts against at least two faces.*

Proof Suppose v is positive. There must be at least two different edges of G^- which are incident to crossing vertices of H at distance one from v , since otherwise the single edge of G^- which is incident to all the crossing vertices at distance one from v would necessarily leave and re-enter the interior of the same G^+ -face, contrary to our assumption of strong normality. Therefore, as we visit in cyclic order the crossing

vertices in H at distance one from v , which edge from G^- they intersect must change at least twice, so v counts against at least two faces of H . \square

Let $d(F) = c(F) + \frac{1}{2}$ (the number of signed vertices counting against F). Given the previous lemma, the following lemma is immediate:

Corollary 4.3 *If (G^+, G^-) is a strongly normal pair of graphs, with H connected, then $\chi(S^2) \leq \sum_F d(F)$.*

We finish by showing that $d(F) \leq 0$ for all F , under the assumption that the width of (G^+, G^-) is greater than two.

Lemma 4.4 *Assume that (G^+, G^-) is a normal pair of graphs and has width greater than two. Let F be a face of H . Then $d(F) \leq 0$.*

Proof Write e for the number of edges of F . Write x for the number of crossing vertices. Now $d(F) \leq 1 + x/4 - e/2 + (e - x)/2 = (4 - x)/4$. We remark that x counts the number of times that the sign of the edge changes as we travel along the boundary of F and thus must be even. It is clear from the above that if F has four or more crossing vertices then we are done.

If F has two or more signed vertices, then it must have four or more crossing vertices. (Due to the contraction discussed in Sect. 2, signed vertices cannot be at distance one, and since the signs of edges alternate, for signed vertices to be at distance two, they would have to have opposite signs, and thus we would have a path of length two.)

If F has no signed vertices, then again it must have at least four crossing vertices (since the signs of the edges alternate, there are an even number of edges, and two edges would mean that the corresponding edges of G^+ and G^- cross twice).

If F has one signed vertex v , it may only have two crossing vertices. In this case, however, F is a triangle, and v does not count against F , so $d(F) = c(F) = 0$. \square

Acknowledgements Francisco Santos was supported in part by the Spanish Ministry of Science through grant MTM2008-04699-C03-02. Tamon Stephen and Hugh Thomas were supported in part by NSERC Discovery Grants. The authors would like to thank the referees for helpful suggestions.

References

1. Matschke, B., Santos, F., Weibel, C.: The width of 5-prismatoids and smaller non-Hirsch polytopes (in preparation)
2. Santos, F.: A counter-example to the Hirsch Conjecture. Preprint [arXiv:1006.2814](https://arxiv.org/abs/1006.2814), June 2010