

**Erratum: Evidence for a Smooth Onset of Deformation in the Neutron-Rich Kr Isotopes**  
**[Phys. Rev. Lett. 108, 062701 (2012)]**

M. Albers, N. Warr, K. Nomura, A. Blazhev, J. Jolie, D. Mücher, B. Bastin, C. Bauer, C. Bernards, L. Bettermann, V. Bildstein, J. Butterworth, M. Cappellazzo, J. Cederkäll, D. Cline, I. Darby, S. Das Gupta, J. M. Daugas, T. Davinson, H. De Witte, J. Diriken, D. Filipescu, E. Fiori, C. Fransen, L. P. Gaffney, G. Georgiev, R. Gernhäuser, M. Hackstein, S. Heinze, H. Hess, M. Huyse, D. Jenkins, J. Konki, M. Kowalczyk, T. Kröll, R. Krücken, J. Litzinger, R. Lutter, N. Marginean, C. Mihai, K. Moschner, P. Napiorkowski, B. S. Nara Singh, K. Nowak, T. Otsuka, J. Pakarinen, M. Pfeiffer, D. Radeck, P. Reiter, S. Rigby, L. M. Robledo, R. Rodríguez-Guzmán, M. Rudigier, P. Sarriuguren, M. Scheck, M. Seidlitz, B. Siebeck, G. Simpson, P. Thöle, T. Thomas, J. Van de Walle, P. Van Duppen, M. Vermeulen, D. Voulot, R. Wadsworth, F. Wenander, K. Wimmer, K. O. Zell, and M. Zielinska

(Received 18 April 2015; published 5 May 2015)

DOI: 10.1103/PhysRevLett.114.189902

PACS numbers: 25.70.De, 27.60.+j, 29.30.Kv, 29.38.Gj, 99.10.Cd

In our Letter, the  $E2$  transition operator used was not  $T^{(E2)} = e_\pi \hat{Q}_\pi + e_\nu \hat{Q}_\nu$ , as quoted in the text on page 3. Instead, the operator  $T^{(E2)} = e_\pi(s_\pi^\dagger \tilde{d}_\pi + d_\pi^\dagger \tilde{s}_\pi)^{(2)} + e_\nu(s_\nu^\dagger \tilde{d}_\nu + d_\nu^\dagger \tilde{s}_\nu)^{(2)} + \chi_\pi(d_\pi^\dagger \tilde{d}_\pi)^{(2)} + \chi_\nu(d_\nu^\dagger \tilde{d}_\nu)^{(2)}$  was used to calculate the  $B(E2)$  values in the IBM-2. This error does not change the main result and the conclusion of the published Letter.