



HAL
open science

The Biased Transformation and its Application in Goodness-of-Fit Tests for the Beta and Gamma Distribution

Mathias Raschke

► **To cite this version:**

Mathias Raschke. The Biased Transformation and its Application in Goodness-of-Fit Tests for the Beta and Gamma Distribution. *Communications in Statistics - Simulation and Computation*, 2009, 38 (09), pp.1870-1890. 10.1080/03610910903152631 . hal-00544826

HAL Id: hal-00544826

<https://hal.science/hal-00544826v1>

Submitted on 9 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The Biased Transformation and its Application in Goodness-of-Fit Tests for the Beta and Gamma Distribution

Journal:	<i>Communications in Statistics - Simulation and Computation</i>
Manuscript ID:	LSSP-2009-0039.R1
Manuscript Type:	Original Paper
Date Submitted by the Author:	22-Jun-2009
Complete List of Authors:	Raschke, Mathias; Fa. M. Raschke
Keywords:	beta distribution, gamma distribution, goodness-of-fit test, transformation
Abstract:	The biased transformation is introduced for continuous random variables. It is proved in the paper that the biased transformation works for the generalized Pareto, the generalized extreme value and the normal distribution. The biased transformation works for continuous distributions of the exponential family as an approximation; this is demonstrated with the aid of simulations. The new transformation can be used for the Anderson-Darling test for the gamma and beta distribution. The power of the new test procedure is validated by simulations. The practical relevance of the new test for the beta distribution is demonstrated in an analysis of meteorological data.



Biased transformation and Goodness-of-Fit Test

The Biased Transformation and its Application in Goodness-of-Fit Tests for the Beta and Gamma Distribution

Mathias Raschke

Fa. M. Raschke

Teichgasse 18, 99423 Weimar, Germany

Phone +49(0)36431903238

E-Mail MathiasRaschke@t-online.de

Abstract: The biased transformation is introduced for continuous random variables. It is proved in the paper that the biased transformation works for the generalized Pareto, the generalized extreme value and the normal distribution. The biased transformation works for important continuous distributions of the exponential family as an approximation; this is demonstrated with the aid of simulations. The new transformation can be used for the Anderson-Darling test for the gamma and beta distribution. The power of the new test procedure is validated by simulations. Finally, a first powerful test is available for the beta distribution although the power function is poor. The χ^2 -test has lower power. The Kolomogorov-Smirnov test for a fully specified distribution has almost no power in the application for the beta distribution with estimated parameters. The practical relevance of the new test for he beta distribution is demonstrated in an analysis of meteorological data.

Keywords: beta distribution, gamma distribution, goodness-of-fit test, transformation

1 Introduction

The following transformation and test procedures have been researched by chance as the author was trying out different transformation methods in flood statistics and was looking into publications for a goodness-of-fit test for the gamma distribution. It is possible to use the found biased transformation to test the goodness-of-fit for the gamma distribution. But there exist already efficient tests for the gamma distribution (Stephens 1986). Except this, there is no practicable and specific goodness-of-fit test for the beta distribution so far. This was the motivation to research the biased transformation for some random distribution and to research the power of the test using biased transformation for the gamma and the beta distribution.

The beta and the gamma distributions are very famous and are explained in different publications e.g. by Johnson et al. (1994, 1995).

Biased transformation and Goodness-of-Fit Test

1
2
3 The goodness-of-fit of an estimated distribution can be tested by the Chi-Squared test for a
4 sample. This and related tests are explained by Moore (1986). There is no special consideration
5 of the type of distribution in this test. The EDF-tests (empirical distribution function tests)
6 consider the type of distribution. Different types of EDF-tests are published like the Anderson
7 Darling test (Anderson and Darling 1952, 1954), Cramer von Mises Test and Watson Test –
8 both introduced by Watson (1961), the Kuiper test (Kuiper 1960) and the famous Kolmogorov-
9 Smirnov test (Kolmogorov 1933). The critical values of EDF-test are published by Stephens in
10 different publications (1974, 1976) for different types of distributions. A summary for the EDF-
11 test for different types of distributions is given by Stephens (1986). The EDF-test for normality
12 with the best power is the Anderson Darling test (Landry and Lepage 1992). The moment tests
13 are not often used. The basics of this test type are explained by Bowman and Shento (1986). The
14 theory of a moment test is discussed for beta distributions by Li and Papadopoulos (2002) but
15 contains a mistake (a moment is divided by itself) and the power of this test approach is small
16 (Li and Papadopoulos 2002, Tab. 1-3).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Generally each continuously variable can be transformed into another random variable by any
34 continuously function which is defined in the ranges of these random variables. The power
35 transformation of Box and Cox (1964) is an important family of transformation. The goal of
36 these transformations is the normalisation of the random variable. A further transformation
37 implies the application of Copulas (Sklar 1959, Carley and Taylor 2000). The continuously
38 random variables are transformed to uniform distributed variables by using the marginal
39 distribution or by using the EDF. The approach to use a transformation for goodness-of-fit test
40 is introduced already by Quesenberry (1986).
41
42
43
44
45
46
47
48
49

50 The biased transformation (BT) is introduced here and is explained in the second section. In the
51 third section is shown that BT works as an approximation for the normal, gamma and beta
52 distribution. Furthermore the BT is applied in an EDF- test procedure for the gamma and the
53 beta distribution in sections 4 and 5. The benefit of the new test procedure for the beta
54 distribution is demonstrated in the section 6 with weather data. The results are summarised and
55 discussed in the last section. The theoretical research for the BT for the generalised Pareto
56
57
58
59
60

Biased transformation and Goodness-of-Fit Test

distribution (GPD), the generalised extreme value distribution (GEV) and the normal distribution is given in the appendix.

2 The Biased Transformation

A continuous random variable X_A with cumulated distribution function (CDF) F_A can generally be transformed to another X_B with CDF F_B with the transformation function $x_B(x_A)$, which is

$$x_B = F_B^{-1}[F_A(x_A)]. \quad (1)$$

F^{-1} is the symbol for the inverse function with $x=F^{-1}(F(x))$. A further biased transformation (BT) is introduced here with

$$x_D = F_C^{-1}[F_B(x_A)]. \quad (2)$$

The type of CDF F_A and F_B is identical and likewise the type of CDF F_C and F_D is identical, but the parameter vectors θ don't need to be equal - $\theta_A \neq \theta_B$ and $\theta_C \neq \theta_D$. Furthermore the type of F_A and F_B need not to be identical to distribution F_C and F_D . The functions and distributions are distinguished by the capitals A to D. All CDF are continuous in the relevant ranges. The probability density function (PDF) is the derivative $f(x)$ of the CDF $F(x)$ with respect to the random variable. This includes the assumption that the CDF is differentiable in the relevant range. The CDF F_B and F_C are of interest and be used in this paper. The corresponding random variables X_B and X_C are not used here.

The transformation implies that

$$F_D[x_D(x_A)] = F_A(x_A) \quad \text{and} \quad (3)$$

$$f_D[x_D(x_A)] = f_A(x_A) f_C[x_D(x_A)] / f_B(x_A). \quad (4)$$

The transformation is called "biased" because $F_A(x_A) \neq F_B(x_A)$ in equation 2. The biases are described by the bias function $F_B(F_A)$ where $F_B(F_A) = F_B(x_A)$ with $F_A(x_A)$. Equation 2 is the transformation $x_D(x_A, \theta_B, \theta_C)$, and from this the functions $\theta_D(\theta_A, \theta_B, \theta_C)$ can be deduced. The transformation is an assumption until it is proved for the concrete distribution types. It has to be shown in a proof that either equation 3 or 4 is right for the functions $x_4(x_1, \theta_2, \theta_3)$ and $\theta_4(\theta_1, \theta_2, \theta_3)$.

The BT works for some distributions. This is demonstrated by the research of the generalised Pareto distribution (GPD) and the generalised extreme value distribution (GEV). **The distributions,**

Biased transformation and Goodness-of-Fit Test

theorems and simple proofs are formulated in the appendix. Furthermore, the BT works exactly for the normal distribution with the PDF

$$f(x) = 1/(\sigma\sqrt{2\pi})\exp\left[-(x-\mu)^2/(2\sigma^2)\right], \quad \sigma > 0 \quad (5)$$

The theorem and the proof for BT of the normal distributed X_A and X_D is described in the appendix.

3 The BT as an Approximation for Some Continuous Distributions of the Exponential Family

The exponential family includes the normal, the gamma and the beta distribution. The PDF for the gamma distribution is denoted $\Gamma(\alpha)$ as a gamma function of α with

$$f(x) = (x/\lambda)^{\alpha-1} \exp(-x/\lambda) / [\lambda \Gamma(\alpha)], \quad x \geq 0, \lambda > 0, \alpha > 0.$$

The PDF for the beta distribution is denoted $B(\alpha, \beta)$ as beta function of α and β with

$$f(x) = x^{\alpha-1} (1-x)^{\beta-1} / B(\alpha, \beta), \quad 0 \leq x \leq 1, \alpha > 0, \beta > 0.$$

There are no elementary solutions for the integral of the CDF of the beta distribution, the gamma distribution and the normal distribution. But the CDF can be computed numerically.

The BT can be used for the gamma, beta and normal distribution as an approximation.

Assumption: If X_A with F_A is normal, beta or gamma distributed, and F_B is the same type of CDF as F_A with $\theta_A \neq \theta_B$ but not with $|\theta_A - \theta_B| \gg 0$, then X_A can be transformed with F_C according equation 2 to X_D . F_C has any θ_c and F_D of X_D can be approximated by the same type of F_C .

The basic idea for the validation is the following. Samples of a uniform distributed random variable U ($0 \leq u \leq 1$) can be simulated. From this sample, the sample of X_A can be computed by using the inverse function with defined and simulated values of θ_A . The sample of X_D can be computed with the sample of X_A and any defined and simulated values of θ_B and θ_C . The parameter vector of the samples of X_A and X_D can be estimated by the maximum likelihood method (ML). The goodness-of-fit of $F_A(x_A, \hat{\theta}_A)$ and $F_D(x_D, \hat{\theta}_D)$ can be evaluated with the real probability which is the simulated and known sample of U . The validation values of goodness-of-fit are defined here with Δ_{\max} and Δ^2

$$\Delta_{\max} = \max_{i=1..n} |u_i - F(x_i, \hat{\theta})|,$$

Biased transformation and Goodness-of-Fit Test

$$\Delta^2 = \frac{\sum_{i=1}^n [u_j - F(x_i, \hat{\theta})]^2}{n}.$$

The definition of Δ_{\max} is comparable with the test value of the Kolmogorov-Smirnov test of fit. The definition of Δ^2 is similar to the mean squared error of parameter estimation. The values Δ_{\max} and Δ^2 are random variables as well. If the assumption is right for the defined range of parameters of transformation θ_B and θ_C , then the distribution of Δ_{\max} and Δ^2 of X_A is equal or very similar to the distribution of Δ_{\max} and Δ^2 of X_D ; and Δ_{\max} and Δ^2 should be strongly correlated for X_A and X_D .

Simulations have been carried out in a broad range of parameters and sample sizes to validate the assumption. θ_A and θ_C are simulated in a defined range. θ_B have to be defined with $\theta_B = \theta_A + \Delta_\theta$ with the simulated θ_A and a simulated Δ_θ in a defined range of Δ_θ which is rather narrow. If the difference Δ_θ is too large the approximations of $F(x)$ for the normal, beta or gamma distribution could result in $F_B(x_A) = 1$ or $F_B(x_A) = 0$. If $F_C^{-1}(1)$ or $F_C^{-1}(0)$ results in infinity or negative infinity, the transformation has to be cancelled.

The empirical distributions of Δ_{\max} and Δ^2 are plotted here for graphical comparisons with the position

$$\hat{P}_i = i/(n+1).$$

The Kolmogorov-Smirnov test for homogeneity (Birnbaum and Hall 1960) is used to validate that the distributions of Δ_{\max} of X_A and Δ_{\max} of X_D are the same population and the distributions of Δ^2 of X_A and Δ^2 of X_D are the same population. The test value D_{n_1, n_2} of the test of homogeneity of the two the samples of Δ^2 is written

$$D_{n_1, n_2} = \max_{\Delta^2} \left\{ \left| \hat{F}_1(\Delta^2) - \hat{F}_2(\Delta^2) \right| \right\}$$

and for Δ_{\max} in the same manner. The test values should be smaller than the critical value for a significance of $\alpha = 10\%$ or higher. H_0 (the same population) is not rejected.

Three variants for the sample size n for X_A and X_D were simulated with $n = 10, 100$ and 1000 . For each variant of sample size and distribution type of F_A and F_D , 1000 simulations were carried out. The ML estimation of the parameters was carried out for beta and gamma distributions with an exactness of $10^{-4}\%$. For the different variants, the following ranges of parameters were defined:

Biased transformation and Goodness-of-Fit Test

- If X_A is gamma distributed, then the parameters were simulated with $\alpha_A=1.0...3.5$ and $\lambda_A=0.5...4.0$ uniformly distributed. The parameters of F_B were simulated with $\alpha_B=\alpha_A+\Delta_\alpha$, $\Delta_\alpha=-0.2...0.2$ uniformly distributed and $\lambda_B=\lambda_A+\Delta_\lambda$, $\Delta_\lambda=-0.2...0.2$ uniformly distributed.
- If X_A is beta distributed the parameters were simulated with $\alpha_A=0.75...4.0$ and $\beta_A=0.75...4.0$ uniformly distributed. The parameters of F_B were simulated with $\alpha_2=\alpha_A+\Delta_\alpha$, $\Delta_\alpha=-0.25...0.25$ uniformly distributed and $\beta_B=\beta_A+\Delta_\beta$, $\Delta_\beta=-0.25...0.25$ uniformly distributed.
- If X_A is normally distributed, the parameters were simulated with $\mu_A=-0.5...0.5$ and $\sigma_A=0.75...1.25$ uniformly distributed. The parameters of F_B were simulated with $\mu_B=\mu_A+\Delta_\mu$, $\Delta_\mu=-0.25...0.25$ uniformly distributed and $\sigma_B=\sigma_A+\Delta_\sigma$, $\Delta_\sigma=-0.25...0.25$ uniformly distributed.

The parameters of F_C were simulated independent of θ_A in the following ranges:

- If $F_C(x)$ is the CDF of a gamma distribution, the parameter α_C was simulated in the range $\alpha_C=0.7...4.0$ uniformly distributed and the parameter λ_C was simulated in the range $\lambda_C=0.7...4.0$ uniformly distributed.
- If $F_C(x)$ is the CDF of a beta distribution, the parameter α_C was simulated in the range $\alpha_C=0.7...4.0$ uniformly distributed and the parameter β_C was simulated in the range $\beta_C=0.7...4.0$ uniformly distributed.
- If $F_C(x)$ is the CDF of a normal distribution, the parameter μ_C was simulated in the range $\alpha_C=-1...1$ uniformly distributed and the parameter σ_C was simulated in the range $\sigma_C=0.5...1$ uniformly distributed.

The important point of the BT is the bias function $F_B(F_A)$. The dimension of the bias functions of the simulations are shown by the extreme of $F_B(x_A)$ for the simulated parameters θ_A and θ_B and the mean values of the absolute difference $|F_B(x_A)-F_A(x_A)|$ for the simulated parameters θ_A and θ_B as function of F_A in Fig. 1. The bias of BT of simulated samples is large. The good results of the simulation can not be explained with a small bias in BT. The result of the simulation is a validation of the assumption. It can be seen in Figures 2a and b with the distributions of Δ_{\max} and Δ^2 for the case that X_A is normally distributed. The plotted values of Δ_{\max} and Δ^2 of the different variants of F_C and X_D result in very similar curves. It is the same for the case that X_A is gamma or beta distributed. The test values of the

Biased transformation and Goodness-of-Fit Test

1
2
3 Kolmogorov-Smirnov test of homogeneity for two samples are in Tab. 1. All values $D_{n_1; n_2}$ are much
4
5 smaller than the critical value of 0.048 for the level of significance of 10% for the 1000 simulations
6
7 (number of simulations = sample size here). H_0 – the same population – is never rejected. The BT
8
9 works for the normal, beta and gamma distribution as a good approximation in the range of the
10
11 simulations. The sample size n does not influence the result. The correlation of the validation values of
12
13 goodness-of-fit is very high and in a small range around $r=0.98$. It is clear in Figures 2a and b that the
14
15 transformation for X_A to X_D as normal distribution is exactly as shown in proof in the appendix.

16
17
18 **If a beta or gamma distribution is transformed into a normal distribution with transformation**
19
20 **parameters of F_B which differ not much from the parameters F_A of X_A , than the fit of an**
21
22 **estimated normal distribution F_D for X_D should not be influenced by the parameters of F_B and F_C**
23
24 **according to the simulation results. The parameters of F_C should not influence the fit of X_D**
25
26 **according to the simulation results.**
27
28

30 31 **4 The Application of the BT for the Anderson-Darling Test** 32 **for the Gamma Distribution** 33

34 There are EDF tests of goodness-of-fit with the hypothesis H_0 that the sample is from the assumed
35
36 distribution with the estimated parameters and with the defined level of significance. The level of
37
38 significance is the share of rejections although the assumption H_0 is correct. The transformed variable
39
40 X_D has almost the equal quality of goodness-of-fit as the variable X_A as shown earlier. The results of
41
42 the goodness-of-fit test should almost be the same for X_D and likewise for X_A . The different shares of
43
44 rejection of the test hypothesis H_0 , even if H_0 is right – an error of the first kind - should be almost
45
46 equal. The power should be almost equal likewise. The power is the share of rejection if H_0 is false.
47
48 The error of second kind is the share of acceptance of a false hypothesis H_0 .

49
50
51 The power function is the probability of rejection dependent on estimated parameter. The parameters
52
53 of CDF are the independent variables in the power function. The probability of rejection should
54
55 increase strong if the absolute difference between the parameters in the power function and the
56
57 estimated parameters increase. This power function is not good in the test with the BT because the
58
59 basics of the BT. But the test with the BT may be used only for the ML-estimated parameters.
60
Furthermore, the power of the test can be very high independent of the poor power function. It is

Biased transformation and Goodness-of-Fit Test

tested with BT if the sample is from a family of distributions. This approach is not new. It has been already introduced by Quesenberry (1986, equation 6.2).

The approach is used concretely for the Anderson-Darling test (Anderson and Darling 1952, Stephens 1974). The test value is A^2 with

$$A^2 = -n - 1/n \sum_{j=1}^n \left\{ (2j-1) \left[\ln(F(x_j, \hat{\theta})) + \ln(1 - F(x_{n+1-j}, \hat{\theta})) \right] \right\}.$$

The test value has to be adopted for normal distributions with

$$A^{*2} = A^2 \left(1 + 0.75/n + 2.25/n^2 \right).$$

The critical values are 1.035 for the level of significance of 1%, 0.752 for the level of significance of 5% and 0.631 for the level of significance of 10%. Only the Anderson-Darling test is used from the EDF-tests because the Anderson-Darling Test has the best behaviour of the tests for normality according Landry and Leparge (1992). But the approach probably works for the Kolmogorov-Smirnov test and the Cramér-von-Mises test for the goodness-of-fit likewise.

The procedure for the test of goodness-of-fit for the gamma distributed X_A by using the BT is the following:

- ML estimation of $\hat{\theta}_A$.
- Using this estimation of $\hat{\theta}_A$ for θ_B .
- Definition of θ_C with F_C as the CDF of a normal distribution with $\mu=0$ and $\sigma=1$.
- Computing the sample of X_D by the BT according equation 2.
- ML estimation of $\hat{\theta}_D$ of the sample of X_D , assumed as normal distribution – H_0 .
- Computing of the A^2 for the sample of X_D and test.
- Rejection of H_0 at the defined level of significance if $A^{*2} > A^2_{\text{critical}}$, otherwise acceptance of H_0 .

The assumed gamma distribution of X_A is rejected with the rejection of the assumed normal distribution of X_D .

The results of the test of the samples of X_D from simulated samples of X_A can be compared with results of the Anderson-Darling test for estimation of the gamma distribution of X_A according to Stephens (1986, critical values form Tab. 4.21). Stephens' test procedure is applied for the case that all parameters are estimated except the location parameter. There is no location parameter in the gamma

Biased transformation and Goodness-of-Fit Test

distribution as formulated above, the location parameter is known and has the value 0 in this paper. Interpolation functions (from regression analysis of values given by Stephens, 1986, Tab. 4.21) are used here for the given critical value for the significance levels of 1%, 5% and 10% with

$$A_{critical1\%}^2 = 1.035 + 0.057\hat{\alpha}^{-1.093},$$

$$A_{critical5\%}^2 = 0.752 + 0.034\hat{\alpha}^{-1.114},$$

$$A_{critical10\%}^2 = 0.631 + 0.0264\hat{\alpha}^{-1.110}.$$

The parameter vector θ_A for X_A – assumed to be gamma distributed has to be estimated by the ML-method. This is defined in the procedure of Stephens, and likewise the ML estimation is defined in the procedure with the BT.

Different constellations of distributions and parameters for X_A are simulated and the estimations are tested. 10000 simulations were carried out for each constellation. X_A was assumed as gamma distribution in each constellation. X_A was simulated in each constellation by a simulation of a sample of the uniformly distributed random variable U and the using of the inverse function. Furthermore, a normal distributed sample was also calculated from the sample of U with $\mu=0$ and $\sigma=1$. The parameters were estimated for this by using the ML method and the Anderson-Darling test is applied as well. The share of rejection for this procedure is the control value for the simulations.

The constellation with gamma distributed X_A was not the only simulation. The logarithmic normal distribution for X_A was simulated as well (the parameters are for $\ln(X_A)$). Furthermore the Burr distribution Type XII and the Pareto distribution were applied with

$$F(x) = 1 - \left[\eta / (\eta - x^\tau) \right]^\lambda, \quad x \geq 0, \eta > 0, \tau > 0, \lambda > 0,$$

$$F(x) = (x+1)^{-\alpha}, \quad x \geq 0, \alpha > 0.$$

These “false” distributions are used to validate the power of the test. The resulting shares of rejections for the different levels of significance are shown in Tables 2 to 4 for the sample sizes of X_A of $n=25$, 50 and 100. The results of the Anderson-Darling test for a gamma distributed random variables are in the column “Classical Test”. The real distribution of X_A and their parameters are listed in the columns labelled as such. It is clear that the Anderson-Darling test with the BT works as well as the Anderson-

Biased transformation and Goodness-of-Fit Test

1
2
3 Darling test with the procedure and the critical values of Stephens (1986) if the hypothesis H_0 is right.
4
5 The shares of rejections are very similar and nearly all are in the 90% range of the estimated share of a
6
7 Bernoulli distribution. The ranges are 0.8%-1.1% for the level of 1%, 4.6%-5.3% for the level of 5%
8
9 and 9.5%-10.4% for the level of 10%. The shares of rejections are out of range in the case that $n=100$
10
11 and the level of significance are defined with 5%. But in this constellation, the control value is also out
12
13 of range.
14

15
16 The results of the classical test and the test with BT are very similar for the cases that X_A is not gamma
17
18 distributed. The differences between the classical test and the test with BT in the cases that X_A are
19
20 gamma distributed are small but partly significant. The new procedure with the BT reject more false
21
22 assumptions if the sample size is not large ($n=25$). The procedure according Stephens has a bit more
23
24 power in the case of larger sample sizes ($n \geq 50$). Independent of this fact - the new procedure works
25
26 well and the power of the test is high. Both test procedures are not very powerful in the case that the
27
28 real distribution is GPD and γ is near 0. The reason for this phenomena is that the GPD for $\gamma=0$
29
30 (Gumbel case) is an exponential distribution which is a special case of gamma distribution with $\alpha=1$.
31
32 The GPDs with γ near 0 and the gamma distributions with α near 1 are similar, no test can be very
33
34 power full in this constellation.
35
36
37
38

39 40 **5 The Application of the BT for the Anderson-Darling Test** 41 **for the Beta Distribution** 42

43
44 The only test specified for the beta distribution which was found in publications is an approach of Li
45
46 and Papadopoulos (2002). The moments are used in this approach for goodness-of-fit tests for
47
48 different distributions. But there is a mistake in the chapter for the beta distribution (a moment is
49
50 divided by itself), and the power of the test is not so high for an exponential distribution. For example,
51
52 7-8% were rejected at a significance level of 5% in the case that H_0 was correct (Li and Papadopoulos,
53
54 2002, Tab. 1). This approach is not considered in this paper. Further tests or critical values for test
55
56 specified for the beta distribution were not found in publications. In this place, the BT can be used for
57
58 a first EDF test of goodness-of-fit for the beta distribution specified for it. The procedure includes the
59
60 following steps:

- ML estimation of $\hat{\theta}_A$ under the assumption of beta distribution for X_A .

Biased transformation and Goodness-of-Fit Test

- 1
- 2
- 3
- 4 - Using this estimation of $\hat{\theta}_A$ for θ_B .
- 5
- 6 - Definition of θ_C with F_C as the CDF of a normal distribution with $\mu=0$ and $\sigma=1$.
- 7
- 8 - Computing the sample of X_D by the BT according equation 2.
- 9
- 10 - ML estimation of $\hat{\theta}_D$ of the sample of X_D , assumed as normal distribution – H_0 .
- 11
- 12 - Computing of the A^2 for the sample of X_D and test.
- 13
- 14
- 15 - Rejection of H_0 at the defined level of significance if $A^{*2} > A^2_{critical}$, otherwise acceptance of H_0 .
- 16

17 The assumption that X_A is beta distributed is rejected with the rejection of H_0 that X_D is normally
 18 distributed. The power function of this test has a poor quality even if the test works well. The test
 19 procedure is evaluated by simulations. Samples of beta distribution with variants of parameters were
 20 simulated (Tables 5-7). And distributions were simulated for X_A which were not beta distributions.
 21 These were the log-normal distributions, the gamma distribution and the normal distribution. The log-
 22 normal and the gamma distributions were truncated at $x=F^{-1}(0.99)$ and scaled, so that $x_{max}=1$. The
 23 normal distribution was truncated with a lower bound $x=F^{-1}(0.01)$ and an upper bound $x=F^{-1}(0.99)$ and
 24 was located and scaled than so that $0 \leq x \leq 1$.

25 The samples were simulated for the size of $n=10, 25, 50$ and 100 . The considered levels of
 26 significance are 1%, 5% and 10%. Furthermore, the χ^2 test and the Kolomogorov-Smirnov test were
 27 applied for the sample sizes of $n=50$ and $n=100$ to compare the power of this test with the Anderson-
 28 Darling test with the BT. The samples were divided in groups with 10 observations for the χ^2 test. The
 29 bound between the groups was selected with the mean of the largest value of the smaller group and the
 30 minimum observation of the larger group. The variant of the Kolomogorov-Smirnov for a full
 31 specified distribution hypothesis was applied for the sample sizes of $n=50$ and $n=100$ likewise. The
 32 critical values of the χ^2 test and the Kolomogorov-Smirnov test are taken from Rinne (2003, Chapter
 33 3.4.5).

34 The shares of rejection from the 10000 simulations of each variant of real distribution are shown in
 35 Tables 5 through 7. The test procedure works well. The share of rejections for the case that H_0 is true
 36 is almost equal to the defined level of significance. But some values in these cases are outside the 90%
 37 range for the estimation of the share of a Bernoulli distribution. The randomness of the simulation is
 38 reasonable for this because the control value is also often out of this range. The control value is again

Biased transformation and Goodness-of-Fit Test

1
2
3 the share of rejections for the real normal distribution with estimated parameters for a normal
4 distribution and the same underlying sample of U used for the other real distributions as well. The
5 Anderson-Darling test for the beta distribution by using the BT has more power than the χ^2 test or the
6 Kolomogorov-Smirnov test as shown in Tables 6 and 7. The χ^2 test rejects less than the defined level
7 of significance of beta distribution than the new procedure for the Anderson-Darling test in the cases
8 that H_0 is false; and the χ^2 test rejects more than the defined level of significance of beta distribution.
9 The error of the first kind is higher for the χ^2 test than for the new procedure. The Kolomogorov-
10 Smirnov test does not work; the power is very small.
11
12
13
14
15
16
17
18
19
20
21

22 **6 Practical application of tests for beta distribution**

23
24 The beta distribution is used for modeling of meteorological random variables. Yao (1974) modeled
25 surface relative humidity of U.S. weather stations by using the beta distribution. Different data sets
26 were analysed for example daily observations for one month. Parameters were estimated by the
27 moment method. The goodness-of-fit was evaluated by using χ^2 -test. The number of observations per
28 classes was not limited with $n_i \geq 10$ contrary to recommendations (Rinne, 2008, chapter 3.4.5.1). The fit
29 is good according to this test results (Yao, 1974, Tab. 1).
30
31
32
33
34
35
36

37 The daily cloud duration was modeled by Chia and Hutchinson (1991) with beta distributions for
38 different places in Australia. The goodness-of-fit was evaluated by a Kolmogorov-Smirnov test for a
39 full specified distribution. The beta distribution can be used for most data according test results (Cia
40 and Hutchinson, 1990, Tab. 3).
41
42
43
44
45

46 The beta distribution was applied furthermore for data of relative sunshine duration of Malaysia by
47 Sulaiman et al. (1999). The goodness-of-fit was evaluated by the Kolmogorov-Smirnov test again. The
48 fit is good for most of the data (Sulaiman et al., 1999, Tab. 2).
49
50
51
52

53 Neither the χ^2 -test nor the Kolmogorov-Smirnov test has a good performance for beta distribution as
54 seen in Tables 6 and 7. Especially the Kolmogorov-Smirnov test has very low power. But till now
55 there existed no better test. A practical example of meteorological data demonstrates the improvement
56 of new test procedure for application of beta distribution. The data of Haarweg Wageningen weather
57 station of Wageningen University (Netherlands) of July 2005 are analysed. The data set includes
58
59
60

Biased transformation and Goodness-of-Fit Test

relative humidity. The relative sunshine duration is calculated by using sunshine minutes of data set and time between sunrise and sunset. The time between sunrise and sunset is assumed with linear function with 998.4min for 1st of July and 931.1min for 31st of July. Parameters of beta distribution are estimated with the ML method. The estimation is $\hat{\alpha}=7.47$ and $\hat{\beta}=4.6255$ for relative humidity. The critical value for the Kolmogorov-Smirnov test is 0.23788 for $n=31$ and 5% significance and is not overstepped with 0.1253. H_0 - beta distributed sample - is accepted. The χ^2 -test is applied with four classes with $n_i=7$ and 8 (although n_i should be ≥ 10). The critical value is 3.841 (5%, one grade of freedom) and is not overstepped by the test value of 2.686. Contrary to this the Anderson-Darling test value $A^{*2}=1.433$ and is much larger the $A_{crit}^2=0.753$ (5% significance). The sample is not beta distributed according to the Anderson-Darling test. The parameters of normal distribution of X_D are $\hat{\mu}=-0.0005$ and $\hat{\sigma}=1.0164$. The results for data of relative sunshine duration per day are similar. Sample size is $n=31$ again. Parameters are $\hat{\alpha}=1.5536$ and $\hat{\beta}=0.7480$. The value of the Kolmogorov-Smirnov test is 0.1706 and doesn't overstep critical value. The value of χ^2 -test is 3.9937. This oversteps the critical value a bit. H_0 would be rejected according to the χ^2 -test but not according to the Kolmogorov-Smirnov test. The Anderson-Darling test rejects the hypothesis of beta distributed data very clearly with $A^{*2}=2.059$. The parameters of the normal distribution of X_D are $\hat{\mu}=-0.0116$ and $\hat{\sigma}=1.0057$. The estimated CDF and observed data are shown in Fig. 3. The fit is not good.

The example shows that the introduced procedure of the Anderson-Darling test for beta distribution by using BT is an improvement of applied statistic. It seems that many data are assumed likely as beta distributed in pasted analysis although they are not beta distributed.

7 Summary and Discussion

The BT is introduced and can be used exactly for each of the GEV, the GPD and the normal distribution. The BT works as an approximation for the gamma, beta and normal distribution as a family of distributions. Because of this the Anderson-Darling Test for normal distributions can be used as a goodness-of-fit test for the gamma and the beta distribution. The new test procedure works with similar power as the classical procedure of Stephens. The power of the new procedure is a bit higher

Biased transformation and Goodness-of-Fit Test

1
2
3 for smaller sample sizes. The classical procedure has a bit more power for larger sample sizes. The
4 same test procedure in principle is used for a first goodness-of-fit test specified for beta distribution.
5
6 This is the first test specified for the beta distribution. The power function of the new test is poor
7 because of the basics of the BT. And it is defined in the test procedure that the ML estimated
8 parameters of assumed the gamma or beta distribution has to be used. The method of parameter
9 estimation is defined in the procedures of Stephens. The general question is: Is it necessary to have a
10 good power function for the whole range of parameters for a powerful test if the estimation method is
11 defined in the test procedure? A good power function is not necessary according to the results of this
12 paper. The Anderson-Darling test for the gamma distribution with the new procedure has a similar
13 power as the classical Anderson-Darling test for the gamma distribution. And the power function of
14 the new procedure is very poor compared with the function of the classical test. The poor power
15 function can result only in wrong test results, if the test procedure is not used right and the parameter
16 vector θ_A is not estimated correct with the ML method.
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 If the new test does not reject the hypothesis, the user can assume that the ML estimation of the
32 parameters leads to an estimated distribution with similar goodness-of-fit as for a sample of normal
33 distribution with the equal underlying sample of U as for the gamma or beta distribution. It was shown
34 in the validation of the BT that the goodness-of-fit of a beta, gamma or normal distributed sample
35 from the same underlying sample of U is almost equal (Figures 2 c and d).
36
37
38
39
40
41

42 The CDF of the normal, gamma and beta distributions are computed by using approximations from
43 different software libraries (Excel was used and the library of www.extremeoctimizations.com). These
44 approximations work well in the common ranges of parameters and percentiles. But if, for example, α
45 of the gamma distribution is small and the percentile is as well, the approximation is wrong. The BT of
46 the normal, gamma and beta distribution cannot work better than the approximation of the CDF of
47 these distributions. This is a reason why larger simulations were not carried out for the validation of
48 the BT for normal, gamma and beta distribution. The probability that the approximation of the CDF
49 does not work in a variant of parameters and sample of U increases with the number of simulations.
50
51
52
53
54
55
56
57
58
59
60 The failure of the approximation is avoided by a limitation of the simulations. And more simulations

Biased transformation and Goodness-of-Fit Test

are not necessary, because the distributions of the values of goodness-of-fit in the validation of the BT were tested with knowledge of the simulation size.

The other EDF-tests specified for the normal distribution could be used in the test procedure with the BT too. The power of such tests could be evaluated in further works. The Anderson-Darling test has the better quality in the test of normality (Landry and Lepage, 1992). That is why only this EDF-test is applied in this paper.

The practical relevance of the new test for beta distributions was demonstrated successfully in metrological data.

The BT is interesting irrespective of the introduced application for the test of the gamma and beta distribution. Perhaps the BT will become a criterion for the classification of distributions or the BT can perhaps be used for estimations of the GPD or GEV or for their tests.

The possibility of BT to estimate the PDF or CDF of a beta distribution by using the normal distribution should be researched in the future.

8 Acknowledgement

The author would like to thank the Federal Ministry for Education and Science of Germany for the financial support of the FloodRisk project in the EraNet Crue initiative; the research for the paper was partly carried out within this project. **Furthermore I would like to express my gratitude to Michael A. Stephens for the helpful communication.**

9 References

- Anderson, T.W., Darling, D. A. (1952). Asymptotic theory of certain "Goodness-of-fit" criteria based on stochastic processes. *Ann. Math. Statist.* 23: 193-212.
- Anderson, T.W., Darling, D. A. (1954). A test for goodness-of-fit. *J. Amer. Statist. Assoc.* 49: 300-310**
- Birnbaum, Z.W., Hall, R.A. (1960). Small Sample Distributions for Multi-Sample Statistics of Smirnov Type. *Ann. Math. Statist.* 31: 710-720
- Bowman, K.O., Shenton, L.R. (1986). Moment ($\sqrt{b_1, b_2}$) Techniques. In: D'Augustino, R.B., Stephens, M.A. ed., *Goodness-of-Fit Techniques. Statistics: textbooks and monographs, Vol. 68: 279-327, Marcel Dekker: New York*
- Box, G.E.P.; Cox, D.R. (1964). An Analysis of Transformations. *J. R. Statist. Soc. B*, 26: 211-246

Biased transformation and Goodness-of-Fit Test

1
2
3 **Carley, H., Taylor, M.D. (2000). A new proof of Sklar's theorem. Conference on Distributions with given**
4 **marginals, Barcelona, July 2000**

5
6
7 Chia, E., Hutchinson, M.F. (1991). The beta distribution as a probability model for daily cloud duration.
8
9 Agriculture and Forest Meteorology, 56, 195-208

10
11 **Johnson, N.L., Kotz, S., Balakrishnan, N. (1994). Continuous Univariate Distributions- Volume 1. 2nd Ed.**
12 **Wiley: New York**

13
14
15 **Johnson, N.L., Kotz, S., Balakrishnan, N. (1995). Continuous Univariate Distributions- Volume 1. 2nd Ed.**
16 **Wiley: New York**

17
18
19 **Kolmogorov, A.N. (1933). Sulla determinazione empirica di una legge di distribuzione. Giorn. Ist. Attuari.**
20 **4: 83-91**

21
22
23 **Kuiper, N.H. (1960). Tests concerning random points on a circle. Proc. Konink. Neder. Akad. Van.**
24 **Wetenschappen, A 63: 38-47.**

25
26
27 Landry, L., Lepage, Y. (1992). Empirical Behavior of some tests for Normality. Commun. Statist. B – Computa.
28
29 Simula. 21: 971-999

30
31 Li, G., Papadopoulos, A. (2002). A note on Goodness of fit test using moments. STATISTICA, 62, n. 1: 72-86

32
33 Moore, D.S. (1986). Tests of Chi-Squared Type. In: D'Augustino, R.B., Stephens, M.A. ed., Goodness-of-Fit
34
35 Techniques. Statistics: textbooks and monographs, Vol. 68: 63-95, Marcel Dekker: New York

36
37 Quesenberry, C.P. (1986). Some transformation methods in Goodness of fit. In: D'Augustino, R.B., Stephens,
38
39 M.A. ed., Goodness-of-Fit Techniques. Statistics: textbooks and monographs, Vol. 68: 235-278, Marcel
40
41 Dekker, New York

42 Rinne, H. (2003). Taschenbuch der Statistik (German; Handbook of the statistic). Harri Deutsch, Frankfurt

43 Sklar, M. (1959). Fonctions de repartition à N dimensions et leurs marges. Publ. Inst. Stat. Paris, 8: 229-231

44
45 Stephens, M.A. (1974). EDF Statistics for Goodness of Fit and Some Comparisons. J. Am. Statist. Assoc. 69, No
46
47 347: 730- 737

48
49
50 **Stephens, M.A. (1976). Asymptotic results for Goodness-of-Fit statistics with unknown parameters. The**
51 **Annals of Statistics 4, no2: 357-369**

52
53
54 Stephens, M.A. (1986). Tests based on EDF statistics. In: D'Augustino, R.B., Stephens, M.A. ed., Goodness-of-
55
56 Fit Techniques. Statistics: textbooks and monographs, Vol. 68: 97-194, Marcel Dekker: New York

57
58 Sulaiman, M.Y., Oo, H.W.M., Wahab, M.A., Zakaria, A. (1999). Application of Beta distribution model to
59
60 Malaysian sunshine data. Renewable Energy 18: 573-579

Biased transformation and Goodness-of-Fit Test

Yao, A.Y.M. (1974). A Statistical Model for the Surface relative humidity. Journal of Applied Meteorology. Vol. 13: 17-21

Wageningen University, Netherlands, Meteorology and Air Quality section (2009, download). Weather station Haarweg Wageningen. Data: <http://www.met.wau.nl/>

Watson, G.S. (1961). Goodness-of-fit tests on a circle. *Biometrika* 48: 109-114.

10 Appendix

The BT for the GEV

The GEV is formulated for the Gumbel case

$$F(x) = \exp\{-\exp[-(x - \mu)/\sigma]\} \quad (6)$$

and is formulated for the Fréchet and Weibull case with

$$F(x) = \exp\left\{-\left[1 + \gamma(x - \mu)/\sigma\right]^{-1/\gamma}\right\} \quad (7)$$

where the parameter is $\sigma > 0$. The lower bound is for the Fréchet case $\gamma > 0$

$$x \geq \mu - \sigma/\gamma.$$

The upper bound is for the Weibull case with $\gamma < 0$

$$x \leq \mu - \sigma/\gamma.$$

Theorem A1: If X_A is GEV-Gumbel distributed and F_B is the CDF of a GED-Gumbel and F_C is the CDF of a GEV-Fréchet/Weibull with any parameters, then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GEV-Fréchet/Weibull distributed with the parameters

$$\gamma_D = \gamma_C \sigma_A / \sigma_B, \quad (8)$$

$$\sigma_D = \sigma_A \sigma_C \exp[\gamma_C(\mu_A - \mu_B)/\sigma_B] / \sigma_B, \quad (9)$$

$$\mu_D = \sigma_C \exp[\gamma_C(\mu_A - \mu_B)/\sigma_B] / \gamma_C - \sigma_C / \gamma_C + \mu_C \quad (10)$$

and the transformation function $x_D(x_A)$ is

$$x_D = \{\sigma_C \exp[\gamma_C(x_A - \mu_B)/\sigma_B] - \sigma_C + \gamma_C \mu_C\} / \gamma_C. \quad (11)$$

Proof. If equations 8-11 are used in $F_D(x_D)$ according to equation 7, then according to equation 6 the CDF $F_A(x_A)$ can be formulated after simplifications so that equation 3 is valid. \square

This transformation works likewise the other way round if the bounds of X_D and X_C are the same with

$$\mu_C - \sigma_C / \gamma_C = \mu_D - \sigma_D / \gamma_D.$$

and γ_C and γ_D have the same sign.

Theorem A2: If X_A is GEV-Gumbel distributed and F_B is the CDF of a GEV-Gumbel and F_C is the CDF of a GEV-Gumbel with any parameters, then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GEV-Gumbel distributed with the parameters

$$\sigma_D = \sigma_A \sigma_C / \sigma_B. \quad (12)$$

$$\mu_D = (-\mu_B \sigma_C + \mu_C \sigma_B + \mu_A \sigma_C) / \sigma_B \quad (13)$$

and the transformation function $x_D(x_A)$ is

$$x_D = (\sigma_B \mu_C + \sigma_C x_A - \mu_B \sigma_C) / \sigma_B. \quad (14)$$

Biased transformation and Goodness-of-Fit Test

Proof: If equations 12-14 are used in $F_D(x_D)$ according to equation 6, then according to equation 6 the function $F_A(x_A)$ can be formulated after simplifications so that equation 3 is valid. \square

Theorem A3: If X_A is GEV-Fréchet/Weibull distributed and F_B is the CDF of a GEV-Fréchet/Weibull and F_C is the CDF of a GEV-Fréchet/Weibull with any parameters but with

$$\sigma_A / \gamma_A - \mu_A = \sigma_B / \gamma_B - \mu_B \tag{15}$$

and γ_A and γ_B have the same sign then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GEV-Fréchet/Weibull distributed with the parameters

$$\gamma_D = \gamma_A \gamma_C / \gamma_B, \tag{16}$$

$$\sigma_D = [\sigma_A \gamma_B / (\gamma_A \sigma_B)]^{\gamma_C / \gamma_B} \sigma_C \gamma_A / \gamma_B, \tag{17}$$

$$\mu_D = \mu_C - \sigma_C / \gamma_C + [\sigma_A \gamma_B / (\gamma_A \sigma_B)]^{\gamma_C / \gamma_B} \sigma_C / \gamma_C \tag{18}$$

and the transformation function $x_D(x_A)$ is

$$x_D = \left\{ \sigma_C [1 + \gamma_B (x_A - \mu_B) / \sigma_B]^{\gamma_C / \gamma_B} - \sigma_C + \gamma_C \mu_C \right\} / \gamma_C. \tag{19}$$

Proof: If equations 16-19 are used in $F_D(x_D)$ according to equation 7, and equation 15 is considered, then according equation 7 the function $F_A(x_A)$ can be formulated after some simplifications so that equation 3 is valid. \square

The BT for the GPD

The GPD is formulated for the Gumbel case

$$F(x) = 1 - \exp(-x / \sigma) \tag{20}$$

and for the Fréchet and Weibull case formulated with

$$F(x) = 1 - (1 + \gamma x / \sigma)^{-1/\gamma}. \tag{21}$$

Generally $x \geq 0$ und $\sigma > 0$. There is an upper bound for the Weibull case $\gamma < 0$ with

$$x \leq -\sigma / \gamma.$$

Theorem A4. If X_A is GPD-Gumbel distributed and F_B is the CDF of a GPD-Gumbel and F_C is the CDF of a GPD-Gumbel with any parameters, then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GPD-Gumbel distributed with the parameters

$$\sigma_D = \sigma_A \sigma_C / \sigma_B \tag{22}$$

and the transformation function $x_D(x_A)$ is

$$x_D = x_A \sigma_C / \sigma_B. \tag{23}$$

Proof: If equations 22-23 are used in $F_D(x_D)$ according equation 20, then according to equation 20 the function $F_A(x_A)$ can be formulated after some simplifications so that equation 3 is valid. \square

Theorem A5: If X_A is GPD-Fréchet/Weibull distributed and F_B is the CDF of a GPD-Fréchet/Weibull and F_C is the CDF of a GPD-Fréchet/Weibull with any parameters but with

$$\sigma_A / \gamma_A = \sigma_B / \gamma_B \tag{24}$$

then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GPD-Fréchet/Weibull distributed with the parameters

$$\gamma_D = \gamma_A \gamma_C / \gamma_B, \tag{25}$$

$$\sigma_D = \gamma_A \sigma_C / \gamma_B \tag{26}$$

and the transformation function $x_D(x_A)$ is

$$x_D = \sigma_C \left\{ (1 + \gamma_B x_A / \sigma_B)^{\gamma_C / \gamma_B} - 1 \right\} / \gamma_C. \tag{27}$$

Biased transformation and Goodness-of-Fit Test

Proof: If equations 25-27 are used in $F_D(x_D)$ according to equation 21 under consideration of equation 24, then according to equation 21, the function $F_A(x_A)$ can be formulated after some simplifications so that equation 3 is valid. \square

The condition of equation 24 has to be also considered for the Fréchet case even this is not a real bound of X .

Theorem A6: If X_A is GPD-Gumbel distributed and F_B is the CDF of a GPD-Gumbel and F_C is the CDF of a GPD- Fréchet/Weibull with any parameters, then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is GPD- Fréchet/Weibull distributed with the parameters

$$\gamma_D = \gamma_C \sigma_A / \sigma_B, \quad (28)$$

$$\sigma_D = \sigma_A \sigma_C / \sigma_B \quad (29)$$

and the transformation function $x_D(x_A)$ is

$$x_A = \sigma_C \{ \exp(x_A \gamma_C / \sigma_B) - 1 \} / \gamma_C. \quad (30)$$

Proof: If equations 28-30 are used in $F_D(x_D)$ according to equation 20, then according to equation 21, the function $F_A(x_A)$ can be formulated after some simplifications so that equation 3 is valid. \square

It is possible to transform X_D to X_A with any $F_C(x)$ and $F_B(x)$. It has only to be ensured that

$$\sigma_C / \gamma_C = \sigma_D / \gamma_D.$$

The Normal Distribution

Theorem A7: If X_A is normally distributed and F_B and F_C are CDF of normal distributions with any parameters, then X_A can be transformed to X_D with $X_D = F_C^{-1}(F_B(X_A))$ and X_D is normally distributed with the parameters

$$\mu_D = \mu_C + \sigma_C (\mu_A - \mu_B) / \sigma_B, \quad (31)$$

$$\sigma_D = \sigma_A \sigma_C / \sigma_B \quad (32)$$

and the transformation function $x_D(x_A)$ is

$$x_D = \mu_C + \sigma_C (x_A - \mu_B) / \sigma_B \quad (33)$$

Proof: If equation 4 is taken and the PDF are replaced by adaptations of equation 5 it can be simplified

$$f_D(x_D) = \frac{\sigma_B}{\sigma_A \sigma_C} \frac{\exp\left\{-\frac{[x_A(x_D) - \mu_A]^2}{2\sigma_A^2}\right\} \exp\left\{-\frac{(x_D - \mu_C)^2}{2\sigma_C^2}\right\}}{\sqrt{2\pi} \exp\left\{-\frac{[x_A(x_D) - \mu_B]^2}{2\sigma_B^2}\right\}}.$$

With equation 32, for the replacing of σ_A it can be modified and written

$$f_D(x_D) = 1 / (\sigma_D \sqrt{2\pi}) \exp\left\{-\frac{[x_A(x_D) - \mu_A]^2 \sigma_C^2}{2\sigma_D^2 \sigma_B^2}\right\} \exp\left\{-\frac{(x_D - \mu_C)^2}{\sigma_C^2} + \frac{[x_A(x_D) - \mu_B]^2}{\sigma_B^2}\right\}^{0.5}.$$

Further $x_A(x_D)$ can be replaced by using equation 33, and it can be simplified and modified to

$$f_D(x_D) = 1 / (\sigma_D \sqrt{2\pi}) \exp\left\{-\frac{[\sigma_B / \sigma_C (x_D - \mu_C) + \mu_B - \mu_A]^2 \sigma_C^2}{2\sigma_D^2 \sigma_B^2}\right\}.$$

The parameter μ_D can be replaced by the using of equation 31. It can be written

$$f_D(x_D) = 1 / (\sigma_D \sqrt{2\pi}) \exp\left\{-\frac{[\sigma_B / \sigma_C (x_D - \mu_C) + \mu_B - \mu_D \sigma_B / \sigma_C + \mu_C \sigma_B / \sigma_C - \mu_B]^2 \sigma_C^2}{2\sigma_D^2 \sigma_B^2}\right\}$$

and simplified to

$$f_D(x_D) = 1 / (\sigma_D \sqrt{2\pi}) \exp\left\{-\frac{(x_D - \mu_D)^2}{2\sigma_D^2}\right\}$$

which is the PDF for X_4 according to equation 5. X_4 is normally distributed. \square

Figures and Tables

Biased transformation and Goodness-of-Fit Test




Figure 1: Bias of the BT for the simulation of $n=1000$ for the different types of F_A and F_B : a) mean of the absolute difference $|F_B(x_A) - F_A(x_A)|$, b) extremes of $F_B(x_A)$ in relation to $F_A(x_A)$




Figure 2: Results of the simulations for $n=1000$ with X_A as normally distributed and the different distribution types of X_D : a) distributions of Δ_{max} , b) distributions of Δ^2 , c) Δ_{max} of X_A to Δ_{max} of X_D , d) Δ^2 of X_A to Δ^2 of X_D

Biased transformation and Goodness-of-Fit Test




Figure 3: Applied beta distribution: a) relative sunshine duration, b) relative humidity

Biased transformation and Goodness-of-Fit Test

Table 1: Test values of the Kolomogorov-Smirnov test for homogeneity for the distributions of Δ_{\max} and Δ^2 for the different variants of sample size, distribution of X_A and distribution of X_D

Sample size	Distribution of X_A	Distribution of Value	Distribution of X_D		
			Gamma	Beta	Normal
n=10	Gamma	Δ_{\max}	0.028	0.030	0.008
		Δ^2	0.021	0.019	0.005
	Beta	Δ_{\max}	0.009	0.008	0.030
		Δ^2	0.027	0.026	0.005
	Normal	Δ_{\max}	0.030	0.031	0.001
		Δ^2	0.033	0.036	0.001
n=100	Gamma	Δ_{\max}	0.013	0.016	0.010
		Δ^2	0.018	0.017	0.010
	Beta	Δ_{\max}	0.015	0.018	0.010
		Δ^2	0.014	0.019	0.009
	Normal	Δ_{\max}	0.016	0.021	0.001
		Δ^2	0.018	0.016	0.001
n=1000	Gamma	Δ_{\max}	0.016	0.019	0.017
		Δ^2	0.016	0.017	0.015
	Beta	Δ_{\max}	0.022	0.020	0.016
		Δ^2	0.021	0.019	0.023
	Normal	Δ_{\max}	0.019	0.020	0.001
		Δ^2	0.016	0.026	0.001

Table 2: Percentage or rejection of H_0 of the classical Anderson-Darling test, the test with the BT for gamma distribution for different significance levels and n=25. The level of significance is written in percentage.

Real Distribution	Parameters	Classical Test			Test with the BT		
		1%	5%	10%	1%	5%	10%
LogNorm	$\mu=1, \sigma^2=0.25$	3.42	10.92	18.19	3.69	11.41	18.89
LogNorm	$\mu=1, \sigma^2=1$	13.26	27.97	38.08	13.43	27.98	38.42
BurrXII	$\eta=4, \tau=8, \lambda=1$	4.74	12.1	19.41	5.28	13.32	21.08
BurrXII	$\eta=1, \tau=1, \lambda=2$	14.62	27.49	36.67	13.95	27.2	36.46
Pareto	$\alpha=8$	1.52	6.63	12.02	1.68	6.93	12.68
Pareto	$\alpha=2$	14.62	27.49	36.67	13.95	27.2	36.46
GPD	$\sigma=1, \gamma=-1/16$	0.91	4.67	9.56	0.92	4.9	10.01
GPD	$\sigma=1, \gamma=-1/4$	1.71	7.68	14.06	1.73	7.57	13.73
Gamma	$\lambda=1, \alpha=0.5$	0.86	4.62	9.46	0.86	4.87	9.72
Gamma	$\lambda=1, \alpha=1$	0.87	4.84	9.58	0.97	5.09	10.05
Gamma	$\lambda=1, \alpha=2.5$	0.89	4.92	9.72	0.98	5.18	10.07
Gamma	$\lambda=1, \alpha=5$	0.89	4.89	9.69	0.97	5.18	10.14
Gamma	$\lambda=1, \alpha=10$	0.89	4.88	9.72	0.98	5.17	10.13
Control	$\mu=0, \sigma=0$	-	-	-	0.97	5.18	10.09

Biased transformation and Goodness-of-Fit Test

Table 3: Percentage or rejection of H_0 of the classical Anderson-Darling test, the test with the BT for gamma distribution for different significance levels and $n=50$. The level of significance is written in percentage.

Real Distribution	Parameters	Classical Test			Test with the BT		
		1%	5%	10%	1%	5%	10%
LogNormal	$\mu=1, \sigma^2=0.25$	6.68	18.23	27.29	6.82	18.26	27.51
LogNormal	$\mu=1, \sigma^2=1$	33.24	53.23	63.57	31.46	51.51	62.1
BurrXII	$\eta=4, \tau=8, \lambda=1$	7.74	18.36	26.82	8.5	19.77	28.22
BurrXII	$\eta=1, \tau=1, \lambda=2$	34.32	50.4	59.21	31.19	47.81	56.89
Pareto	$\alpha=8$	2.66	8.43	14.56	2.46	8.26	14.44
Pareto	$\alpha=2$	34.32	50.4	59.21	31.19	47.81	56.89
GPD	$\sigma=1, \gamma=-1/16$	0.98	4.95	9.75	1	4.84	9.71
GPD	$\sigma=1, \gamma=-1/4$	3.2	11.59	20.09	3.04	10.94	18.81
Gamma	$\lambda=1, \alpha=0.5$	0.96	4.67	9.51	0.95	4.64	9.61
Gamma	$\lambda=1, \alpha=1$	0.97	4.68	9.4	1.04	4.72	9.61
Gamma	$\lambda=1, \alpha=2.5$	0.97	4.7	9.51	1.06	4.83	9.65
Gamma	$\lambda=1, \alpha=5$	0.97	4.69	9.49	1.05	4.81	9.59
Gamma	$\lambda=1, \alpha=10$	1	4.67	9.46	1.05	4.84	9.57
Control	$\mu=0, \sigma=0$	-	-	-	1.03	4.83	9.54

Table 4: Percentage or rejection of H_0 of the classical Anderson-Darling test, the test with the BT for gamma distribution for different significance levels and $n=100$. The level of significance is written in percentage.

Real Distribution	Parameters	Classical Test			Test with the BT		
		1%	5%	10%	1%	5%	10%
LogNormal	$\mu=1, \sigma^2=0.25$	14.22	31.41	43.44	14.09	31.28	43.15
LogNormal	$\mu=1, \sigma^2=1$	66.84	83.64	89.61	63.46	81.46	88.09
BurrXII	$\eta=4, \tau=8, \lambda=1$	14.59	28.65	38.66	15.57	29.83	40.13
BurrXII	$\eta=1, \tau=1, \lambda=2$	63.73	78.29	84.17	58.14	74.91	81.56
Pareto	$\alpha=8$	3.91	11.31	18.46	3.45	10.68	17.82
Pareto	$\alpha=2$	63.73	78.29	84.17	58.14	74.91	81.56
GPD	$\sigma=1, \gamma=-1/16$	1.14	5.27	10.56	1.12	5.3	10.33
GPD	$\sigma=1, \gamma=-1/4$	7.7	22.29	33.52	6.68	20.13	31.11
Gamma	$\lambda=1, \alpha=0.5$	1.04	4.54	9.33	0.92	4.34	9.71
Gamma	$\lambda=1, \alpha=1$	0.98	4.4	9.64	0.96	4.47	9.73
Gamma	$\lambda=1, \alpha=2.5$	0.93	4.44	9.8	0.99	4.58	9.85
Gamma	$\lambda=1, \alpha=5$	0.94	4.48	9.8	1.01	4.63	9.84
Gamma	$\lambda=1, \alpha=10$	0.95	4.52	9.79	1	4.61	9.81
Control	$\mu=0, \sigma=0$	-	-	-	1.01	4.6	9.84

Biased transformation and Goodness-of-Fit Test

Table 5: Percentage of rejections of H_0 of the goodness-of-fit test for the assumed beta distributions for different levels of significance [%] for the sample size $n=10$ and 25 (*distribution is truncated, scaled and/or moved)

Real Distribution	Parameter	BT and Anderson-Darling Test for Normality for $n=10$			BT and Anderson-Darling Test for Normality for $n=25$		
		1%	5%	10%	1%	5%	10%
Beta	$\alpha=0.5, \beta=0.5$	1.00	5.33	10.76	0.82	4.43	9.31
Beta	$\alpha=1, \beta=1$	1.04	5.51	10.73	0.86	4.50	9.38
Beta	$\alpha=2, \beta=2$	1.04	5.57	10.64	0.83	4.53	9.52
Beta	$\alpha=4, \beta=4$	1.09	5.54	10.64	0.85	4.58	9.49
Beta	$\alpha=1, \beta=4$	1.05	5.46	10.77	0.88	4.53	9.42
Beta	$\alpha=2, \beta=4$	1.06	5.53	10.65	0.85	4.56	9.53
Beta	$\alpha=0.5, \beta=1$	0.99	5.35	10.65	0.85	4.48	9.40
Beta	$\alpha=0.5, \beta=2$	1.01	5.31	10.62	0.83	4.54	9.32
Beta	$\alpha=0.5, \beta=3$	1.02	5.28	10.64	0.86	4.55	9.37
LogNormal*	$\mu=1, \sigma^2=0.25$	5.70	14.70	22.33	16.88	31.52	41.57
LogNormal*	$\mu=1, \sigma^2=1.0$	6.72	16.80	25.30	22.34	39.47	49.90
Gamma*	$\gamma=3, \lambda=1$	3.29	9.75	16.62	7.35	17.19	25.13
Gamma*	$\gamma=1, \lambda=1$	2.31	8.10	14.45	4.26	12.35	19.24
Normal*	$\mu=0, \sigma^2=1$	2.49	9.36	15.18	3.99	12.24	19.62
Control	$\mu=0, \sigma^2=1$	1.12	5.56	10.70	0.84	4.58	9.42

Table 6: Percentage of rejections of H_0 of the goodness-of-fit tests for the assumed beta distributions for different levels of significance [%] for the sample size $n=50$. (*distribution is truncated, scaled and/or moved)

Real Distribution	Parameters	χ^2 Test			Kolmogorov-Smirnov Test			BT and Anderson-Darling Test		
		1%	5%	10%	1%	5%	10%	1%	5%	10%
Beta	$\alpha=0.5, \beta=0.5$	2.05	7.71	14.76	0	0.06	0.31	0.86	4.34	9.3
Beta	$\alpha=1, \beta=1$	2.13	7.99	15.08	0	0.05	0.2	0.88	4.49	9.5
Beta	$\alpha=2, \beta=2$	2.12	8.19	15.52	0	0.03	0.15	0.82	4.56	9.52
Beta	$\alpha=4, \beta=4$	2.15	8.19	15.12	0	0.01	0.07	0.84	4.56	9.53
Beta	$\alpha=1, \beta=4$	2.21	8.36	15.81	0	0.03	0.18	0.88	4.54	9.59
Beta	$\alpha=2, \beta=4$	2.25	8.5	16.21	0	0.02	0.1	0.84	4.54	9.57
Beta	$\alpha=0.5, \beta=1$	2.14	8.06	15.06	0	0.09	0.27	0.85	4.36	9.28
Beta	$\alpha=0.5, \beta=2$	2.2	8.16	15.33	0	0.1	0.37	0.84	4.29	9.38
Beta	$\alpha=0.5, \beta=3$	2.13	8.21	15.22	0	0.08	0.27	0.84	4.28	9.35
LogNormal*	$\mu=1, \sigma^2=0.25$	12.54	28.68	40.75	0.63	4.93	11.28	37.06	55.86	65.86
LogNormal*	$\mu=1, \sigma^2=1.0$	19.08	37.61	49.53	2.1	11.46	21.74	49.25	68.12	76.32
Gamma*	$\gamma=3, \lambda=1$	6.18	17.15	27.35	0.06	1.28	3.81	14.16	29.11	39.11
Gamma*	$\gamma=1, \lambda=1$	4.83	14.22	23.38	0.04	0.8	2.32	7.55	18.92	27.99
Normal*	$\mu=0, \sigma^2=1$	5.02	14.96	24.18	0	0.24	1.03	7.04	18.07	26.67
Control	$\mu=0, \sigma^2=1$	-	-	-	-	-	-	0.84	4.61	9.54

Biased transformation and Goodness-of-Fit Test

Table 7: Percentage of rejections of H_0 of the goodness-of-fit tests for the assumed beta distributions for different levels of significance [%] for the sample size $n=100$. (*distribution is truncated, scaled and/or moved.)

Real Distribution	Parameters	χ^2 Test			Kolmogorov-Smirnov Test			BT and Anderson-Darling Test		
		1%	5%	10%	1%	5%	10%	1%	5%	10%
Beta	$\alpha=0.5, \beta=0.5$	2.24	7.46	13.32	0	0.08	0.49	1.27	5.1	10.11
Beta	$\alpha=1, \beta=1$	2.29	7.48	13.23	0	0.04	0.35	1.27	5.12	10.3
Beta	$\alpha=2, \beta=2$	2.37	7.72	13.7	0	0.03	0.16	1.29	5.22	10.3
Beta	$\alpha=4, \beta=4$	2.55	7.73	13.6	0	0.01	0.1	1.29	5.12	10.29
Beta	$\alpha=1, \beta=4$	2.31	7.77	13.76	0	0.05	0.25	1.28	5.1	10.21
Beta	$\alpha=2, \beta=4$	2.57	8.18	14.31	0	0.01	0.13	1.31	5.16	10.25
Beta	$\alpha=0.5, \beta=1$	2.24	7.47	13.31	0	0.1	0.56	1.23	5.06	9.99
Beta	$\alpha=0.5, \beta=2$	2.32	7.46	13.43	0	0.08	0.46	1.25	5.03	9.98
Beta	$\alpha=0.5, \beta=3$	2.32	7.6	13.29	0	0.07	0.41	1.22	5.06	9.93
LogNormal*	$\mu=1, \sigma^2=0.25$	25.19	45.77	58.76	4.26	16.05	27.64	69.88	84.72	89.94
LogNormal*	$\mu=1, \sigma^2=1.0$	39.17	61.08	71.75	11.27	35.3	52.58	84.01	93.39	96.09
Gamma*	$\gamma=3, \lambda=1$	8.8	21.39	31.32	0.74	4.51	9.11	29.8	49.67	60.75
Gamma*	$\gamma=1, \lambda=1$	5.92	15.51	23.76	0.25	2.49	5.88	15.89	31.18	42.14
Normal*	$\mu=0, \sigma^2=1$	4.93	14.31	22.88	0.04	0.43	1.51	12.79	27.82	38.96
Control	$\mu=0, \sigma^2=1$	-	-	-	-	-	-	1.25	5.08	10.28