## ACOUSTIC VIBRATION PROBLEM FOR DISSIPATIVE FLUIDS

### FELIPE LEPE, SALIM MEDDAHI, DAVID MORA, AND RODOLFO RODRÍGUEZ

Abstract. In this paper we analyze a finite element method for solving a quadratic eigenvalue problem derived from the acoustic vibration problem for a heterogeneous dissipative fluid. The problem is shown to be equivalent to the spectral problem for a noncompact operator and a thorough spectral characterization is given. The numerical discretization of the problem is based on Raviart-Thomas finite elements. The method is proved to be free of spurious modes and to converge with optimal order. Finally, we report numerical tests which allow us to assess the performance of the method.

### 1. INTRODUCTION

This paper deals with the numerical approximation of an acoustic dissipative fluid system. This kind of problem has attracted much interest, since it is frequently encountered in engineering applications([\[3,](#page-25-0) [10,](#page-25-1) [15\]](#page-26-0)). One typical example is to achieve optimal designs that reduce noise and vibrations in fluid-structure systems like cars, aircraft or dams.

Although dissipation is usually neglected in standard acoustics, modeling this phenomenon is essential to study the effect of noise reduction techniques. Indeed, in most real situations, damping mechanisms that transform mechanical energy into heat do exist. Sometimes these mechanisms are based on surface damping arising from viscoelastic materials placed on the boundary of the propagation domain. In these cases, the dissipative effects are typically included in the model by means of a surface impedance in the boundary conditions (see, for instance, [\[2,](#page-25-2) [4,](#page-25-3) [5\]](#page-25-4)). The present paper addresses damping when it arises in the propagation media itself due to friction and heat conduction. A general approach to this topic can be found in the books by Landau and Lifshitz [\[12\]](#page-25-5), Morse [\[14\]](#page-26-1), and Pierce [\[19\]](#page-26-2), all of which include extensive bibliographic references on the subject.

This paper focus on computing the (complex) vibration frequencies and modes of an acoustic dissipative fluid system within a rigid cavity. One motivation for considering this problem is that it constitutes a stepping stone towards the more challenging goal of devising numerical approximations for coupled systems involving fluid-structure interaction between viscous fluids and solid structures. The natural model for the fluid system should be based on the Stokes equations for compressible fluids. However, since in real applications the viscosity is typically very small, the

<sup>2000</sup> Mathematics Subject Classification. Primary 65N25, 65N30, 76M10.

Key words and phrases. Spectral problems, dissipative fluid, finite elements, error estimates.

The first author was supported by a CONICYT fellowship (Chile).

The second author was supported by Spain's Ministry of Economy Project MTM2013-43671-P. The third author was partially supported by CONICYT-Chile through FONDECYT project 1140791 (Chile) and by DIUBB through project 151408 GI/VC, Universidad del Bío-Bío, (Chile).

The fourth author was partially supported by BASAL project CMM, Universidad de Chile (Chile).

resulting problem turns out a singular perturbation of that for an inviscid fluid. This fact leads to a kind of dilemma, since appropriate finite elements for the Stokes equations introduce spurious modes in the limit case of a vanishing viscosity, whereas the finite elements that avoid such spectral pollution fail when applied to the Stokes equation.

To circumvent this drawback, we resort to an alternative model based on a curlfree displacement formulation (see [\[6\]](#page-25-6) for the derivation of a similar model in the time domain from basic mechanical laws). Let us remark that in principle the fluid displacement does not need to be curl-free. However, since the viscosity term due to vorticity is typically very small, except perhaps near the walls of the enclosure, it may be neglected in the interior of the enclosure and eventually modeled as a wall impedance on its boundary (see [\[15\]](#page-26-0) for a similar model).

The numerical solution of the vibration problem for an inviscid acoustic homogeneous fluid is nowadays a well known subject (see, for instance, [\[3\]](#page-25-0)). In its turn, as is shown in Remark 2.1 of the present paper, the vibration frequencies of a viscous homogeneous irrotational fluid within a rigid cavity can be algebraically computed from those of the analogous inviscid fluid and the corresponding vibration modes coincide. However, this is not the case for a heterogeneous fluid and this is the reason why we choose this as our model problem. In particular, we consider the acoustic vibration problem for a dissipative fluid system that consists of two homogeneous viscous immiscible fluids contained in a rigid cavity.

We begin with a variational formulation of the spectral problem relying only on the fluid displacement, which leads to a quadratic eigenvalue problem. For the theoretical analysis, this is transformed into an equivalent double-size linear eigenvalue problem. We introduce a convenient functional framework to analyze it and prove that the nonlinear eigenvalue problem is equivalent to the spectral problem for a nonselfadjoint, noncompact bounded operator. Thus, the essential spectrum not necessarily reduces to zero (as is the case for compact operators). This means that the spectrum may now contain nonzero eigenvalues of infinite-multiplicity, nonzero accumulation points, continuous spectrum, etc. Thus, following [\[11\]](#page-25-7), our first task is to prove that the relevant eigenvalues can be isolated from the essential spectrum, at least for sufficiently small values of the viscosity that are realistic in practice. Then, we propose a conforming discretization based on Raviart-Thomas finite elements. By appropriately adapting the abstract spectral approximation theory for noncompact operators developed in [\[7,](#page-25-8) [8\]](#page-25-9), we establish that the resulting scheme provides a correct approximation of the spectrum and prove error estimates for the eigenfunctions and a double order for the eigenvalues. Moreover, the discrete quadratic eigenvalue problem is shown to be equivalent to a well posed generalized eigenvalue problem which can be solved by standard eigensolvers like eigs from MATLAB, which is based on Arnoldi iterations.

The paper is organized as follows: in Section [2,](#page-2-0) we introduce the spectral problem and the corresponding variational formulation, which leads to a quadratic eigenvalue problem. We introduce an auxiliary unknown to transform the quadratic eigenvalue problem into a linear one. Moreover, we introduce the corresponding solution operator for the spectral problem. In Section [3,](#page-7-0) we provide a thorough spectral characterization of the solution operator, based on the theory developed in [\[11\]](#page-25-7). We also consider the limit problem (i.e., the case when the viscosity vanishes) and the relation between the solutions of the dissipative and non-dissipative problems. In Section [4,](#page-10-0) we introduce a finite element discretization using Raviart-Thomas elements for both fluids and imposing the continuity of the corresponding normal components on the interface. We analyze the discrete spectral problem analogously as in the continuous case and introduce the corresponding discrete solution operator. We use the abstract theory from [\[7\]](#page-25-8) to prove the convergence. We also prove error estimates for our problem by adapting the arguments from [\[2\]](#page-25-2). Finally, in Section [5,](#page-19-0) we report some numerical tests which allow us to asses the performance of the proposed method.

Throughout the paper,  $\Omega$  is a generic Lipschitz bounded domain of  $\mathbb{R}^d$  (*d* = 2, 3), with outer unit normal vector n. We denote by  $\mathcal{D}(\Omega)$  the space of infinitely smooth function compactly supported in  $\Omega$ . For  $r \geq 0$ ,  $\|\cdot\|_{r,\Omega}$  stands indistinctly for the norm of the Hilbertian Sobolev spaces  $H^r(\Omega)$  or  $H^r(\Omega)^d$  with the convention  $\mathrm{H}^{0}(\Omega) := \mathrm{L}^{2}(\Omega)$ . We also define the Hilbert space  $\mathrm{H}(\mathrm{div};\Omega) := \{v \in \mathrm{L}^{2}(\Omega)^{d}$ :  $\mathrm{div}\,\boldsymbol{v}\in\mathrm{L}^2(\Omega)\},\,\text{whose norm is given by }\|\boldsymbol{v}\|_{\mathrm{d}}^2$  $\frac{2}{\mathrm{div},\Omega}:=\left\|\bm{v}\right\|_{0,\Omega}^{2}+\left\|\mathrm{div}\,\bm{v}\right\|_{0}^{2}$  $\epsilon_{0,\Omega}^2$ , and its subspace  $H_0(\text{div}; \Omega) := \{v \in H(\text{div}; \Omega) : v \cdot n = 0 \text{ on } \partial\Omega\}.$ 

<span id="page-2-0"></span>Finally, C represents a generic constant independent of the discretization parameters, which may take different values at different places.

## 2. The model problem

We take as our model problem the case of two immiscible fluids within a rigid cavity. Let  $\Omega_i$  with  $i = 1, 2$  be the polygonal (in the 2D case) or polyhedral (in the 3D case) Lipschitz domains occupied by each of the fluids. Let  $\rho_i$  be the corresponding densities,  $\nu_i$  the fluid viscosities, and  $c_i$  the acoustic speeds, which we consider all constant,  $\rho_i$  and  $c_i$  strictly positive and  $\nu_i$  non negative. We denote by  $n_i$  the outward unit normal vectors corresponding to each subdomain. We define  $\Omega := (\Omega_1 \cup \Omega_2)^\circ, \Gamma := \partial \Omega_1 \cap \partial \Omega_2$ , and  $\Gamma_i := \partial \Omega_i \cap \partial \Omega, i = 1, 2$ . We assume that each domain  $\Omega_i$  as well as  $\Omega$  are simply connected (see Figure [1\)](#page-2-1).



<span id="page-2-1"></span>FIGURE 1. 2D sketch of the polygonal domains for the fluids.

We consider small displacements of a compressible viscous fluid at rest neglecting convective terms. The equation of motion derived from the Stokes equation reads

$$
\rho_i \ddot{\mathbf{U}}_i = 2\nu_i \Delta \dot{\mathbf{U}}_i - \nabla P_i \quad \text{in } \Omega_i,
$$

where  $U_i$  denotes the fluid displacement and  $P_i$  the pressure fluctuation in the domain  $\Omega_i$ ,  $i = 1, 2$ . The dot represents derivation with respect to time. Moreover, since the fluid is compressible, we consider the isentropic relation

$$
P_i + \rho_i c_i^2 \operatorname{div} \boldsymbol{U}_i = 0 \qquad \text{in } \Omega_i.
$$

Since we are considering irrotational fluids, we assume that  $\text{curl } U_i = 0$ . Hence, considering the identity  $\Delta \vec{U}_i = \nabla(\text{div } \vec{U}_i) - \text{curl}(\text{curl } \vec{U}_i)$ , we conclude that  $\Delta \dot{U}_i = \nabla(\text{div }\dot{U}_i)$ . Then, the equations of our model problem are the following:

<span id="page-3-0"></span>(2.1)  $\rho_1 \ddot{\mathbf{U}}_1 - 2\nu_1 \nabla (\text{div } \dot{\mathbf{U}}_1) + \nabla P_1 = 0 \text{ in } \Omega_1 \times (0, T),$ 

<span id="page-3-2"></span>(2.2) 
$$
P_1 + \rho_1 c_1^2 \operatorname{div} U_1 = 0 \quad \text{in } \Omega_1 \times [0, T],
$$

<span id="page-3-1"></span>(2.3) 
$$
\rho_2 \ddot{U}_2 - 2\nu_2 \nabla (\text{div } \dot{U}_2) + \nabla P_2 = 0 \text{ in } \Omega_2 \times (0, T),
$$

<span id="page-3-3"></span>(2.4) 
$$
P_2 + \rho_2 c_2^2 \operatorname{div} U_2 = 0 \quad \text{in } \Omega_2, \times [0, T],
$$

(2.5) 
$$
U_1 \cdot n_1 + U_2 \cdot n_2 = 0 \quad \text{on } \Gamma \times [0, T],
$$

(2.6) 
$$
(2\nu_1 \operatorname{div} \dot{\mathbf{U}}_1 + P_1) - (2\nu_2 \operatorname{div} \dot{\mathbf{U}}_2 + P_2) = 0 \text{ on } \Gamma \times (0, T),
$$

$$
U_1 \cdot \mathbf{n}_1 = 0 \quad \text{on } \Gamma_1 \times (0,T),
$$

<span id="page-3-7"></span>(2.8) U<sup>2</sup> · n<sup>2</sup> = 0 on Γ<sup>2</sup> × (0, T ).

Let us remark that a similar argument leads exactly to the same equations in 2D.

Multiplying equations [\(2.1\)](#page-3-0) and [\(2.3\)](#page-3-1) by a test function  $v \in H_0(\text{div}; \Omega)$ , integrating by parts, and using the boundary conditions and the transmission conditions on Γ, we obtain

(2.9) 
$$
\int_{\Omega} \rho \ddot{\mathbf{U}} \cdot \mathbf{v} + 2 \int_{\Omega} \nu \operatorname{div} \dot{\mathbf{U}} \operatorname{div} \mathbf{v} - \int_{\Omega} P \operatorname{div} \mathbf{v} = 0 \quad \forall \mathbf{v} \in H_0(\operatorname{div}, \Omega),
$$

where

<span id="page-3-4"></span>
$$
U := \begin{cases} U_1 & \text{in } \Omega_1, \\ U_2 & \text{in } \Omega_2, \end{cases} \quad P := \begin{cases} P_1 & \text{in } \Omega_1, \\ P_2 & \text{in } \Omega_2, \end{cases} \quad \nu := \begin{cases} \nu_1 & \text{in } \Omega_1, \\ \nu_2 & \text{in } \Omega_2, \end{cases}
$$

$$
\rho := \begin{cases} \rho_1 & \text{in } \Omega_1, \\ \rho_2 & \text{in } \Omega_2, \end{cases} \quad \text{and} \quad c := \begin{cases} c_1 & \text{in } \Omega_1, \\ c_2 & \text{in } \Omega_2. \end{cases}
$$

<span id="page-3-6"></span>Using  $(2.2)$  and  $(2.4)$  we eliminate P in  $(2.9)$  and write

$$
(2.10)\quad \int_{\Omega} \rho \ddot{\boldsymbol{U}} \cdot \boldsymbol{v} + 2 \int_{\Omega} \nu \operatorname{div} \dot{\boldsymbol{U}} \operatorname{div} \boldsymbol{v} + \int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{U} \operatorname{div} \boldsymbol{v} = 0 \qquad \forall \boldsymbol{v} \in H_0(\operatorname{div}, \Omega).
$$

The *vibration modes* of this problem are complex solutions of the form  $U(x, t) =$  $e^{\lambda t}u(x)$  with  $\lambda \in \mathbb{C}$ . Looking for this kind of solutions leads to the following quadratic eigenvalue problem:

<span id="page-3-5"></span>**Problem 1.** Find  $\lambda \in \mathbb{C}$  and  $\theta \neq u \in H_0(\text{div}; \Omega)$  such that

$$
\lambda^2 \int_{\Omega} \rho \mathbf{u} \cdot \overline{\mathbf{v}} + 2\lambda \int_{\Omega} \nu \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} + \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} = 0 \qquad \forall \mathbf{v} \in H_0(\operatorname{div}; \Omega).
$$

Let us remark that in absence of viscosity (i.e.,  $\nu = 0$ ) we are left with the free vibration problem of two inviscid fluids in contact (whose numerical approximation has not been analyzed either). The eigenvalues  $\lambda^2$  of such a problem are negative real numbers (as will be proved below), so that  $\lambda$  are purely imaginary, namely,  $\lambda = \pm i\omega$  with  $\omega$  being the so called *natural vibration frequencies*, which correspond to periodic in time solutions  $U(x,t) = e^{-i\omega t} u(x)$  of the time domain problem. This

is the reason why, for  $\nu = 0$ , Problem [1](#page-3-5) is usually written as follows: Find  $\omega > 0$ and  $\underline{0} \neq \mathbf{u} \in H_0(\text{div}; \Omega)$  such that

<span id="page-4-0"></span>(2.11) 
$$
\int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u} \operatorname{div} \overline{\boldsymbol{v}} = \omega^2 \int_{\Omega} \rho \boldsymbol{u} \cdot \overline{\boldsymbol{v}} \qquad \forall \boldsymbol{v} \in H_0(\operatorname{div}; \Omega).
$$

In the applications,  $\nu$  is typically very small. As we will show below, in such a case there are eigenvalues  $\lambda$  of Problem [1](#page-3-5) that lie close to  $\pm i\omega$  with  $\omega$  being a natural vibration frequency (i.e., a solution of [\(2.11\)](#page-4-0)). Actually, we will prove below that those  $\lambda$  converge to  $\pm i\omega$  as  $\|\nu\|_{\infty,\Omega}$  goes to zero. On solving Problem [1,](#page-3-5) the aim is to compute the eigenvalues  $\lambda$  close to the smallest natural vibration frequencies  $\omega > 0$ , which are the most relevant in the applications.

**Remark 2.1.** In the case of a homogeneous viscous fluid,  $\rho$ , c and  $\nu$  are constant in the whole  $\Omega$ . Then, Problem [1](#page-3-5) can be written as

$$
\lambda^2 \int_{\Omega} \rho \mathbf{u} \cdot \overline{\mathbf{v}} + \frac{2\lambda \nu + \rho c^2}{\rho c^2} \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} = 0 \quad \forall \mathbf{v} \in H_0(\operatorname{div}, \Omega).
$$

Hence, in such a case,  $(\lambda, \mathbf{u})$  is an eigenpair of Problem [1](#page-3-5) if and only if  $-\frac{\lambda^2 \rho c^2}{2\lambda \nu + \rho c^2}$  $\omega^2$  with  $(\omega, \mathbf{u})$  being a solution to problem [\(2.11\)](#page-4-0). Therefore, for a homogeneous viscous fluid,  $\lambda$  can be algebraically computed from the solution of [\(2.11\)](#page-4-0) as follows:

$$
\lambda = \frac{-\nu\omega^2 \pm \sqrt{\nu^2\omega^4 - \rho^2c^4\omega^2}}{\rho c^2}
$$

.

We denote  $\mathcal{H} := L^2(\Omega)^d$  endowed with the weighted inner product

$$
(\boldsymbol{v},\boldsymbol{w})_{\mathcal{H}}:=\int_{\Omega}\rho\boldsymbol{v}\cdot\overline{\boldsymbol{w}}
$$

and  $V := H_0(\text{div}; \Omega)$  with the inner product

$$
(\boldsymbol{v},\boldsymbol{w})_{\mathcal{V}} := \int_{\Omega} \rho \boldsymbol{v} \cdot \overline{\boldsymbol{w}} + \int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{v} \operatorname{div} \overline{\boldsymbol{w}}.
$$

Notice that the inner products in H and V induce norms  $\|\cdot\|_{\mathcal{H}}$  and  $\|\cdot\|_{\mathcal{V}}$  on each of these spaces equivalent to the classical  $L^2(\Omega)^d$  and  $H(\text{div}; \Omega)$  norms, respectively. Therefore, when it might be convenient, we will use these classical norms.

Clearly  $\lambda = 0$  is an eigenvalue of Problem [1](#page-3-5) with associated eigenspace

$$
\mathcal{K} = H_0(\mathrm{div}^0, \Omega) := \{ \mathbf{v} \in H_0(\mathrm{div}; \Omega) : \text{ div } \mathbf{v} = 0 \text{ in } \Omega \}.
$$

We define:

$$
\mathcal{G} := \mathcal{K}^{\perp_{\mathcal{V}}} = \{ \mathbf{v} \in \mathcal{V} : (\mathbf{v}, \mathbf{w})_{\mathcal{V}} = 0 \quad \forall \mathbf{w} \in \mathcal{K} \}.
$$

Since K is a closed subspace of V, clearly  $\mathcal{V} = \mathcal{G} \oplus \mathcal{K}$ . Notice that  $\mathcal{G}$  and K are also orthogonal in the  $H$  inner product. Hence,

$$
\mathcal{G} = \{ \boldsymbol{v} \in \mathcal{V} : (\boldsymbol{v}, \boldsymbol{w})_{\mathcal{H}} = 0 \ \forall \boldsymbol{w} \in \mathcal{K} \}.
$$

The following result brings a characterization of the space  $\mathcal{G}$ .

<span id="page-4-1"></span>Lemma 2.1. There holds

$$
\mathcal{G}=\frac{1}{\rho}\nabla(\mathrm{H}^1(\Omega))\cap \mathcal{V}.
$$

*Proof.* We will prove this result by checking the double inclusion. Let  $v \in \mathcal{G}$ . Then, for all  $\psi \in \mathcal{D}(\Omega)^d$ , since **curl**  $\psi \in \mathcal{K}$ , we have

$$
0 = \int_{\Omega} \rho \operatorname{curl} \boldsymbol{\psi} \cdot \overline{\boldsymbol{v}} = \int_{\Omega} \boldsymbol{\psi} \cdot \operatorname{curl} \left( \rho \overline{\boldsymbol{v}} \right).
$$

Thus, curl( $\rho v$ ) = 0 in  $\Omega$ . Since  $\Omega$  is simply connected, this implies that there exists  $\varphi \in H^1(\Omega)$  such that  $\rho v = \nabla \varphi$ . Hence,  $v \in \frac{1}{\rho} \nabla(H^1(\Omega)) \cap \mathcal{V}$ . Conversely, let  $v \in \frac{1}{\rho} \nabla(\mathrm{H}^1(\Omega)) \cap \mathcal{V}$  and  $w \in \mathcal{K}$ . Let  $\varphi \in \mathrm{H}^1(\Omega)$  be such that  $v = \frac{1}{\rho} \nabla \varphi$ . Then,

$$
(\mathbf{v}, \mathbf{w})_{\mathcal{H}} = \int_{\Omega} \rho \left( \frac{1}{\rho} \nabla \varphi \right) \cdot \overline{\mathbf{w}} = - \int_{\Omega} \varphi \operatorname{div} \overline{\mathbf{w}} + \int_{\partial \Omega} \varphi (\overline{\mathbf{w}} \cdot \mathbf{n}) = 0.
$$

Therefore,  $v \in \mathcal{G}$ . The proof is complete.

In what follows we prove additional regularity for the functions in  $\mathcal{G}$ . From now and on, s will denote a positive number such that the following lemma holds true.

<span id="page-5-0"></span>**Lemma 2.2.** There exists  $s > 0$  (with s depending on  $\rho$  and  $\Omega$ ) such that  $v \in$  $\mathrm{H}^s(\Omega)^d$  for all  $\boldsymbol{v} \in \mathcal{G}$  and

(2.12) kvks,<sup>Ω</sup> ≤ Ck div vk<sup>0</sup>,<sup>Ω</sup>,

where C is a positive constant independent of  $v$ .

*Proof.* According to Lemma [2.1,](#page-4-1) there exists  $\varphi \in H^1(\Omega)$  such that  $\mathbf{v} = \frac{1}{\rho} \nabla \varphi$ . Consequently,  $\varphi \in H^1(\Omega)/\mathbb{C}$  is the unique solution of the following well-posed Neumann problem:

<span id="page-5-1"></span>
$$
\operatorname{div}\left(\frac{1}{\rho}\nabla\varphi\right) = \operatorname{div}\mathbf{v} \quad \text{in } \Omega,
$$

$$
\frac{1}{\rho}\frac{\partial\varphi}{\partial\mathbf{n}} = 0 \qquad \text{on } \partial\Omega.
$$

Hence, in the 3D case, the theorem follows from [\[16,](#page-26-3) Lemma 2.20] with

$$
s:=\frac{1}{2\pi}\min\left\{\min_{\Omega}\{\rho\},\frac{1}{\max_{\Omega}\{\rho\}}\right\}>0.
$$

For the 2D case, the theorem follows by applying [\[16,](#page-26-3) Lemma 4.3]. (See [\[18\]](#page-26-4) for more details.)  $\Box$ 

From the physical point of view, the time domain problem [\(2.10\)](#page-3-6) is dissipative in the sense that its solution should decay as  $t$  increases. The latter happens if and only if the so called *decay rate*,  $\text{Re}(\lambda)$ , is negative. The following result shows that this is the case in our formulation.

**Lemma 2.3.** Let  $(\lambda, u) \in \mathbb{C} \times \mathcal{V}$  be a solution of Problem [1.](#page-3-5) If  $\lambda \neq 0$ , then  $Re(\lambda) < 0.$ 

*Proof.* Testing Problem [1](#page-3-5) with  $\mathbf{v} = \mathbf{u}$ , we have that  $A\lambda^2 + B\lambda + C = 0$ , with

$$
A:=\int_{\Omega}\rho|\textbf{\textit{u}}|^2,\quad B:=2\int_{\Omega}\nu|\operatorname{div}\textbf{\textit{u}}|^2,\quad\text{and}\quad C:=\int_{\Omega}\rho c^2|\operatorname{div}\textbf{\textit{u}}|^2
$$

.

We observe that  $A > 0$ ,  $B \ge 0$ , and  $C \ge 0$ . Moreover, since  $\lambda = 0$  if and only if  $u \in \mathcal{K} = H_0(\text{div}^0, \Omega)$ , for  $\lambda \neq 0$  we have that  $B, C > 0$ , too. Therefore, since  $\lambda = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}$ , it is immediate to check that  $\text{Re}(\lambda) < 0$ . <span id="page-6-5"></span>**Remark 2.2.** Any eigenpair  $(\lambda, \mathbf{u})$  of Problem [1](#page-3-5) satisfies

$$
\lambda^2 \int_{\Omega} \rho \mathbf{u} \cdot \overline{\mathbf{v}} + \int_{\Omega} (2\lambda \nu + \rho c^2) \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} = 0 \quad \forall \mathbf{v} \in \mathcal{V}.
$$

Since the coefficients are constant in each subdomain, if  $2\lambda \nu + \rho c^2 \neq 0$  in  $\Omega_i$ , by testing with  $\mathbf{v} \in \mathcal{D}(\Omega_i)^d$  we obtain that  $\text{div } \mathbf{u}|_{\Omega_i} \in H^1(\Omega_i)$ ,  $i = 1, 2$ . On the other hand, if  $2\lambda \nu + \rho c^2 = 0$  in  $\Omega_i$  (i = 1 or 2), then, for  $\lambda \neq 0$ , by testing again with  $v \in \mathcal{D}(\Omega_i)^d$ , we obtain that  $u = \underline{\theta}$  in  $\Omega_i$ . Thus, in any case, div  $u|_{\Omega_i} \in H^1(\Omega_i)$ ,<br> $i = 1, 2$  $i = 1, 2.$ 

For the theoretical analysis it is convenient to transform Problem [1](#page-3-5) into a linear eigenvalue problem. With this aim we introduce the new variable  $\hat{u} := \lambda u$ , as usual in quadratic problems, and the space  $\tilde{\mathcal{V}} := \mathcal{V} \times \mathcal{H}$  endowed with the corresponding product norm, which carry us to the following:

<span id="page-6-0"></span>**Problem 2.** Find  $\lambda \in \mathbb{C}$  and  $\theta \neq (\mathbf{u}, \widehat{\mathbf{u}}) \in \widetilde{\mathcal{V}}$  such that

<span id="page-6-1"></span>(2.13) 
$$
\int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} = \lambda \left( -2 \int_{\Omega} \nu \operatorname{div} \mathbf{u} \operatorname{div} \overline{\mathbf{v}} - \int_{\Omega} \rho \widehat{\mathbf{u}} \cdot \overline{\mathbf{v}} \right) \qquad \forall \mathbf{v} \in \mathcal{V},
$$
  
(2.14) 
$$
\int_{\Omega} \rho \widehat{\mathbf{u}} \cdot \overline{\widehat{\mathbf{v}}} = \lambda \int_{\Omega} \rho \mathbf{u} \cdot \overline{\widehat{\mathbf{v}}} \qquad \forall \widehat{\mathbf{v}} \in \mathcal{H}.
$$

<span id="page-6-2"></span>We observe that  $\lambda = 0$  is an eigenvalue of Problem [2](#page-6-0) and its associated eigenspace is  $\widetilde{\mathcal{K}} := \mathcal{K} \times \{0\}$ . Let  $\widetilde{\mathcal{G}}$  be the orthogonal complement of  $\widetilde{\mathcal{K}}$  in  $\mathcal{V} \times \mathcal{H}$ . Notice that  $\widetilde{\mathcal{G}} = \mathcal{G} \times \mathcal{H}.$ 

We introduce the sesquilinear continuous form  $a: \mathcal{V} \times \mathcal{V} \to \mathbb{C}$  defined by

$$
a(\boldsymbol{u},\boldsymbol{v}) := \int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u} \operatorname{div} \overline{\boldsymbol{v}},
$$

and the sesquilinear continuous forms  $\tilde{a}, \tilde{b} : \tilde{V} \to \tilde{V}$  defined as follows:

$$
\widetilde{a}((\boldsymbol{u},\widehat{\boldsymbol{u}}),(\boldsymbol{v},\widehat{\boldsymbol{v}})) := \int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u} \operatorname{div} \overline{\boldsymbol{v}} + \int_{\Omega} \rho \widehat{\boldsymbol{u}} \cdot \overline{\widehat{\boldsymbol{v}}},
$$

$$
\widetilde{b}((\boldsymbol{u},\widehat{\boldsymbol{u}}),(\boldsymbol{v},\widehat{\boldsymbol{v}})) := -2 \int_{\Omega} \nu \operatorname{div} \boldsymbol{u} \operatorname{div} \overline{\boldsymbol{v}} - \int_{\Omega} \rho \widehat{\boldsymbol{u}} \cdot \overline{\boldsymbol{v}} + \int_{\Omega} \rho \boldsymbol{u} \cdot \overline{\widehat{\boldsymbol{v}}}.
$$

In what follows we prove that  $a(\cdot, \cdot)$  and  $\widetilde{a}(\cdot, \cdot)$  are elliptic in  $\mathcal G$  and  $\widetilde{\mathcal G}$ , respectively.

<span id="page-6-3"></span>**Lemma 2.4.** The sesquilinear form  $a : \mathcal{G} \times \mathcal{G} \to \mathbb{C}$  is  $\mathcal{G}$ -elliptic and, consequently,  $\widetilde{a} : \widetilde{\mathcal{G}} \times \widetilde{\mathcal{G}} \to \mathbb{C}$  is  $\widetilde{\mathcal{G}}$ -elliptic.

*Proof.* For  $v \in \mathcal{G}$  we have

<span id="page-6-4"></span>(2.15) 
$$
a(\mathbf{v}, \mathbf{v}) = \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{v} \operatorname{div} \overline{\mathbf{v}} \ge \min_{\Omega} \{ \rho c^2 \} || \operatorname{div} \mathbf{v} ||_{0,\Omega}^2.
$$

Then, the G-ellipticity of  $a(\cdot, \cdot)$  follows from Lemma [2.2.](#page-5-0) From this, the ellipticity of  $\widetilde{a}(\cdot,\cdot)$  in  $\widetilde{\mathcal{G}} = \mathcal{G} \times \mathcal{H}$  is immediate.

Let  $T: \widetilde{\mathcal{V}} \to \widetilde{\mathcal{V}}$  be the bounded linear operator defined by  $T(f, g) := (u, \widehat{u}) \in \widetilde{\mathcal{G}}$ , where  $(u,\hat{u})$  is the unique solution of the following problem:

$$
\widetilde{a}((\boldsymbol{u},\widehat{\boldsymbol{u}}),(\boldsymbol{v},\widehat{\boldsymbol{v}}))=\widetilde{b}((\boldsymbol{f},\boldsymbol{g}),(\boldsymbol{v},\widehat{\boldsymbol{v}}))\qquad\forall(\boldsymbol{v},\widehat{\boldsymbol{v}})\in\widetilde{\mathcal{G}}.
$$

It is easy to check that

<span id="page-7-1"></span>
$$
\widehat{\mathbf{u}} = \mathbf{f} \quad \text{in } \Omega,
$$

and

<span id="page-7-6"></span>(2.17) 
$$
\int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u} \operatorname{div} \overline{\boldsymbol{v}} = -2 \int_{\Omega} \nu \operatorname{div} \boldsymbol{f} \operatorname{div} \overline{\boldsymbol{v}} - \int_{\Omega} \rho \boldsymbol{g} \cdot \overline{\boldsymbol{v}} \qquad \forall \boldsymbol{v} \in \mathcal{G}.
$$

As a consequence of the above equalities, we have that  $\mu = 0$  is an eigenvalue of T with associated eigenspace  $\{0\} \times \mathcal{G}^{\perp_{\mathcal{H}}}$ , which is nontrivial since  $\mathcal{G}^{\perp_{\mathcal{H}}} \supset \mathcal{K}$ . The Following lemma shows that the nonzero eigenvalues of  $T$  are exactly the reciprocals of the nonzero eigenvalues of Problem [2](#page-6-0) with the same corresponding eigenfunctions.

<span id="page-7-3"></span>**Lemma 2.5.** There holds that  $(\mu, (\boldsymbol{u}, \hat{\boldsymbol{u}}))$  is an eigenpair of **T** (i.e.  $T(\boldsymbol{u}, \hat{\boldsymbol{u}}) =$  $\mu(\mathbf{u},\hat{\mathbf{u}})$ ) with  $\mu \neq 0$  if and only if  $(\lambda,(\mathbf{u},\hat{\mathbf{u}}))$  is a solution of Problem [2](#page-6-0) with  $\lambda =$  $1/\mu \neq 0.$ 

*Proof.* Let  $(\mu, (\boldsymbol{u},\hat{\boldsymbol{u}}))$  be an eigenpair of T with  $\mu \neq 0$ . Hence

<span id="page-7-2"></span>(2.18) 
$$
\widetilde{a}((\boldsymbol{u},\widehat{\boldsymbol{u}}),(\boldsymbol{v},\widehat{\boldsymbol{v}}))=\frac{1}{\mu}\widetilde{b}((\boldsymbol{u},\widehat{\boldsymbol{u}}),(\boldsymbol{v},\widehat{\boldsymbol{v}})) \qquad \forall (\boldsymbol{v},\widehat{\boldsymbol{v}})\in \widetilde{\mathcal{G}}.
$$

Then, according to [\(2.16\)](#page-7-1) we have that  $\hat{u} = \frac{1}{\mu}u \in \mathcal{G}$ . Hence, for  $(v, \hat{v}) \in \tilde{\mathcal{K}}$  =  $\mathcal{K} \times \{0\}$ , clearly  $b((\boldsymbol{u}, \hat{\boldsymbol{u}}),(\boldsymbol{v}, \hat{\boldsymbol{v}})) = 0$  and  $\tilde{a}((\boldsymbol{u}, \hat{\boldsymbol{u}}),(\boldsymbol{v}, \hat{\boldsymbol{v}})) = 0$ . So, [\(2.18\)](#page-7-2) holds for all  $(v, \hat{v}) \in \tilde{V} = \tilde{\mathcal{G}} \oplus \tilde{\mathcal{K}}$ ; namely,  $(\lambda, (\mathbf{u}, \hat{\mathbf{u}}))$  with  $\lambda = 1/\mu$  is a solution to Problem [2.](#page-6-0)

Conversely, let  $(\lambda, (\boldsymbol{u},\hat{\boldsymbol{u}}))$  be a solution of Problem [2](#page-6-0) with  $\lambda \neq 0$ . Taking  $\boldsymbol{v} \in \mathcal{K}$ in [\(2.13\)](#page-6-1), we have that  $\int_{\Omega} \rho \hat{\mathbf{u}} \cdot \mathbf{v} = 0$ , which implies that  $\hat{\mathbf{u}} \in \mathcal{G}$ . On the other hand, we observe that [\(2.14\)](#page-6-2) implies that  $\lambda u = \hat{u} \in \mathcal{G}$ . Hence it is easy to check that  $T(u, \hat{u}) = u(u, \hat{u})$  with  $u = 1/\lambda$ .  $T(u, \hat{u}) = \mu(u, \hat{u})$  with  $\mu = 1/\lambda$ .

### 3. Spectral Characterization

<span id="page-7-0"></span>The goal of this section is to characterize the spectrum of the solution operator **T**. Since the inclusion  $H_0(\text{div}; \Omega) \hookrightarrow L^2(\Omega)^d$  is not compact, it is easy to check from  $(2.16)$  that  $\boldsymbol{T}$  is not compact either. However, we will show that the essential spectrum, has to lie in a small region of the complex plane, well separated from the isolated eigenvalues which, according to Lemma [2.5,](#page-7-3) correspond to the solutions of Problem [2.](#page-6-0) With this aim, we will resort to the theory described in [\[11\]](#page-25-7) to decompose appropriately T. Let  $T_1, T_2 : \mathcal{G} \to \mathcal{G}$  be the operators given by

<span id="page-7-5"></span>(3.1) 
$$
T_1f = u_1 \in \mathcal{G}: \quad a(u_1, v) = 2 \int_{\Omega} \nu \operatorname{div} f \operatorname{div} \overline{v} \quad \forall v \in \mathcal{G},
$$

<span id="page-7-4"></span>(3.2) 
$$
T_2g = u_2 \in \mathcal{G} : a(u_2, v) = \int_{\Omega} \rho g \cdot \overline{v} \quad \forall v \in \mathcal{G}.
$$

It is easy to check that these operators are self-adjoint with respect to  $a(\cdot, \cdot)$ . Moreover  $T_1$  is non-negative and  $T_2$  is positive with respect to  $a(\cdot, \cdot)$  (namely,  $a(\mathbf{T}_1 v, v) \geq 0 \ \forall v \in \mathcal{G}$  and  $a(\mathbf{T}_2 v, v) > 0 \ \forall v \in \mathcal{G}, v \neq 0$ ). Moreover, we have the following result following result.

# **Lemma 3.1.** The operator  $T_2 : \mathcal{G} \to \mathcal{G}$  is compact.

*Proof.* Since  $a(\cdot, \cdot)$  is G-elliptic (cf. Lemma [2.4\)](#page-6-3), applying Lax-Milgram's Lemma, we know that problem [\(3.2\)](#page-7-4) is well posed and has a unique solution  $u_2 \in \mathcal{G}$ . Moreover, according to Lemma [2.2,](#page-5-0) we know that there exists  $s > 0$  such that

 $u_2 \in H^s(\Omega)^d$ . On the other hand, notice that  $(3.2)$  also holds for  $v \in \mathcal{K}$ , since in such a case  $a(\boldsymbol{u}_2, \boldsymbol{v}) = 0 = \int_{\Omega} \rho \boldsymbol{g} \cdot \boldsymbol{\overline{v}}$  for  $\boldsymbol{g} \in \mathcal{G}$ . Hence, since  $\mathcal{V} = \mathcal{G} \oplus \mathcal{K}$ , we have that

$$
a(\boldsymbol{u}_2,\boldsymbol{v})=\int_{\Omega}\rho\boldsymbol{g}\cdot\overline{\boldsymbol{v}}\quad\forall\boldsymbol{v}\in\mathcal{V}.
$$

Then, by testing this equation with  $\mathbf{v} \in \mathcal{D}(\Omega)^d \subset \mathcal{V}$ , we have that  $-\nabla(\rho c^2 \operatorname{div} \mathbf{u}_2) =$  $\rho g$  in  $\Omega$ , so that  $\rho c^2$  div  $u_2 \in H^1(\Omega)$ . Therefore, since  $\rho$  and c are positive constants in each subdomain  $\Omega_1$  and  $\Omega_2$ , we have that div  $u_2|_{\Omega_i} \in H^1(\Omega_i)$ ,  $i = 1, 2$ . Since the inclusions  $\{v \in \mathcal{L}^2(\Omega) : v|_{\Omega_i} \in \mathcal{H}^1(\Omega_i), i = 1, 2\} \subset \mathcal{L}^2(\Omega)$  and  $\mathcal{H}^s(\Omega)^d \subset \mathcal{L}^2(\Omega)^d$ , are compact, we derive that  $T_2$  is compact too.

The operator T can be written in terms of the operators  $T_1$  and  $T_2$  given above as follows:

$$
\boldsymbol{T} = \begin{pmatrix} -\boldsymbol{T}_1 & -\boldsymbol{T}_2 \\ \boldsymbol{I} & 0 \end{pmatrix}.
$$

Moreover, by defining as in [\[11\]](#page-25-7) the operators

$$
\boldsymbol{S}:=\begin{pmatrix}\boldsymbol{I}&0\\0&\boldsymbol{T}^{1/2}_2\end{pmatrix}\qquad\text{and}\qquad \boldsymbol{H}:=\begin{pmatrix}-\boldsymbol{T}_1&-\boldsymbol{T}^{1/2}_2\\ \boldsymbol{T}^{1/2}_2&0\end{pmatrix},
$$

we have that  $ST = HS$ . We note that the eigenvalues of T and H and their algebraic multiplicities coincide. Moreover the corresponding Jordan chains have the same length. In fact, let  $\{x_k\}_{k=1}^r$  be a Jordan chain associated with the eigenvalue  $\mu$  of  $\bm{T}$ . Then, using the identities above, we observe that

$$
H S x_k = ST x_k = S(\mu x_k + x_{k-1}) = \mu S x_k + S x_{k-1}, \quad k=1,\ldots,r.
$$

This shows that  $\{S_x\}_{k=1}^r$  is a Jordan chain of  $H$  of the same length. Actually, the whole spectra of  $T$  and  $H$  coincide as is shown in the following result, which has been proved in Lemma 3.2 of [\[2\]](#page-25-2).

<span id="page-8-0"></span>Lemma 3.2. There holds

$$
Sp(T) = Sp(H).
$$

Moreover,  $Sp_{ess}(T) = Sp_{ess}(H)$ , too.

The operator  $H$  can be written as the sum of a self-adjoint operator  $B$  and a compact operator C:

$$
\boldsymbol{H} = \boldsymbol{B} + \boldsymbol{C} \quad \text{with} \quad \boldsymbol{B} := \begin{pmatrix} -\boldsymbol{T}_1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \boldsymbol{C} := \begin{pmatrix} 0 & -\boldsymbol{T}_2^{1/2} \\ \boldsymbol{T}_2^{1/2} & 0 \end{pmatrix}.
$$

Then, applying the classical Weyl's Theorem (see [\[20\]](#page-26-5)), we have that  $Sp_{ess}(H)$  =  $Sp_{ess}(B)$  and the rest of the spectrum  $Sp_{disc}(H) := Sp(H) \backslash Sp_{ess}(H)$  consists of isolated eigenvalues with finite algebraic multiplicity. Moreover,  $Sp_{ess}(B)$  =  $Sp_{\textrm{ess}}(-T_1) \cup \{0\}.$ 

Our next goal is to show that the essential spectrum of  $T_1$  must lie in a small region of the complex plane. Actually, we will localize the whole spectrum of  $T_1$ . With this aim, we analyze separately for which values  $\mu \in \mathbb{C}$ , the operator  $(\mu I - T_1)$ is not necessarily one-to-one and for which values it is not necessarily onto.

• If  $(\mu I - T_1)$  is not one-to-one, then there exists  $f \in \mathcal{G}, f \neq 0$ , such that  $T \cdot f = \mu f$  pamely  $T_1f = \mu f$ , namely,

$$
\mu \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{f} \operatorname{div} \overline{\mathbf{v}} = 2 \int_{\Omega} \nu \operatorname{div} \mathbf{f} \operatorname{div} \overline{\mathbf{v}} \qquad \forall \mathbf{v} \in \mathcal{G}.
$$

Then, testing with  $v = f$  and using that in each subdomain the coefficients  $\rho$  and c are positive, we deduce that

$$
\mu = \frac{2 \int_{\Omega} \nu \left| \text{div} \, \boldsymbol{f} \right|^2}{\int_{\Omega} \rho c^2 \left| \text{div} \, \boldsymbol{f} \right|^2}
$$

(we recall that for  $0 \neq f \in \mathcal{G}$ ,  $\int_{\Omega} |\text{div } f|^2 > 0$  because of Lemma [2.2\)](#page-5-0). Hence,

$$
\mu \in \left[\frac{2\min_{\Omega}\{\nu\}}{\max_{\Omega}\{\rho c^2\}}, \frac{2\max_{\Omega}\{\nu\}}{\min_{\Omega}\{\rho c^2\}}\right].
$$

• On the other hand,  $(\mu I - T_1)$  is onto if and only if for any  $g \in \mathcal{G}$  there exists  $f \in \mathcal{G}$  such that  $T_1 f = \mu f - g$ , which from [\(3.1\)](#page-7-5) reads

$$
\int_{\Omega} \rho c^2 \operatorname{div} \mathbf{g} \operatorname{div} \overline{\mathbf{v}} = \int_{\Omega} (-2\nu + \mu \rho c^2) \operatorname{div} \mathbf{f} \operatorname{div} \overline{\mathbf{v}} \qquad \forall \mathbf{v} \in \mathcal{G}.
$$

By writing  $\mu = \alpha + \beta i$  with  $\alpha, \beta \in \mathbb{R}$ , the equation above reads:

$$
\int_{\Omega} (-2\nu + \alpha \rho c^2 + \rho c^2 \beta i) \operatorname{div} \mathbf{f} \operatorname{div} \overline{\mathbf{v}} = \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{g} \operatorname{div} \overline{\mathbf{v}} \qquad \forall \mathbf{v} \in \mathcal{G}.
$$

We observe that for all  $\beta \neq 0$  the problem above has a solution and hence the operator  $(\mu I - T_1)$  is onto. On the other hand, if  $\beta = 0$ , then  $\mu$  has to be real. In such a case, the operator  $T_1$  will still be onto when  $(-2\nu + \mu \rho c^2)$ has the same sign in the whole domain  $\Omega$ . This happens at least in two cases:

(i) when 
$$
\mu > \frac{2 \max_{\Omega} \{\nu\}}{\min_{\Omega} \{\rho c^2\}}
$$
, in which case  $-2\nu + \mu \rho c^2 > 0$ , (ii) when  $\mu < \frac{2 \min_{\Omega} \{\nu\}}{\max_{\Omega} \{\rho c^2\}}$ , in which case  $-2\nu + \mu \rho c^2 < 0$ .

Therefore, if  $(\mu I - T_1)$  is not onto, then  $\mu \in \left[\frac{2 \min_{\Omega} \{\nu\}}{\max_{\Omega} {\{\rho c^2\}}} \frac{2 \max_{\Omega} {\{\nu\}}}{\min_{\Omega} {\{\rho c^2\}}} \right]$  $\Big]$ , too.

Now we are in position to write the following spectral characterization of the solution operator T.

<span id="page-9-0"></span>**Theorem 3.1.** The spectrum of  $T$  consists of

$$
\mathrm{Sp}_{\mathrm{ess}}(\boldsymbol{T})=\mathrm{Sp}(\boldsymbol{T}_1)\cup\{0\}
$$

with

$$
\mathrm{Sp}(\boldsymbol{T}_1)\subset \left[\frac{2\min_{\Omega}\{\nu\}}{\max_{\Omega}\{\rho c^2\}}, \frac{2\max_{\Omega}\{\nu\}}{\min_{\Omega}\{\rho c^2\}}\right]
$$

and  $\text{Sp}_{disc}(T) := \text{Sp}(T) \setminus \text{Sp}_{ess}(T)$ , which is a set of isolated eigenvalues of finite algebraic multiplicity.

Proof. As a consequence of the classical Weyl's Theorem (see [\[20\]](#page-26-5)) and Lemma [3.2,](#page-8-0)

$$
Sp_{ess}(\boldsymbol{T}) = Sp_{ess}(\boldsymbol{H}) = Sp_{ess}(\boldsymbol{B}) = Sp_{ess}(-\boldsymbol{T}_1) \cup \{0\},
$$

whereas the inclusion follows from the above analyis.  $\hfill \Box$ 

In what follows, we will show that for  $\nu$  small enough some of the eigenvalues of **T** are well separated from its essential spectrum. With this end, given  $f \in \mathcal{G}$ , by testing [\(3.1\)](#page-7-5) with  $v = u_1 \in \mathcal{G}$  and using [\(2.15\)](#page-6-4), we have that

$$
\min_{\Omega} \{ \rho c^2 \} ||u_1||_{\text{div},\Omega}^2 \le a(u_1, u_1) \le 2||\nu||_{\infty,\Omega} ||\text{ div } f||_{0,\Omega} ||u_1||_{\text{div},\Omega}.
$$

Therefore  $||T_1||_{\mathcal{L}(\mathcal{G}\times\mathcal{G})}\to 0$  as  $||\nu||_{\infty,\Omega}$  goes to zero. Consequently, H converges in norm to the operator

$$
\boldsymbol{H}_0:=\begin{pmatrix}0&-{\boldsymbol{T}}_2^{1/2}\\{\boldsymbol{T}}_2^{1/2}&0\end{pmatrix}
$$

as  $\|\nu\|_{\infty,\Omega}$  goes to zero. Thus, from the classical spectral approximation theory (see [\[9\]](#page-25-10)), the isolated eigenvalues of  $H$  converge to those of  $H_0$ .

Since the isolated eigenvalues of  $H$  and  $T$  coincide (cf. Lemma [3.2\)](#page-8-0), in order to localize those of T, we begin by characterizing those of  $H_0$ . Let  $\mu$  be an isolated eigenvalue of  $H_0$  and  $(u,\hat{u}) \in \mathcal{G} \times \mathcal{G}$  the corresponding eigenfunction. It is easy to check that

<span id="page-10-1"></span>(3.3) 
$$
H_0\begin{pmatrix}u\\ \hat{u}\end{pmatrix}=\mu\begin{pmatrix}u\\ \hat{u}\end{pmatrix} \iff T_2u=-\mu^2u \text{ and } T_2^{1/2}u=\mu\hat{u}.
$$

Since  $T_2$  is compact, self-adjoint, and positive, its spectrum consists of a sequence of positive eigenvalues that converge to zero and 0 itself. Notice that the spectrum of  $T_2$  is related with the solution of the eigenvalue problem [\(2.11\)](#page-4-0). In fact, this problem has 0 as an eigenvalue with corresponding eigenspace  $K$ . The rest of the eigenvalues  $\omega^2$  are strictly positive and the corresponding eigenfunctions  $u \in$  $\mathcal{K}^{\perp_{\mathcal{V}}} =: \mathcal{G}$ , so that they are also solutions of the following problem: Find  $\omega > 0$  and  $u \in \mathcal{G}$  such that

$$
a(\boldsymbol{u},\boldsymbol{v}) = \omega^2 \int_{\Omega} \rho \boldsymbol{u} \cdot \overline{\boldsymbol{v}} \qquad \forall \boldsymbol{v} \in \mathcal{G}.
$$

Clearly  $(\omega^2, u)$  is an eigenpair of the above problem with  $\omega > 0$  if and only if  $T_2u = \frac{1}{\omega^2}u$ . Thus, by virtue of [\(3.3\)](#page-10-1), we have that the eigenvalues of  $H_0$  are given by  $\pm i/\omega$  and hence they are purely imaginary.

Now we are in a position to establish the following result.

**Theorem 3.2.** For each isolated eigenvalue  $\pm i/\omega$  of  $T_2$  of algebraic multiplicity m, let  $r > 0$  be such that the disc  $D_r := \{z \in \mathbb{C} : |z \pm i/\omega| < r\}$  intersects  $\text{Sp}(T_2)$  only in  $\pm i/\omega$ . Then, there exists  $\delta > 0$  such that if  $||\nu||_{\infty,\Omega} < \delta$ , there exist m eigenvalues of  $T$ ,  $\mu_1, \ldots, \mu_m$ , (repeated according to their respective algebraic multiplicities) lying in the disc  $D_r$ . Moreover,  $\mu_1, \ldots, \mu_m \to \frac{i}{\omega}$  as  $\|\nu\|_{\infty,\Omega}$  goes to zero.

As claimed above, the eigenvalues of  $T$  that are relevant in the applications, are those which are close to  $\pm i/\omega$  for the smallest positive vibration frequencies  $\omega$  of [\(2.11\)](#page-4-0). According to the above theorem, these eigenvalues are well separated from the real axis and, hence, from the essential spectrum of  $T$  (cf. Theorem [3.1\)](#page-9-0).

### 4. Spectral Approximation

<span id="page-10-0"></span>In this section, we propose and analyze a finite element method to approximate the solutions of Problem [1.](#page-3-5) With this end, we introduce appropriate discrete spaces. Let  ${\{\mathcal{T}_h(\Omega)\}_{h>0}}$  be a family of regular partitions of  $\Omega$  such that  $\mathcal{T}_h(\Omega_i) := \{T \in$  $\mathcal{T}_h : T \subset \Omega_i$  are partitions of  $\Omega_i$ ,  $i = 1, 2$ . We introduce the lowest-order Raviart-Thomas finite element space:

$$
\mathcal{V}_h:=\{\boldsymbol{v}\in\mathcal{V}: \boldsymbol{v}|_T(\boldsymbol{x})=\boldsymbol{a}+b\boldsymbol{x},\ \boldsymbol{a}\in\mathbb{R}^d,\ b\in\mathbb{R},\ \boldsymbol{x}\in T\}.
$$

The discretization of Problem [1](#page-3-5) reads as follows:

### 12 FELIPE LEPE, SALIM MEDDAHI, DAVID MORA, AND RODOLFO RODRÍGUEZ

<span id="page-11-0"></span>**Problem 3.** Find  $\lambda_h \in \mathbb{C}$  and  $\mathcal{Q} \neq \mathbf{u}_h \in \mathcal{V}_h$  such that

$$
\lambda_h^2 \int_{\Omega} \rho \mathbf{u}_h \cdot \overline{\mathbf{v}}_h + 2\lambda_h \int_{\Omega} \nu \operatorname{div} \mathbf{u}_h \operatorname{div} \overline{\mathbf{v}}_h + \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u}_h \operatorname{div} \overline{\mathbf{v}}_h = 0 \qquad \forall \mathbf{v}_h \in \mathcal{V}_h.
$$

We proceed as we did in the continuous case and introduce a new discrete variable  $\hat{u}_h := \lambda_h u_h$  to rewrite the problem above in the following equivalent form:

<span id="page-11-1"></span>**Problem 4.** Find  $\lambda_h \in \mathbb{C}$  and  $\mathcal{Q} \neq (\mathbf{u}_h, \widehat{\mathbf{u}}_h) \in \mathcal{V}_h \times \mathcal{V}_h$  such that

$$
\int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u}_h \operatorname{div} \overline{\boldsymbol{v}}_h = \lambda_h \left( -2 \int_{\Omega} \nu \operatorname{div} \boldsymbol{u}_h \operatorname{div} \overline{\boldsymbol{v}}_h - \int_{\Omega} \rho \widehat{\boldsymbol{u}}_h \cdot \overline{\boldsymbol{v}}_h \right) \qquad \forall \boldsymbol{v}_h \in \mathcal{V}_h,
$$
\n
$$
\int_{\Omega} \rho \widehat{\boldsymbol{u}}_h \cdot \overline{\widehat{\boldsymbol{v}}}_h = \lambda_h \int_{\Omega} \rho \boldsymbol{u}_h \cdot \overline{\widehat{\boldsymbol{v}}}_h \qquad \forall \widehat{\boldsymbol{v}}_h \in \mathcal{V}_h.
$$

We observe that  $\lambda_h = 0$  is an eigenvalue of this problem and its associated eigenspace is  $\tilde{\mathcal{K}}_h := \mathcal{K}_h \times \{0\}$  with  $\mathcal{K}_h := \mathcal{K} \cap \mathcal{V}_h$  being the eigenspace of  $\lambda_h = 0$ in Problem [3.](#page-11-0) At the beginning of Section [5,](#page-19-0) we will show that Problem [4](#page-11-1) is well posed, in the sense that it is equivalent to a generalized matrix eigenvalue problem with a symmetric positive definite right-hand side matrix.

We introduce the well known Raviart-Thomas interpolation operator,  $\Pi_h : \mathcal{V} \cap$  $H^{r}(\Omega)^{d} \to V_{h}, r \in (0,1]$  (see [\[13\]](#page-25-11)), for which there holds the approximation result

(4.1) 
$$
\|v - \Pi_h v\|_{0,\Omega} \leq Ch^r(\|v\|_{r,\Omega} + \|\operatorname{div} v\|_{0,\Omega})
$$

and the commuting diagram property

(4.2) 
$$
\operatorname{div}(\Pi_h \boldsymbol{v}) = \mathcal{P}_h(\operatorname{div} \boldsymbol{v}),
$$

where

<span id="page-11-5"></span><span id="page-11-3"></span><span id="page-11-2"></span>
$$
\mathcal{P}_h: \mathcal{L}^2(\Omega) \to \mathcal{U}_h := \{ v_h \in \mathcal{L}^2(\Omega) : \ v_h|_T \in \mathcal{P}_0(T) \ \forall T \in \mathcal{T}_h \}
$$

is the standard L<sup>2</sup>-orthogonal projector. Then, for any  $r \in (0, 1]$  we have that

(4.3) 
$$
||q - \mathcal{P}_h q||_{0,\Omega} \leq Ch^r ||q||_{r,\Omega} \quad \forall q \in H^r(\Omega).
$$

Let  $\mathcal{G}_h$  be the orthogonal complement of  $\mathcal{K}_h$  in  $\mathcal{V}_h$ , and  $\widetilde{\mathcal{G}}_h := \mathcal{G}_h \times \mathcal{G}_h \subset \widetilde{\mathcal{V}} = \mathcal{V} \times \mathcal{H}$ endowed with the corresponding product norm. Note that  $\mathcal{G}_h \nsubseteq \mathcal{G}$  and hence  $\widetilde{\mathcal{G}}_h \nsubseteq \widetilde{\mathcal{G}}.$ 

The following result provides estimates for the terms in the Helmholtz decomposition of functions in  $\mathcal{G}_h$ .

<span id="page-11-4"></span>**Lemma 4.1.** For any  $v_h \in \mathcal{G}_h$ ,

$$
\boldsymbol{v}_h = \frac{1}{\rho} \nabla \xi + \chi
$$

with  $\frac{1}{\rho} \nabla \xi \in \mathcal{H}^s(\Omega)^d$  and  $\chi \in \mathcal{K}$  satisfying

$$
\left\|\frac{1}{\rho}\nabla\xi\right\|_{s,\Omega}\leq C\|\operatorname{div}\boldsymbol{v}_h\|_{0,\Omega}\qquad\text{and}\qquad\|\chi\|_{0,\Omega}\leq Ch^s\|\operatorname{div}\boldsymbol{v}_h\|_{0,\Omega}.
$$

*Proof.* Let  $\xi \in H^1(\Omega)/\mathbb{C}$  be a solution of the following well-posed Neumann problem:

$$
\operatorname{div}\left(\frac{1}{\rho}\nabla\xi\right) = \operatorname{div}\boldsymbol{v}_h \quad \text{in } \Omega,
$$

$$
\frac{1}{\rho}\frac{\partial\xi}{\partial\mathbf{n}} = 0 \qquad \text{on } \partial\Omega.
$$

Thanks to Lax-Milgram's Lemma, there exists a unique solution  $\xi \in H^1(\Omega)/\mathbb{C}$ of this problem. Moreover, according to Lemmas [2.1](#page-4-1) and [2.2,](#page-5-0)  $\frac{1}{\rho} \nabla \xi \in H^s(\Omega)^d$  and  $\|\frac{1}{\rho}\nabla\xi\|_{s,\Omega} \leq C \|\mathop{\mathrm{div}}\nolimits \boldsymbol{v}_h\|_{0,\Omega}$ . Now, let  $\chi := \boldsymbol{v}_h - \frac{1}{\rho}\nabla \xi$ . Clearly  $\mathop{\mathrm{div}}\nolimits \chi = 0$  and  $\chi \cdot \mathbf{n} = 0$ , so that  $\chi \in \mathcal{K}$ . On the other hand

$$
\|\chi\|_{\mathcal{H}}^2 = \int_{\Omega} \rho \chi \cdot \left(\mathbf{v}_h - \frac{1}{\rho} \nabla \xi\right).
$$

Since  $\frac{1}{\rho} \nabla \xi \in \mathcal{V} \cap \mathrm{H}^s(\Omega)^d$ , we have that  $\Pi_h(\frac{1}{\rho} \nabla \xi)$  is well defined. Hence,

$$
\|\chi\|_{\mathcal{H}}^2 = \underbrace{\int_{\Omega} \rho \chi \cdot \left[\boldsymbol{v}_h - \Pi_h \left(\frac{1}{\rho} \nabla \xi\right)\right]}_{(I)} + \underbrace{\int_{\Omega} \rho \chi \cdot \left[\Pi_h \left(\frac{1}{\rho} \nabla \xi\right) - \frac{1}{\rho} \nabla \xi\right]}_{(II)}.
$$

For (I), thanks to [\(4.2\)](#page-11-2),  $\text{div}(\mathbf{v}_h - \Pi_h(\frac{1}{\rho}\nabla \xi)) = \text{div}\,\mathbf{v}_h - \mathcal{P}_h(\text{div}(\frac{1}{\rho}\nabla \xi)) = 0$ . Therefore,  $(\boldsymbol{v}_h - \Pi_h(\frac{1}{\rho}\nabla \xi)) \in \mathcal{K}_h \subset \mathcal{K}$ . Since  $\frac{1}{\rho}\nabla \xi \in \mathcal{G}$  and  $\boldsymbol{v}_h \in \mathcal{G}_h$ , we obtain

$$
(4.4) (I) = \int_{\Omega} \rho v_h \cdot \left[ v_h - \Pi_h \left( \frac{1}{\rho} \nabla \xi \right) \right] - \int_{\Omega} \rho \left( \frac{1}{\rho} \nabla \xi \right) \cdot \left[ v_h - \Pi_h \left( \frac{1}{\rho} \nabla \xi \right) \right] = 0.
$$
  
For (II) since we have already proved that  $||^1 \nabla \xi|| \leq C || \dim ||$ 

For  $(II)$ , since we have already proved that  $\|\frac{1}{\rho}\nabla\xi\|_{s,\Omega} \leq C \|\mathop{\rm div} \mathbf{v}_h\|_{0,\Omega}$  and  $\operatorname{div}(\frac{1}{\rho}\nabla \xi) = \operatorname{div} \boldsymbol{v}_h$ , from [\(4.1\)](#page-11-3) we obtain

$$
(II) \leq \|\chi\|_{0,\Omega} \left\| \Pi_h \left( \frac{1}{\rho} \nabla \xi \right) - \frac{1}{\rho} \nabla \xi \right\|_{0,\Omega} \leq C h^s \|\chi\|_{0,\Omega} \|\operatorname{div} \boldsymbol{v}_h\|_{0,\Omega},
$$

which allows us to complete the proof.

Now, we will prove that  $a(\cdot, \cdot)$  and  $\widetilde{a}(\cdot, \cdot)$  are elliptic in  $\mathcal{G}_h$  and  $\widetilde{\mathcal{G}}_h$ , respectively.

**Lemma 4.2.** The sesquilinear form  $a : \mathcal{G}_h \times \mathcal{G}_h \to \mathbb{C}$  is  $\mathcal{G}_h$ -elliptic, with ellipticity constant not depending on h. Consequently,  $\tilde{a}$  :  $\tilde{G}_h \times \tilde{G}_h \to \mathbb{C}$  is  $\tilde{G}_h$ -elliptic uniformly in h.

*Proof.* Let  $v_h \in \mathcal{G}_h$ . We have that

$$
a(\boldsymbol{v}_h, \boldsymbol{v}_h) = \int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{v}_h \operatorname{div} \overline{\boldsymbol{v}}_h \ge \min_{\Omega} \{ \rho c^2 \} || \operatorname{div} \boldsymbol{v}_h ||_{0,\Omega}^2.
$$

Now, from Lemma [4.1](#page-11-4) we write  $\mathbf{v}_h = \frac{1}{\rho} \nabla \xi + \chi$  with  $\frac{1}{\rho} \nabla \xi \in H^s(\Omega)^d$  and  $\chi \in \mathcal{K}$ . Then, using Lemma [4.1](#page-11-4) again we obtain

$$
\|\boldsymbol{v}_h\|_{0,\Omega}\leq \left\|\frac{1}{\rho}\nabla \xi\right\|_{0,\Omega}+\|\chi\|_{0,\Omega}\leq C\|\operatorname{div}\boldsymbol{v}_h\|_{0,\Omega},
$$

which together with the previous inequality allow us to conclude that  $a(\cdot, \cdot)$  is  $\mathcal{G}_h$ elliptic. The  $\mathcal{G}_h$ -ellipticity of  $\tilde{a}(\cdot, \cdot)$  is a direct consequence of the  $\mathcal{G}_h$ -ellipticity of  $a(\cdot, \cdot)$ .  $a(\cdot, \cdot)$ .

Now, we are in position to introduce the discrete version of the operator  $T$ . Let  $T_h : \widetilde{V} \to \widetilde{V}$  be defined by  $T_h(f,g) := (u_h,\widehat{u}_h)$  with  $(u_h,\widehat{u}_h) \in \widetilde{\mathcal{G}}_h$  being the solution of

<span id="page-12-0"></span>
$$
\widetilde{a}((\boldsymbol{u}_h,\widehat{\boldsymbol{u}}_h),(\boldsymbol{v}_h,\widehat{\boldsymbol{v}}_h))=\widetilde{b}((\boldsymbol{f},\boldsymbol{g}),(v_h,\widehat{\boldsymbol{v}}_h))\qquad\forall (\boldsymbol{v}_h,\widehat{\boldsymbol{v}}_h)\in \widetilde{\mathcal{G}}_h.
$$

It is easy to check that  $(u_h,\hat{u}_h) = T_h(f,g)$  if and only if

$$
\widehat{\boldsymbol{u}}_h = \mathcal{P}_{\mathcal{G}_h} \boldsymbol{f},
$$

$$
\Box
$$

where  $\mathcal{P}_{\mathcal{G}_h}$  is the H-orthogonal projection onto  $\mathcal{G}_h$ , and  $\mathbf{u}_h \in \mathcal{G}_h$  solves

<span id="page-13-4"></span>(4.6) 
$$
\int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u}_h \operatorname{div} \boldsymbol{v}_h = -2 \int_{\Omega} \nu \operatorname{div} \boldsymbol{f} \operatorname{div} \overline{\boldsymbol{v}}_h - \int_{\Omega} \rho \boldsymbol{g} \cdot \overline{\boldsymbol{v}}_h \qquad \forall \boldsymbol{v}_h \in \mathcal{G}_h.
$$

Since  $T_h(\mathcal{V}) \subset \mathcal{G}_h$ , there holds  $\text{Sp}(T_h) = \text{Sp}(T_h |_{\widetilde{\mathcal{G}}_h}) \cup \{0\}$  (cf. [\[1,](#page-25-12) Lemma 4.1]). Thus, we will restrict our attention to  $T_h|_{\widetilde{\mathcal{G}}_h}$ .

As claimed above, at the beginning of Section [5,](#page-19-0) Problem [4](#page-11-1) will be shown to be equivalent to a well posed generalized matrix eigenvalue problem. This problem has  $\lambda_h = 0$  as an eigenvalue with corresponding eigenspace  $\mathcal{K}_h$ . The rest of the eigenvalues are related with the spectrum of  $T_h|_{\widetilde{\mathcal{G}}_h}$  according to the following lemma.

**Lemma 4.3.** There holds that  $(\mu_h, (\boldsymbol{u}_h, \widehat{\boldsymbol{u}}_h))$  is an eigenpair of  $\boldsymbol{T}_h|_{\widetilde{\mathcal{G}}_h}$  with  $\mu_h \neq 0$ if and only if  $(\lambda_h,(\boldsymbol{u}_h,\widehat{\boldsymbol{u}}_h))$  is a solution of Problem [4](#page-11-1) with  $\lambda_h = 1/\mu_h$ .

Proof. The proof follows essentially as that of Lemma [2.5,](#page-7-3) by using the fact that  $\mathcal{V}_h \times \mathcal{V}_h = \widetilde{\mathcal{G}}_h \oplus (\mathcal{K}_h \times \mathcal{K}_h).$ 

Our next goal is to show that any isolated eigenvalue of  $T$  with algebraic multiplicity m is approximated by exactly m eigenvalues of  $T_h$  (repeated according to their respective algebraic multiplicities) and that spurious eigenvalues do not arise. With this end, we will adapt to our problem the theory from [\[2\]](#page-25-2), which in turn use arguments introduced in [\[7,](#page-25-8) [8\]](#page-25-9) to deal with non compact operators. From now on, let  $\mu \in \mathrm{Sp}_{disc}(\mathbf{T}), \mu \neq 0$ , be a fixed isolated eigenvalue of finite algebraic multiplicity m. Let  $\mathcal E$  be the invariant subspace of  $T$  corresponding to  $\mu$ . Our analysis will be based on proving the following two properties:

P1. 
$$
\|\boldsymbol{T} - \boldsymbol{T}_h\|_h := \sup_{\substack{\boldsymbol{\mathcal{Q}} \neq (\boldsymbol{f}_h, \boldsymbol{g}_h) \in \widetilde{\mathcal{G}}_h}} \frac{\|(\boldsymbol{T} - \boldsymbol{T}_h)(\boldsymbol{f}_h, \boldsymbol{g}_h)\|_{\widetilde{\mathcal{V}}}}{\|(\boldsymbol{f}_h, \boldsymbol{g}_h)\|_{\widetilde{\mathcal{V}}}} \to 0 \text{ as } h \to 0;
$$
  
P2. 
$$
\forall (\boldsymbol{v}, \widehat{\boldsymbol{v}}) \in \mathcal{E}, \inf_{(\boldsymbol{v}_h, \widehat{\boldsymbol{v}}_h) \in \widetilde{\mathcal{G}}_h} \|(\boldsymbol{v}, \widehat{\boldsymbol{v}}) - (\boldsymbol{v}_h, \widehat{\boldsymbol{v}}_h)\|_{\widetilde{\mathcal{V}}} \to 0 \text{ as } h \to 0.
$$

Let  $(f_h, g_h) \in \mathcal{G}_h$  and  $(u, \hat{u}) := T(f_h, g_h)$ . From [\(2.17\)](#page-7-6), we can write  $u =$  $u_1 + u_2$  with  $u_1, u_2 \in \mathcal{G}$  satisfying

<span id="page-13-0"></span>(4.7) 
$$
\mathbf{u}_1 \in \mathcal{G}: \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u}_1 \operatorname{div} \overline{\mathbf{v}} = -2 \int_{\Omega} \nu \operatorname{div} \mathbf{f}_h \operatorname{div} \overline{\mathbf{v}} \quad \forall \mathbf{v} \in \mathcal{G},
$$

and

(4.8) 
$$
\mathbf{u}_2 \in \mathcal{G}: \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u}_2 \operatorname{div} \overline{\mathbf{v}} = -\int_{\Omega} \rho \mathbf{g}_h \cdot \overline{\mathbf{v}} \quad \forall \mathbf{v} \in \mathcal{G}.
$$

<span id="page-13-1"></span>The following result states some properties of the solutions of the problems above.

<span id="page-13-2"></span>**Lemma 4.4.** For  $(f_h, g_h) \in \widetilde{\mathcal{G}}_h$ , let  $(u, \widehat{u}) := T(f_h, g_h)$  and consider the decomposition  $\mathbf{u} = \mathbf{u}_1 + \mathbf{u}_2$  as above. Hence,  $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{H}^s(\Omega)^d$ , div  $\mathbf{u}_1 \in \mathcal{U}_h$ ,  $\text{div } u_2|_{\Omega_i} \in H^{1+s}(\Omega_i)$ ,  $i = 1, 2$ , and the following estimates hold

<span id="page-13-3"></span>(4.9) ku1ks,<sup>Ω</sup> ≤ Ck(f<sup>h</sup> , g<sup>h</sup> )kV<sup>e</sup>,

(4.10)  $\|u_2\|_{s,\Omega} + \|\text{div } u_2\|_{1+s,\Omega_1} + \|\text{div } u_2\|_{1+s,\Omega_2} \leq C \|(f_h, g_h)\|_{\widetilde{\mathcal{V}}}$ 

*Proof.* Since  $u_1 \in \mathcal{G}$ , due to Lemma [2.2](#page-5-0) we have that  $u_1 \in H^s(\Omega)^d$  and  $||u_1||_{s,\Omega} \le$  $C\|(f_h, g_h)\|_{\widetilde{\mathcal{V}}}$ . Moreover, note that [\(4.7\)](#page-13-0) also holds for  $v \in \mathcal{K}$  and hence for all  $v \in V$ . Then, we write

$$
\int_{\Omega} (\rho c^2 \operatorname{div} \boldsymbol{u}_1 + 2\nu \operatorname{div} \boldsymbol{f}_h) \operatorname{div} \boldsymbol{\overline{v}} = 0 \qquad \forall \boldsymbol{v} \in \mathcal{V}.
$$

Thus, taking test functions in  $\mathcal{D}(\Omega)^d \subset \mathcal{V}$  we have  $\nabla(\rho c^2 \operatorname{div} \mathbf{u}_1 + 2\nu \operatorname{div} \mathbf{f}_h) = 0$ . Since  $\rho, c, \nu$  and div  $f_h$  are piecewise constant, we have that div  $u_1$  is piecewise constant as well; namely, div  $u_1 \in \mathcal{U}_h$ .

On the other hand, since  $u_2 \in \mathcal{G}$ , by applying Lemma [2.2](#page-5-0) again we have that  $u_2 \in H^s(\Omega)^d$  and  $||u_2||_{s,\Omega} \leq C ||(f_h, g_h)||_{\widetilde{\mathcal{V}}}$ . To prove additional regularity for div  $u_2$ , we use Lemma [4.1](#page-11-4) to write  $g_h = \frac{1}{\rho} \nabla \xi + \chi$  with  $\chi \in \mathcal{K}, \frac{1}{\rho} \nabla \xi \in H^s(\Omega)^d$  and  $\|\frac{1}{\rho}\nabla\xi\|_{s,\Omega} \leq C \|\mathop{\mathrm{div}} \mathbf{g}_h\|_{0,\Omega}$ . Moreover, since  $\rho$  is constant in each subdomain  $\Omega_i$ , also  $\nabla \xi |_{\Omega_i} \in H^s(\Omega_i)^d$ ,  $i = 1, 2$ . Then, from [\(4.8\)](#page-13-1) we have that

$$
\int_{\Omega} \rho c^2 \operatorname{div} \boldsymbol{u}_2 \operatorname{div} \boldsymbol{\overline{v}} = -\int_{\Omega} \nabla \xi \cdot \boldsymbol{\overline{v}} \qquad \forall \boldsymbol{v} \in \mathcal{G}.
$$

Since the above equation trivially holds for  $v \in \mathcal{K}$  too, it holds for all  $v \in \mathcal{V}$ . Then, by testing it with  $\mathbf{v} \in \mathcal{D}(\Omega)^d$  we have that  $\nabla(\rho c^2 \text{ div } \mathbf{u}_2) = -\nabla \xi \in \Omega$ . Therefore, by restricting to  $\Omega_i$ ,  $i = 1, 2$ , we have that  $\nabla (\rho c^2 \operatorname{div} \boldsymbol{u}_2 |_{\Omega_i}) = -\nabla (\xi |_{\Omega_i}) \in \mathrm{H}^s(\Omega_i)^d$ . Since  $\rho$  and c are piecewise constant, we conclude that div  $u_2|_{\Omega_i} \in H^{1+s}(\Omega_i)$ ,  $i =$ 1, 2, and

$$
\|\,\mathrm{div}\,\bm{u}_2\|_{1+s,\Omega_1} + \|\,\mathrm{div}\,\bm{u}_2\|_{1+s,\Omega_2} \leq C\|\nabla\xi\|_{0,\Omega} \leq C\|\,\mathrm{div}\,\bm{g}_h\|_{0,\Omega}.
$$

Hence, we conclude the proof.

We consider a similar decomposition in the discrete case. For 
$$
(\mathbf{f}_h, \mathbf{g}_h) \in \widetilde{\mathcal{G}}_h
$$
, let  $(\mathbf{u}_h, \widehat{\mathbf{u}}_h) := \mathbf{T}_h(\mathbf{f}_h, \mathbf{g}_h)$ . We write  $\mathbf{u}_h = \mathbf{u}_{1h} + \mathbf{u}_{2h}$  with  $\mathbf{u}_{1h}$  and  $\mathbf{u}_{2h}$  satisfying  
(4.11)  $\mathbf{u}_{1h} \in \mathcal{G}_h$ :  $\int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u}_{1h} \operatorname{div} \overline{\mathbf{v}}_h = -2 \int_{\Omega} \nu \operatorname{div} \mathbf{f}_h \operatorname{div} \overline{\mathbf{v}}_h \qquad \forall \mathbf{v}_h \in \mathcal{G}_h$ ,

<span id="page-14-0"></span>and

<span id="page-14-1"></span>(4.12) 
$$
\mathbf{u}_{2h} \in \mathcal{G}_h: \quad \int_{\Omega} \rho c^2 \operatorname{div} \mathbf{u}_{2h} \operatorname{div} \overline{\mathbf{v}}_h = -\int_{\Omega} \rho \mathbf{g}_h \cdot \overline{\mathbf{v}}_h \qquad \forall \mathbf{v}_h \in \mathcal{G}_h.
$$

These are the finite element discretization of problems [\(4.7\)](#page-13-0) and [\(4.8\)](#page-13-1), respectively, and the following error estimates hold true.

<span id="page-14-5"></span>**Lemma 4.5.** Let  $(f_h, g_h) \in \widetilde{\mathcal{G}}_h$ . Let  $u_1, u_2$  be the solutions of problems [\(4.7\)](#page-13-0) and [\(4.8\)](#page-13-1), respectively, and  $u_{1h}, u_{2h}$  those of problems [\(4.11\)](#page-14-0) and [\(4.12\)](#page-14-1), respectively. Then, the following estimates hold true:

<span id="page-14-3"></span>(4.13) 
$$
\|\boldsymbol{u}_1-\boldsymbol{u}_{1h}\|_{\text{div},\Omega}\leq Ch^s\|(\boldsymbol{f}_h,\boldsymbol{g}_h)\|_{\widetilde{\mathcal{V}}},
$$

<span id="page-14-4"></span>(4.14) 
$$
\|u_2-u_{2h}\|_{\text{div},\Omega}\leq Ch^s \|(f_h,g_h)\|_{\widetilde{\mathcal{V}}}.
$$

*Proof.* Since  $\mathcal{G}_h \nsubseteq \mathcal{G}$ , we will resort to the second Strang Lemma, which for problems [\(4.7\)](#page-13-0) and [\(4.11\)](#page-14-0) reads as follows:

<span id="page-14-2"></span>
$$
(4.15) \t ||\boldsymbol{u}_1 - \boldsymbol{u}_{1h}||_{\text{div},\Omega} \leq C \left[\inf_{\boldsymbol{v}_h \in \mathcal{G}_h} \|\boldsymbol{u}_1 - \boldsymbol{v}_h\|_{\text{div},\Omega} + \sup_{\underline{0} \neq \boldsymbol{v}_h \in \mathcal{G}_h} \frac{a(\boldsymbol{u}_1 - \boldsymbol{u}_{1h}, \boldsymbol{v}_h)}{\|\boldsymbol{v}_h\|_{\text{div},\Omega}}\right].
$$

$$
\Box
$$

Because of Lemma [4.4,](#page-13-2)  $\Pi_h \mathbf{u}_1$  is well defined. Since  $\Pi_h \mathbf{u}_1 \in \mathcal{V}_h = \mathcal{G}_h \oplus \mathcal{K}_h$ , there exists  $\tilde{\mathbf{u}}_{1h} \in \mathcal{G}_h$  and  $\tilde{\mathbf{u}}_h \in \mathcal{K}_h$  such that  $\Pi_h \mathbf{u}_1 = \tilde{\mathbf{u}}_{1h} + \tilde{\mathbf{u}}_h$ . Then, since  $\mathbf{u}_1 - \tilde{\mathbf{u}}_{1h}$  is orthogonal to  $\check{u}_h$ , we observe that

$$
\|u_1 - \widetilde{u}_{1h}\|_{\mathcal{V}}^2 \le \|u_1 - \widetilde{u}_{1h}\|_{\mathcal{V}}^2 + \|\breve{u}_h\|_{\mathcal{V}}^2 = \|(\widetilde{u}_{1h} - u_1) + \breve{u}_h\|_{\mathcal{V}}^2 = \|u_1 - \Pi_h u_1\|_{\mathcal{V}}^2 \le C \left( \|u_1 - \Pi_h u_1\|_{0,\Omega}^2 + \|\operatorname{div} u_1 - \operatorname{div}(\Pi_h u_1)\|_{0,\Omega}^2 \right).
$$

The first term on the right hand side above is bounded as follows:

 $||u_1 - \Pi_h u_1||_{0,\Omega} \le Ch^s(||u_1||_{s,\Omega} + ||\text{ div } u_1||_{0,\Omega}) \le Ch^s||(f_h, g_h)||_{\widetilde{\mathcal{V}}}$ 

where we have used [\(4.1\)](#page-11-3), [\(4.9\)](#page-13-3), and the fact that  $\|\text{div }u_1\|_{0,\Omega} \leq C \|\text{div }f_h\|_{0,\Omega}$ , which in turn follows from [\(4.7\)](#page-13-0) by taking  $\mathbf{v} = \mathbf{f}_h$ . On the other hand, the second term vanishes because of [\(4.2\)](#page-11-2) since div  $u_1 \in \mathcal{U}_h$  (cf. Lemma [4.4\)](#page-13-2). Hence,  $||u_1 \tilde{\mathbf{u}}_{1h}$   $\|\text{div}, \Omega \leq Ch^s \|(\boldsymbol{f}_h, \boldsymbol{g}_h)\|_{\tilde{\mathcal{V}}}$ , which allows us to control the approximation term in  $(4.15).$  $(4.15).$ 

For the consistency term, it is enough to recall that [\(4.7\)](#page-13-0) holds for all  $v \in V$ . Then, by using [\(4.11\)](#page-14-0), it is easy to check that  $a(\mathbf{u}_1 - \mathbf{u}_{1h}, \mathbf{v}_h) = 0$  for all  $\mathbf{v}_h \in \mathcal{G}_h \subset$ V. From this, the Strang estimate for  $||u_1 - u_{1h}||_{div,\Omega}$  reads as follows:

$$
\|\boldsymbol{u}_1-\boldsymbol{u}_{1h}\|_{\mathrm{div},\Omega}\leq C\inf\limits_{\boldsymbol{v}_h\in\mathcal{G}_h}\|\boldsymbol{u}_1-\boldsymbol{v}_h\|_{\mathrm{div},\Omega}\leq Ch^s\|(\boldsymbol{f}_h,\boldsymbol{g}_h\|_{\widetilde{\mathcal{V}}}.
$$

Thus [\(4.13\)](#page-14-3) holds true.

<span id="page-15-0"></span>To prove [\(4.14\)](#page-14-4), we resort again to the second Strang Lemma:

$$
(4.16)\t||\boldsymbol{u}_2-\boldsymbol{u}_{2h}\|_{\text{div},\Omega}\leq C\left[\inf_{\boldsymbol{v}_h\in\mathcal{G}_h}\|\boldsymbol{u}_2-\boldsymbol{v}_h\|_{\text{div},\Omega}+\sup_{\underline{0}\neq\boldsymbol{v}_h\in\mathcal{G}_h}\frac{a(\boldsymbol{u}_2-\boldsymbol{u}_{2h},\boldsymbol{v}_h)}{\|\boldsymbol{v}_h\|_{\text{div},\Omega}}\right].
$$

Since  $u_2 \in H^s(\Omega)^d$  (cf. Lemma [4.4\)](#page-13-2), we have that  $\Pi_h u_2$  is well defined. We proceed as above and write  $\Pi_h u_2 = \tilde{u}_{2h} + \tilde{u}_h$  with  $\tilde{u}_{2h} \in \mathcal{G}_h$  and  $\tilde{u}_h \in \mathcal{K}_h$  to obtain

$$
(4.17) \t ||\boldsymbol{u}_2 - \widetilde{\boldsymbol{u}}_{2h}||_{\text{div},\Omega} \leq C \left[\|\boldsymbol{u}_2 - \Pi_h \boldsymbol{u}_2||_{0,\Omega} + \|\text{div}\,\boldsymbol{u}_2 - \text{div}(\Pi_h \boldsymbol{u}_2)||_{0,\Omega}\right].
$$

For the first term on the right hand side above, [\(4.1\)](#page-11-3) and Lemma [4.4](#page-13-2) yield

$$
\|\bm{u}_2-\Pi_h\bm{u}_2\|_{0,\Omega}\leq Ch^s(\|\bm{u}_2\|_{s,\Omega}+\|\,\mathrm{div}\,\bm{u}_2\|_{0,\Omega})\leq Ch^s\|(\bm{f}_h,\bm{g}_h)\|_{\widetilde{\mathcal{V}}}.
$$

For the second term, we have from [\(4.3\)](#page-11-5) and from Lemma [4.4](#page-13-2) again

 $\|\operatorname{div}\boldsymbol{u}_2-\operatorname{div}\Pi_h\boldsymbol{u}_2\|_{0,\Omega}^2=\|\operatorname{div}\boldsymbol{u}_2-\mathcal{P}_h(\operatorname{div}\boldsymbol{u}_2)\|_{0,\Omega}^2,$ 

$$
\leq Ch(\|\mathop{\rm div}\nolimits\bm{u}_2\|_{1,\Omega_1}+\|\mathop{\rm div}\nolimits\bm{u}_2\|_{1,\Omega_2})\leq Ch\|(\bm{f}_h,\bm{g}_h)\|_{\widetilde{\mathcal{V}}}.
$$

Hence,  $\|\boldsymbol{u}_2 - \widetilde{\boldsymbol{u}}_{2h}\|_{\text{div},\Omega} \leq Ch^s \|(\boldsymbol{f}_h, \boldsymbol{g}_h)\|_{\widetilde{\mathcal{V}}}$ , which allows us to bound the approximation term in [\(4.16\)](#page-15-0).

For the consistency term, given  $v_h \in \mathcal{G}_h$  we apply Lemma [4.1](#page-11-4) to write  $v_h$  =  $\frac{1}{\rho} \nabla \xi + \chi$  with  $\frac{1}{\rho} \nabla \xi \in H^s(\Omega)^d$ ,  $\chi \in \mathcal{K}$ , and  $\|\chi\|_{0,\Omega} \le Ch^s \|\text{div} \, \boldsymbol{v}_h\|_{0,\Omega}$ . Then, from  $(4.8)$  we have

$$
a(\boldsymbol{u}_2,\boldsymbol{v}_h)=\int_{\Omega}\rho c^2 \operatorname{div}\boldsymbol{u}_2 \operatorname{div}\boldsymbol{\overline{v}}_h=\int_{\Omega}\rho c^2 \operatorname{div}\boldsymbol{u}_2 \operatorname{div}\left(\frac{1}{\rho}\nabla\overline{\xi}\right)=\int_{\Omega}\boldsymbol{g}_h\cdot\nabla\overline{\xi}.
$$

On the other hand, from [\(4.12\)](#page-14-1),

$$
a(\boldsymbol{u}_{2h},\boldsymbol{v}_{h})=\int_{\Omega}\rho c^{2} \operatorname{div}\boldsymbol{u}_{2h} \operatorname{div}\overline{\boldsymbol{v}}_{h}=\int_{\Omega}\rho \boldsymbol{g}_{h}\cdot\overline{\boldsymbol{v}}_{h}=\int_{\Omega}\boldsymbol{g}_{h}\cdot\nabla \overline{\xi}+\int_{\Omega}\rho \boldsymbol{g}_{h}\cdot\overline{\chi}.
$$

Therefore,

$$
a(\boldsymbol{u}_2-\boldsymbol{u}_{2h},\boldsymbol{v}_h)=-\int_{\Omega}\rho \boldsymbol{g}_h\cdot \overline{\chi}\leq Ch^s\|\boldsymbol{g}_h\|_{0,\Omega}\|\boldsymbol{v}_h\|_{{\rm div},\Omega}
$$

and, hence,

$$
\sup_{\underline{0}\neq\boldsymbol{v}_h\in\mathcal{G}_h}\frac{a(\boldsymbol{u}_2-\boldsymbol{u}_{2h},\boldsymbol{v}_h)}{\|\boldsymbol{v}_h\|_{\text{div},\Omega}}\leq Ch^s\|\boldsymbol{g}_h\|_{0,\Omega},
$$

which allows us to complete the proof.  $\square$ 

Now, we are in a position to establish the following result.

<span id="page-16-5"></span>Lemma 4.6. Property P1 holds true. Moreover,

$$
||\boldsymbol{T}-\boldsymbol{T}_h||_h \leq Ch^s.
$$

Proof. For  $(f_h, g_h) \in \mathcal{G}_h$ , let  $(u, \hat{u}) := T(f_h, g_h)$  and  $(u_h, \hat{u}_h) := T_h(f_h, g_h)$ . From [\(2.16\)](#page-7-1) and [\(4.5\)](#page-12-0) we have that  $\hat{u} - \hat{u}_h = f_h - \mathcal{P}_{\mathcal{G}_h} f_h = 0$ . Hence, by writing  $u = u_1 + u_2$  and  $u_h = u_{1h} + u_{2h}$  as in Lemma [4.5,](#page-14-5) we have from this lemma

$$
\|\boldsymbol{T} - \boldsymbol{T}_h\|_h \le \sup_{\substack{\boldsymbol{0} \ne (\boldsymbol{g}_h, \boldsymbol{f}_h) \in \widetilde{\mathcal{G}}_h}} \frac{\|\boldsymbol{u}_1 - \boldsymbol{u}_{1h}\|_{\text{div},\Omega} + \|\boldsymbol{u}_2 - \boldsymbol{u}_{2h}\|_{\text{div},\Omega}}{\|(\boldsymbol{f}_h, \boldsymbol{g}_h)\|_{\widetilde{\mathcal{V}}}} \le Ch^s.
$$
\nThus, we conclude the proof.

Our next goal is to prove property P2. With this aim, first we will prove the following additional regularity result.

<span id="page-16-4"></span>**Lemma 4.7.** Let  $(\mathbf{u}, \widehat{\mathbf{u}}) \in \mathcal{E}$ . Then,  $\mathbf{u}, \widehat{\mathbf{u}} \in \mathcal{G} \subset \mathrm{H}^s(\Omega)^d$ , div  $\mathbf{u}, \mathrm{div} \widehat{\mathbf{u}} \in \mathrm{H}^{1+s}(\Omega_i)$ ,  $i = 1, 2, and$ 

<span id="page-16-0"></span>(4.18) 
$$
\|u\|_{s,\Omega}+\|\operatorname{div} u\|_{1+s,\Omega_1}+\|\operatorname{div} u\|_{1+s,\Omega_2}\leq C\|(u,\widehat{u})\|_{\widetilde{\mathcal{V}}},
$$

<span id="page-16-1"></span>
$$
(4.19) \t\t ||\widehat{\boldsymbol{u}}||_{s,\Omega} + ||\operatorname{div} \widehat{\boldsymbol{u}}||_{1+s,\Omega_1} + ||\operatorname{div} \widehat{\boldsymbol{u}}||_{1+s,\Omega_2} \leq C ||(\boldsymbol{u},\widehat{\boldsymbol{u}})||_{\widetilde{\mathcal{V}}}.
$$

Proof. We prove the above inequalities for all the generalized eigenfunctions of T. Let  $\{(\mathbf{u}_k, \hat{\mathbf{u}}_k)\}_{k=1}^p$  be a Jordan chain of the operator  $T$  associated with  $\mu$ . Then,  $T(u_k,\hat{u}_k) = \mu(u_k,\hat{u}_k) + (u_{k-1},\hat{u}_{k-1}), k = 1,\ldots,p$ , with  $(u_0,\hat{u}_0) = 0$ . We will use  $\mu(x_k, \omega_k) = \mu(x_k, \omega_k) + (\omega_{k-1}, \omega_{k-1}), \ldots$  and  $\mu_{k-1}$  and  $\omega_k$ –1 belong to G and satisfy<br>an induction argument on k. Assume that  $u_k$ –1 and  $\hat{u}_{k-1}$  belong to G and satisfy  $(4.18)$  and  $(4.19)$ , respectively (which obviously hold for  $k = 1$ ). First note that, because of the boundedness of  $T$ , we have

<span id="page-16-3"></span>
$$
||(u_{k-1},\widehat{u}_{k-1})||_{\widetilde{\mathcal{V}}} \leq C||(u_k,\widehat{u}_k)||_{\widetilde{\mathcal{V}}}.
$$

On the other hand, by using  $(2.16)$  and  $(2.17)$  we have that

$$
\mu \widehat{\boldsymbol{u}}_k + \widehat{\boldsymbol{u}}_{k-1} = \boldsymbol{u}_k \quad \text{in } \Omega
$$

and that  $\mu u_k + u_{k-1} \in \mathcal{G}$  satisfies

<span id="page-16-2"></span>
$$
\int_{\Omega} \rho c^2 \operatorname{div}(\mu \mathbf{u}_k + \mathbf{u}_{k-1}) \operatorname{div} \overline{\mathbf{v}} = -2 \int_{\Omega} \nu \operatorname{div} \mathbf{u}_k \operatorname{div} \overline{\mathbf{v}} - \int_{\Omega} \rho \widehat{\mathbf{u}}_k \cdot \overline{\mathbf{v}} \qquad \forall \mathbf{v} \in \mathcal{G}.
$$

Hence,  $u_k, \hat{u}_k \in \mathcal{G}$ .

We observe that the equation above also holds for any  $v \in \mathcal{K}$ . Then,

$$
\int_{\Omega} \rho c^2 \operatorname{div}(\mu \mathbf{u}_k + \mathbf{u}_{k-1}) \operatorname{div} \overline{\mathbf{v}} = -2 \int_{\Omega} \nu \operatorname{div} \mathbf{u}_k \operatorname{div} \overline{\mathbf{v}} - \int_{\Omega} \rho \widehat{\mathbf{u}}_k \cdot \overline{\mathbf{v}} \qquad \forall \mathbf{v} \in \mathcal{V}.
$$

Thus, considering test functions in  $\mathcal{D}(\Omega)^d \subset \mathcal{V}$  we obtain

(4.22) 
$$
\nabla((\mu \rho c^2 + 2\nu) \operatorname{div} \boldsymbol{u}_k) = \rho \widehat{\boldsymbol{u}}_k - \nabla(\rho c^2 \operatorname{div} \boldsymbol{u}_{k-1}).
$$

Let us assume that  $\mu \rho c^2 + 2\nu \neq 0$  in both  $\Omega_1$  and  $\Omega_2$  (we discuss the other case at the end of the proof). Hence, since  $\rho$ , c, and  $\nu$  are constant in each  $\Omega_i$ ,  $\rho_i \hat{u}_k - \nabla(-2, 1)$  $\nabla(\rho_i c_i^2 \text{ div } \mathbf{u}_{k-1}) \in \mathcal{L}^2(\Omega_i)^d$ , div  $\mathbf{u}_k|_{\Omega_i} \in \mathcal{H}^1(\Omega_i)$ , and

<span id="page-17-1"></span><span id="page-17-0"></span>
$$
\|\operatorname{div} \boldsymbol{u}_k\|_{1,\Omega_i} \leq C \left( \|\operatorname{div} \boldsymbol{u}_{k-1}\|_{1,\Omega_i} + \|(\boldsymbol{u}_k,\widehat{\boldsymbol{u}}_k)\|_{\widetilde{\mathcal{V}}} \right), \qquad i=1,2.
$$

Now, since  $u_k \in \mathcal{G}$ , due to Lemma [2.2](#page-5-0) we have that  $u_k \in H^s(\Omega)^d$ . Then, from [\(2.12\)](#page-5-1) and the previous estimate we have

$$
(4.23) \t\t ||\boldsymbol{u}_k||_{s,\Omega} \leq C \left( \|\operatorname{div} \boldsymbol{u}_{k-1}\|_{1,\Omega_1} + \|\operatorname{div} \boldsymbol{u}_{k-1}\|_{1,\Omega_2} + \|(\boldsymbol{u}_k,\widehat{\boldsymbol{u}}_k)\|_{\widetilde{\mathcal{V}}} \right).
$$

On the other hand, from [\(4.21\)](#page-16-2) we obtain

(4.24) 
$$
\|\widehat{\boldsymbol{u}}_k\|_{s,\Omega} \leq \frac{1}{\mu} \left(\|\widehat{\boldsymbol{u}}_{k-1}\|_{s,\Omega} + \|\boldsymbol{u}_k\|_{s,\Omega}\right)
$$

and, from [\(4.22\)](#page-17-0),

(4.25) 
$$
\|\text{div } \mathbf{u}_k\|_{1+s,\Omega_i} \leq C(\|\text{div } \mathbf{u}_{k-1}\|_{1+s,\Omega_i} + \|\widehat{\mathbf{u}}_k\|_{s,\Omega}), \qquad i=1,2.
$$

Finally, from [\(4.21\)](#page-16-2) again,

<span id="page-17-2"></span>
$$
(4.26) \quad \|\dim \widehat{\mathbf{u}}_k\|_{1+s,\Omega_i} \leq \frac{1}{\mu} (\|\dim \mathbf{u}_k\|_{1+s,\Omega_i} + \|\dim \widehat{\mathbf{u}}_{k-1}\|_{1+s,\Omega_i}), \qquad i=1,2.
$$

Hence, from inequalities  $(4.23)$ – $(4.26)$ , the inductive assumption, and  $(4.20)$ , we derive [\(4.18\)](#page-16-0) and [\(4.19\)](#page-16-1) provided  $\mu \rho c^2 + 2\nu \neq 0$  in both  $\Omega_1$  and  $\Omega_2$ .

In case that  $\mu \rho c^2 + 2\nu$  vanishes in  $\Omega_i$ ,  $i = 1$  or 2, arguing as in Remark [2.2](#page-6-5) we obtain that  $u_1|_{\Omega_i} = \hat{u}_1|_{\Omega_i} = 0$  and, once again, an induction argument allow us to conclude that  $u_k$ ,  $\hat{u}_k = 0$  in  $\Omega_i$ ,  $k = 1, ..., p$ . The proof is complete.

Now, we are in position to establish property P2.

<span id="page-17-3"></span>**Lemma 4.8.** Property P2 holds true. Moreover, for any  $(u, \hat{u}) \in \mathcal{E}$ , there exists  $\widetilde{\boldsymbol{u}}_h$ ,  $\widehat{\boldsymbol{u}}_h \in \mathcal{G}_h$  such that

$$
\|\boldsymbol{u}-\widetilde{\boldsymbol{u}}_h\|_{\mathrm{div},\Omega}+\|\widehat{\boldsymbol{u}}-\widetilde{\hat{\boldsymbol{u}}}_h\|_{\mathrm{div},\Omega}\leq Ch^s\|(\boldsymbol{u},\widehat{\boldsymbol{u}})\|_{\widetilde{\mathcal{V}}}.
$$

*Proof.* Let  $(u, \hat{u}) \in \mathcal{E}$ . According to Lemma [4.7](#page-16-4)  $u, \hat{u} \in H^s(\Omega)^d$  and div  $u, \text{div } \hat{u} \in H^1(s(\Omega))$ .  $H^{1+s}(\Omega_i)$ ,  $i=1,2$ . Let  $\Pi_h \boldsymbol{u} \in \mathcal{V}_h$  be the Raviart-Thomas interpolant of  $\boldsymbol{u}$ . Since  $\mathcal{V}_h = \mathcal{G}_h \oplus \mathcal{K}_h$ , we decompose  $\Pi_h \mathbf{u} = \widetilde{\mathbf{u}}_h + \widetilde{\mathbf{u}}_h$  with  $\widetilde{\mathbf{u}}_h \in \mathcal{G}_h$  and  $\widetilde{\mathbf{u}}_h \in \mathcal{K}_h$ . The same arguments from the proof of Lemma [4.5](#page-14-5) that lead to [\(4.14\)](#page-14-4) apply in this case and combined with Lemma [4.7](#page-16-4) allow us to prove that  $||u - \tilde{u}_h||_{div,\Omega} \leq C h^s ||(u, \hat{u})||_{\tilde{\mathcal{V}}}$ . A similar procedure can be used to define  $\hat{u}_h$  and to prove that  $\|\hat{u} - \hat{u}_h\|_{div,\Omega} \leq Ch^s \|(u, \hat{u})\|_{\tilde{\Omega}}$ .  $Ch^s \|(u,\widehat{u})\|_{\widetilde{\mathcal{V}}}$ .

We also have the following auxiliary result when the source terms are in  $\mathcal{E}$ .

<span id="page-17-4"></span>**Lemma 4.9.** For  $(f, g) \in \mathcal{E}$ , let  $(u, \hat{u}) := T(f, g)$  and  $(u_h, \hat{u}_h) := T_h(f, g)$ . Then,

$$
\|\boldsymbol{u}-\boldsymbol{u}_h\|_{\mathrm{div},\Omega}+\|\widehat{\boldsymbol{u}}-\widehat{\boldsymbol{u}}_h\|_{0,\Omega}\leq Ch^s\|( \boldsymbol{f},\boldsymbol{g})\|_{\widetilde{\mathcal{V}}}.
$$

*Proof.* Since  $\mathcal{G}_h \nsubseteq \mathcal{G}$ , we resort once more to the second Strang Lemma, which applied now to [\(2.17\)](#page-7-6) and [\(4.6\)](#page-13-4) leads to

$$
\|\boldsymbol{u}-\boldsymbol{u}_h\|_{\mathrm{div},\Omega} \leq C \left[\inf_{\boldsymbol{v}_h\in\mathcal{G}_h} \|\boldsymbol{u}-\boldsymbol{v}_h\|_{\mathrm{div},\Omega}+\sup_{\underline{0}\neq\boldsymbol{v}_h\in\mathcal{G}_h} \frac{a(\boldsymbol{u}-\boldsymbol{u}_h,\boldsymbol{v}_h)}{\|\boldsymbol{v}_h\|_{\mathrm{div},\Omega}}\right].
$$

From Lemma [4.8](#page-17-3) we know that there exists  $\widetilde{u}_h \in \mathcal{G}_h$  such that

$$
\|\boldsymbol{u}-\widetilde{\boldsymbol{u}}_h\|_{\text{div},\Omega}\leq Ch^s\|(\boldsymbol{u},\widehat{\boldsymbol{u}})\|_{\widetilde{\mathcal{V}}}\leq Ch^s\|(\boldsymbol{f},\boldsymbol{g})\|_{\widetilde{\mathcal{V}}}.
$$

Moreover, the consistency term above vanishes. In fact, consider  $v_h \in \mathcal{G}_h$  and the decomposition  $v_h = \frac{1}{\rho} \nabla \xi + \chi$  as in Lemma [4.1.](#page-11-4) Using the same arguments as in the proof of Lemma  $4.5$ , we prove that

$$
a(\boldsymbol{u}-\boldsymbol{u}_h,\boldsymbol{v}_h)=\int_{\Omega}\rho \boldsymbol{g}\cdot \overline{\chi}=0,
$$

where the last equality holds because  $g \in \mathcal{G}$  and  $\chi \in \mathcal{K}$ .

On the other hand, we know from [\(2.16\)](#page-7-1) and [\(4.5\)](#page-12-0) that  $\hat{u} = f$  and  $\hat{u}_h = \mathcal{P}_{\mathcal{G}_h} f$ , respectively. Then, since  $\mathcal{P}_{\mathcal{G}_h}$  is the H-orthogonal projection onto  $\mathcal{G}_h$ , we have that  $\|\hat{\mathbf{u}} - \hat{\mathbf{u}}_h\|_{\mathcal{H}} \leq \|\hat{\mathbf{u}} - \hat{\mathbf{u}}_h\|_{\mathcal{H}}$ , with  $\hat{\mathbf{u}}_h \in \mathcal{G}_h$  as in Lemma [4.8.](#page-17-3) Hence, we obtain

$$
\|\widehat{\boldsymbol{u}}-\widehat{\boldsymbol{u}}_h\|_{0,\Omega}\leq Ch^s\|(\boldsymbol{u},\widehat{\boldsymbol{u}})\|_{\widetilde{\mathcal{V}}}\leq Ch^s\|(\boldsymbol{f},\boldsymbol{g})\|_{\widetilde{\mathcal{V}}}.
$$

The proof is complete.  $\Box$ 

The above lemmas are the ingredients to prove spectral convergence and to obtain error estimates. Our first result is the following theorem which has been proved in [\[7\]](#page-25-8) as a consequence of property P1 (cf. Lemma [4.6\)](#page-16-5) and which shows that the proposed method is free of spurious modes.

**Theorem 4.1.** Let  $K \subset \mathbb{C}$  be a compact set such that  $K \cap \text{Sp}(T) = \emptyset$ . Then, there exists  $h_0 > 0$  such that, for all  $h \leq h_0$ ,  $K \cap \mathrm{Sp}(T_h) = \emptyset$ .

Let  $D \subset \mathbb{C}$  be a closed disk centered at  $\mu$ , such that  $D \cap \text{Sp}(T) = {\mu}.$  Let  $\mu_{1h}, \ldots, \mu_{m(h)h}$  be the eigenvalues of  $T_h$  contained in D (repeated according to their algebraic multiplicities). Under assumptions P1 and P2, it is proved in [\[7\]](#page-25-8) that  $m(h) = m$  for h small enough and that  $\lim_{h\to 0} \mu_{kh} = \mu$  for  $k = 1, \ldots, m$ .

On the other hand the arguments used in Section 5 of [\[2\]](#page-25-2) can be readily adapted to our problem, to obtain error estimates. We recall the definition of the gap between two closed subspaces  $W$  and  $Y$  of  $V$ :

$$
\widehat{\delta}(\mathcal{W}, \mathcal{Y}) := \max{\{\delta(\mathcal{W}, \mathcal{Y}), \delta(\mathcal{Y}, \mathcal{W})\}},
$$

with

$$
\delta(\mathcal{W},\mathcal{Y}):=\sup_{\substack{(\phi,\psi)\in\mathcal{W}\\ \Vert(\phi,\psi)\Vert_{\widetilde{\mathcal{V}}} = 1}}\left[\inf_{(\widehat{\phi},\widehat{\psi})\in\mathcal{Y}}\Vert(\phi,\psi)-(\widehat{\phi},\widehat{\psi})\Vert_{\widetilde{\mathcal{V}}}\right].
$$

Let  $\mathcal{E}_h$  be the invariant subspace of  $T_h$  relative to the eigenvalues  $\mu_{1h}, \ldots, \mu_{mh}$ converging to  $\mu$ . From Lemmas [4.6](#page-16-5)[–4.9,](#page-17-4) we derive the following results for which we do not include proofs to avoid repeating step by step those of [\[2,](#page-25-2) Section 5].

**Theorem 4.2.** There exist constants  $h_0 > 0$  and  $C > 0$  such that, for all  $h \leq h_0$ ,

$$
\widehat{\delta}(\mathcal{E}_h, \mathcal{E}) \leq C h^s.
$$

**Theorem 4.3.** There exist constants  $h_0 > 0$  and  $C > 0$  such that, for all  $h \leq h_0$ ,

$$
\left| \mu - \frac{1}{m} \sum_{k=1}^{m} \mu_{kh} \right| \le Ch^{2s},
$$
  

$$
\left| \frac{1}{\mu} - \frac{1}{m} \sum_{k=1}^{m} \frac{1}{\mu_{kh}} \right| \le Ch^{2s},
$$
  

$$
\max_{k=1,\dots,m} |\mu - \mu_{kh}| \le Ch^{2s/p},
$$

<span id="page-19-0"></span>where p is the ascent of the eigenvalue  $\mu$  of  $\mathbf{T}$ .

# 5. Numerical Results

We report in this section the results of some numerical tests, in order to assess the performance of the proposed method. With this end, first we introduce a convenient matrix form of the discrete problem which allows us to use standard eigensolvers. As a by-product, this matrix form also allows us to prove that Problems [3](#page-11-0) and [4](#page-11-1) are well posed.

Let  $\{\phi_j\}_{j=1}^N$  be a nodal basis of  $\mathcal{V}_h$ . We define the matrices  $\mathbf{K}_1 := (\mathbf{K}_{ij}^{(1)})$ ,  $\mathbf{K}_2 := (\mathbf{K}_{ij}^{(2)})$  and  $\mathbf{M} := (\mathbf{M}_{ij})$  as follows:  $\mathbf{K}_{ij}^{(1)} := 2\int_{\Omega}\nu\operatorname{div}\phi_i\operatorname{div}\phi_j,\ \ \ \mathbf{K}_{ij}^{(2)} := \int_{\Omega}\rho c^2\operatorname{div}\phi_i\operatorname{div}\phi_j,\ \ \text{ and }\ \ \mathbf{M}_{ij} := \int_{\Omega}\rho\phi_i\cdot\phi_j.$ 

The matrix form of Problem [3](#page-11-0) reads

(5.1) 
$$
(\lambda_h^2 \mathbf{M} + \lambda_h \mathbf{K}_1 + \mathbf{K}_2) \vec{\boldsymbol{u}}_h = 0
$$

where we denote by  $\vec{u}_h$  the vector of components of  $u_h$  in the nodal basis of  $\mathcal{V}_h$ . Analogously, the matrix form of Problem [4](#page-11-1) reads

,

<span id="page-19-1"></span>
$$
\begin{pmatrix} \mathbf{K}_2 & 0 \\ 0 & \mathbf{M} \end{pmatrix} \begin{pmatrix} \vec{u}_h \\ \vec{\tilde{u}}_h \end{pmatrix} = \lambda_h \begin{pmatrix} -\mathbf{K}_1 & -\mathbf{M} \\ \mathbf{M} & 0 \end{pmatrix} \begin{pmatrix} \vec{u}_h \\ \vec{\tilde{u}}_h \end{pmatrix},
$$

with  $\vec{\hat{u}}_h$  being the vector of components of  $\hat{u}_h$ . However, this problems is not suitable to be solved with standard eigensolvers, since neither the right-hand side nor the left-hand side matrix are Hermitian and positive definite.

Alternatively, for  $\lambda_h \neq 0$ , let  $\mu_h := \frac{1}{\lambda_h}$ . Then, problem [\(5.1\)](#page-19-1) is equivalent to

$$
(\mathbf{M} + 2\mu_h \mathbf{K}_1 + \mu_h^2 \mathbf{K}_2) \vec{\boldsymbol{u}}_h = \underline{0}.
$$

Introducing  $\vec{w}_h := \mu_h \vec{u}_h$ , the problem above is equivalent to

$$
\begin{pmatrix} \mathbf{M} & \mathbf{0} \\ \mathbf{0} & \mathbf{M} \end{pmatrix} \begin{pmatrix} \vec{u}_h \\ \vec{w}_h \end{pmatrix} = \mu_h \begin{pmatrix} -\mathbf{K}_1 & -\mathbf{K}_2 \\ \mathbf{M} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \vec{u}_h \\ \vec{w}_h \end{pmatrix},
$$

which in turn is equivalent to

$$
\begin{pmatrix}\n-\mathbf{K}_1 & -\mathbf{K}_2 \\
\mathbf{M} & 0\n\end{pmatrix}\n\begin{pmatrix}\n\vec{u}_h \\
\vec{w}_h\n\end{pmatrix} = \lambda_h \begin{pmatrix}\n\mathbf{M} & 0 \\
0 & \mathbf{M}\n\end{pmatrix}\n\begin{pmatrix}\n\vec{u}_h \\
\vec{w}_h\n\end{pmatrix}.
$$

Thus, the last problem is equivalent to Problem [3](#page-11-0) except for  $\lambda_h = 0$  and the matrix in its right-hand side is Hermitian and positive definite. Hence, it is well posed and can be safely solved by standard eigensolvers.

We implemented the proposed method in a MATLAB code. We applied it to a 2D rectangular rigid cavity filled with two fluids with different physical parameters as shown in Figure [2.](#page-20-0) The domain occupied by the fluids are  $\Omega_1 := (0, A) \times (0, H)$ and  $\Omega_2 := (0, A) \times (H, B)$ . For such a simple geometry, it is possible to calculate an analytical solution which will be used to validate our method.



<span id="page-20-0"></span>FIGURE 2. Two fluids in a rectangular rigid cavity.

Let  $u \in H_0(\text{div}, \Omega)$  be a solution of Problem [1.](#page-3-5) Testing it with  $v \in \mathcal{D}(\Omega)^2$  we have  $\nabla((2\lambda\nu + \rho c^2) \operatorname{div} \boldsymbol{u}) = -\lambda^2 \rho \boldsymbol{u} \in \mathcal{L}^2(\Omega)^2$ . Then,  $\widehat{p} := -(2\nu\lambda + \rho c^2) \operatorname{div} \boldsymbol{u} \in \mathcal{H}^1(\Omega)$ . Hence,  $\hat{p}_1|_{\Gamma} = \hat{p}_2|_{\Gamma}$ . Moreover,  $\boldsymbol{u} = -\frac{1}{\lambda^2}$  $\frac{1}{\lambda^2 \rho} \nabla \widehat{p}$ , which implies that  $\frac{1}{\rho_1}$  $\frac{\partial \widehat{p}_1}{\partial \nu} = \frac{1}{\rho_2}$  $\rho_2$  $\frac{\partial \widehat{p}_2}{\partial \widehat{p}_2}$ ∂ν on Γ. Then, we write problem  $(2.1)$ – $(2.8)$ , in terms of  $\hat{p}_i$  as follows:

$$
\Delta \widehat{p}_i = \frac{\lambda^2 \rho_i}{\rho_i c_i^2 + 2\nu_i \lambda} \widehat{p}_i \quad \text{in } \Omega_i, \ i = 1, 2,
$$
  
\n
$$
\frac{\partial \widehat{p}_i}{\partial \mathbf{n}_i} = 0 \quad \text{on } \Gamma_i, \ i = 1, 2,
$$
  
\n
$$
\widehat{p}_1 = \widehat{p}_2 \quad \text{on } \Gamma,
$$
  
\n
$$
\frac{1}{\rho_1} \frac{\partial \widehat{p}_1}{\partial \mathbf{n}} = \frac{1}{\rho_2} \frac{\partial \widehat{p}_2}{\partial \mathbf{n}} \quad \text{on } \Gamma.
$$

We proceed by separation of variables. Assuming that  $\hat{p}_i(x, y) = X_i(x)Y_i(y)$ , we are left with the following problem:

<span id="page-20-1"></span>(5.2) 
$$
\frac{X_i''(x)}{X_i(x)} + \frac{Y_i''(y)}{Y_i(y)} = \frac{\lambda^2 \rho_i}{\rho_i c_i^2 + 2\nu_i \lambda} \quad \text{in } \Omega_i,
$$

<span id="page-20-4"></span>(5.3) 
$$
X'_{i}(0) = X'_{i}(A) = 0, \qquad i = 1, 2,
$$

<span id="page-20-5"></span>(5.4) 
$$
Y'_1(0) = Y'_2(B) = 0,
$$

<span id="page-20-2"></span>(5.5) 
$$
\frac{1}{\rho_1} X_1(x) Y_1'(H) = \frac{1}{\rho_2} X_2(x) Y_2'(H), \quad 0 < x < A,
$$

<span id="page-20-3"></span>(5.6) 
$$
X_1(x)Y_1(H) = X_2(x)Y_2(H), \quad 0 < x < A.
$$

From [\(5.2\)](#page-20-1) we have that  $X_i(x)''/X_i(x)$  and  $Y_i(y)''/Y_i(y)$  are constant. Moreover, from [\(5.5\)](#page-20-2) and [\(5.6\)](#page-20-3), it is easy to check that  $Y_i(H)$  and  $Y'_i(H)$  cannot vanish simultaneously and  $X_1(x) = X_2(x)$  (actually, it is derived that  $X_1(x) = CX_2(x)$ , but the constant  $C$  can be chosen equal to one without loss of generality).

From the fact that  $X_i(x)''/X_i(x)$  is constant and [\(5.3\)](#page-20-4), we have that

$$
X_1(x) = X_2(x) = \cos\left(\frac{m\pi x}{A}\right), \qquad m = 0, 1, 2, ....
$$

On the other hand, from the fact that  $Y_i(y)''/Y_i(y)$  is also constant and [\(5.4\)](#page-20-5) we derive

<span id="page-21-0"></span>(5.7) 
$$
Y_1(y) = C_1 \cosh(r_m^{(1)}(\lambda)y)
$$
 and  $Y_2(y) = C_2 \cosh(r_m^{(2)}(\lambda)(y - B)),$ 

where  $C_1$  and  $C_2$  are constants and

$$
r_m^{(i)} := \sqrt{\frac{\lambda^2 \rho_i}{\rho_i c_i^2 + 2\nu_i \lambda} + \frac{m^2 \pi^2}{A^2}}, \qquad m = 0, 1, 2, \dots, \qquad i = 1, 2.
$$

Since  $Y_i(H)$  and  $Y'_i(H)$  cannot vanish simultaneously, [\(5.5\)](#page-20-2) and [\(5.6\)](#page-20-3) lead to

$$
\frac{1}{\rho_1} Y_1'(H) = \frac{1}{\rho_2} Y_2'(H) \quad \text{and} \quad Y_1(H) = Y_2(H),
$$

respectively. Thus, substituting [\(5.7\)](#page-21-0) into these equation yields the following linear system for the coefficients  $C_1$  and  $C_2$ :

$$
C_1 \cosh(r_m^{(1)}(\lambda)H) = C_2 \cosh(r_m^{(2)}(\lambda)(H - B)),
$$
  

$$
\frac{C_1 r_m^{(1)}(\lambda)}{\rho_1} \sinh(r_m^{(1)}(\lambda)H) = \frac{C_2 r_m^{(2)}(\lambda)}{\rho_2} \sinh(r_m^{(2)}(\lambda)(H - B)).
$$

For this system to have non trivial solutions, its determinant must vanish, which yields the following non linear equation in  $\lambda$  for  $m = 0, 1, 2, \ldots$ , whose roots are the eigenvalues of Problem [1:](#page-3-5)

$$
f_m(\lambda) := \frac{r_m^{(1)}(\lambda)}{\rho_1} \sinh(r_m^{(1)}(\lambda)H) \cosh(r_m^{(2)}(\lambda)(H - B)) - \frac{r_m^{(2)}(\lambda)}{\rho_2} \sinh(r_m^{(2)}(\lambda)(H - B)) \cosh(r_m^{(1)}(\lambda)H) = 0.
$$

We have computed some roots of the above equation and used these roots as exact eigenvalues to compare those obtained with the method proposed in this paper. For the geometrical parameters, we have taken  $A = 1$  m,  $B = 2$  m and  $H = 1.25$  m.

We have used physical parameters of water and air for the density and acoustic speed of the fluids in  $\Omega_1$  and  $\Omega_2$ , respectively:  $c_1 = 1430$  m/s,  $\rho_1 = 1000$  kg/m<sup>3</sup>,  $c_2 = 340$  m/s and  $\rho_2 = 1$  kg/m<sup>3</sup>. We have used uniform meshes as those shown in Figure [3.](#page-22-0) The refinement parameter N refers to the number of elements per width of the rectangle.



<span id="page-22-0"></span>FIGURE 3. Meshes for  $N = 4$  (left) and  $N = 8$  (right).

In presence of dissipation ( $\nu \neq 0$ ), the eigenvalues  $\lambda$  are complex numbers  $\lambda =$  $\eta + i\omega$ , with  $\eta < 0$  being the decay rate and  $\omega$  the vibration frequency. In absence of dissipation ( $\nu = 0$ ), the eigenvalues  $\lambda$  are purely imaginary ( $\eta = 0$ ). The same holds for the computed eigenvalues  $\lambda_h$ .

In our first test, we neglected the viscosity damping effects by taking  $\nu_1 = \nu_2 = 0$ . In this case, the eigenvalues  $\lambda$  are actually purely imaginary as can be seen from Figures [4](#page-22-1) and [5,](#page-23-0) which shows contour plots of  $\log(|f_m(\lambda)|)$  for the smallest values of  $m (0 \leq m \leq 3)$ . Accurate values of the zeros of  $f_m(\lambda)$  have been obtained with the MATLAB command fminsearch applied to  $|f_m(\lambda)|$ .



<span id="page-22-1"></span>FIGURE 4. Contour plots of  $\log(|f_m(\lambda)|)$  for  $m = 0$  and  $m = 1$ with vanishing viscosity  $(\nu = 0)$ .



<span id="page-23-0"></span>FIGURE 5. Contour plots of  $log(|f_m(\lambda)|)$  for  $m = 2$  and  $m = 3$ with vanishing viscosity  $(\nu = 0)$ .

Table [1](#page-23-1) shows the eigenvalues computed with the proposed method on successively refined meshes that approximate those shown in Figures [4](#page-22-1) and [5.](#page-23-0) Accurate values of the latter obtained with the MATLAB command fminsearch applied to  $|f_m(\lambda)|$  are also reported on the last line of the table as 'exact' eigenvalues.

m					$\Omega$				$\Omega$		$\mathcal{D}_{\mathcal{L}}$		$\mathcal{D}_{\mathcal{L}}$	
$N=8$		1066.07 i		1418.42 i		1784.37 <i>i</i>		1796.61 i		2118.35 i		2573.86 i		
$N = 16$		1067.78 i		1422.52 i		1781.49i		1797.09 i		2131.94 i		2569.90 i		
$N=32$		1068.21 i		1423.54 i		1780.73i		1797.21 i		2135.36 i		2568.40 i		
$N = 64$			1068.33 i		1423.79i		1780.55 i		1797.23 i		2136.22 i		2568.09 i	
Order			2.00	2.00		1.99		2.00		1.99		1.94		
Exact		1068.36 i		1423.87 i		1780.49 i		1797.24 i		2136.50 i		2568.54 i		
	m		$\Omega$					3		3		$\mathcal{D}_{\mathcal{L}}$		
	$N=8$		2807.28 i		3021.26 i		3142.54 i		3492.47i		3582.49 i			
	$N = 16$		2837.76 i		3037.38 i		3189.22i		3503.56 i		3568.16i			
	$N = 32$		2845.75 i		3041.02 i		3200.79 i		3506.19 i		3562.70 i			
	$N=64$		2848.88 i		3041.89 i		3204.78 i		3506.83 i		3561.22 i			
	Order		1.99		2.02		1.99		1.99		1.94			
	Exact		2849.56 i		3042.18i		3205.74 i		3507.16 i		3560.72 i			

<span id="page-23-1"></span>Table 1. Computed and exact eigenvalues for dissipative fluids in a rigid cavity.

As predicted by the theory, these eigenvalues are purely imaginary. The high accuracy of the computed eigenvalues can be observed from Table [1](#page-23-1) even for the coarsest mesh. We have used a least squares fitting to estimate the convergence rate for each eigenvalue, which are also reported in Table [1.](#page-23-1) A clear order  $\mathcal{O}(h^2)$ can be seen in all cases.

For the second test we have used the same physical parameters as above for both fluids, but considering now non vanishing viscosities. In order to make the dissipation effects more visible, we have used unrealistically large viscosity values:

 $\nu_1 = 9 \text{ N/m} s^2$  and  $\nu_2 = 1 \text{ N/m} s^2$ . We have repeated the scheme used for the first test. Figures [6](#page-24-0) and [7](#page-24-1) show the localization of all the exact eigenvalues  $\lambda$ . Notice that now all  $\lambda$  have negative real parts (the decay rate) as predicted by the theory.



<span id="page-24-0"></span>FIGURE 6. Contour plots of  $\log(|f_m(\lambda)|)$  for  $m = 0$  and  $m = 1$ with non-vanishing viscosity  $(\nu \neq 0)$ .



<span id="page-24-1"></span>FIGURE 7. Contour plots of  $log(|f_m(\lambda)|)$  for  $m = 2$  and  $m = 3$ with non-vanishing viscosity ( $\nu \neq 0$ ).

We report in Table [2](#page-25-13) the computed and 'exact' eigenvalues and the estimated convergence rates, which are in accordance with the theory once again.

m		1		$\Omega$		1	$\Omega$		
$N=8$		$-9.83 + 1066.03 i$		$-17.39 + 1418.31 i$		$-27.54 + 1784.16 i$	$-0.05 + 1796.61 i$		
$N=16$		$-9.86 + 1067.74i$		$-17.49 + 1422.41 i$		$-27.45 + 1781.27i$	$-0.05 + 1797.08 i$		
$N=32$		$-9.87 + 1068.17i$		$-17.51 + 1423.43 i$		$-27.43 + 1780.53 i$	$-0.05 + 1797.20 i$		
$N=64$		$-9.87 + 1068.38 i$		$-17.52 + 1423.78i$		$-27.42 + 1780.34 i$	$-0.05 + 1797.23 i$		
Order		2.00		2.00		1.99	2.00		
Exact	$-9.87 + 1068.32 i$			$-17.52 + 1423.76 i$		$-27.42 + 1780.27i$	$-0.05 + 1797.24 i$		
m		$\mathcal{D}_{\mathcal{L}}$		$\mathcal{D}$					
$N=8$		$-38.82 + 2118.00 i$		$-57.31 + 2573.22 i$		$-68.17 + 2806.45 i$	$-78.96 + 3020.24 i$		
$N=16$		$-39.32 + 2131.58 i$	$-57.13 + 2569.26 i$		$-69.65 + 2836.91 i$		$-79.80 + 3036.34 i$		
$N=32$		$-39.44 + 2135.00 i$		$-57.06 + 2567.76 i$	$-70.05 + 2844.09 i$		$-80.00 + 3039.96 i$		
$N=64$		$-39.58 + 2135.95 i$		$-57.04 + 2567.36 i$	$-70.15 + 2846.92 i$		$-80.04 + 3040.84 i$		
Order		1.99		1.94	1.99		2.02		
Exact		$-39.49 + 2136.14 i \mid -57.04 + 2567.22 i$			$-70.18 + 2847.60 i$		$-80.06 + 3041.13 i$		
	m	3		3			$\mathcal{D}$		
	$N=8$ $-85.43 + 3141.39 i$ $N=16$ $-87.99 + 3188.01 i$			$-105.51 + 3490.88 i$		$-111.02 + 3580.77 i$ $-110.14 + 3566.47 i$			
				$-106.18 + 3501.95 i$					
	$N=32$ $-88.62 + 3199.56 i$		$-106.34 + 3504.57 i$			$-109.80 + 3561.01 i$			
	$N=64$ $-88.88 + 3202.45 i$			$-106.38 + 3505.22 i$			$-109.70 + 3559.53 i$		
	Order 1.99				1.99		1.94		
	$-88.84 + 3203.41 i$ Exact			$-106.40 + 3505.44 i$			$-109.68 + 3559.03 i$		

<span id="page-25-13"></span>Table 2. Computed and exact eigenvalues for dissipative fluids in a rigid cavity.

#### **REFERENCES**

- <span id="page-25-12"></span>[1] A. Bermúdez, R. Durán, M.A. Muschietti, R. Rodríguez, and J. Solomin, Finite element vibration analysis of fluid-solid systems without spurious modes, SIAM J. Numer. Anal., 32 (1995) 1280–1295.
- <span id="page-25-2"></span>[2] A. Bermúdez, R. Durán, R. Rodríguez, and J. Solomin, Finite element analysis of a quadratic eigenvalue problem arising in dissipative acoustics, SIAM J. Numer. Anal., 38 (2000) 267– 291.
- <span id="page-25-0"></span>[3] A. Bermúdez, P. Gamallo, L. Hervella-Nieto, R. Rodríguez, and D. Santamarina, Fluidstructure acoustic interaction, in Computational Acoustics of Noise Propagation in Fluids. Finite and Boundary Element Methods. S. Marburg, B. Nolte, eds., Springer, 2008, pp. 253– 286.
- <span id="page-25-3"></span>[4] A. Bermúdez and R. Rodríguez, Numerical computation of elastoacoustic vibrations with interface damping, in Équations aux Dérivées Partielles et Applications, Gauthier-Villars, Paris, 1998, pp. 165–187.
- <span id="page-25-4"></span>[5] A. Bermúdez and R. Rodríguez, Modeling and numerical solution of elastoacoustic vibrations with interface damping, Internat. J. Numer. Methods Engng., 46 (1999) 1763–1779.
- <span id="page-25-6"></span>[6] A. Bermúdez, R. Rodríguez, and D. Santamarina, Two discretization schemes for a timedomain dissipative acoustics problem, Math. Models Methods Appl. Sci., 16 (2006) 1559– 1598.
- <span id="page-25-8"></span>[7] J. Descloux, N. Nassif, and J. Rappaz, On spectral approximation. Part 1: The problem of convergence. RAIRO Anal. Numér., 12 (1978) 97-112.
- <span id="page-25-9"></span>[8] J. Descloux, N. Nassif, and J. Rappaz, On spectral approximation. Part 2: Error estimates for the Galerkin method. RAIRO Anal. Numér.,  $12$  (1978) 113-119.
- <span id="page-25-10"></span><span id="page-25-1"></span>[9] T. Kato, Perturbation Theory for Linear Operators, Springer-Verlag, Berlin, 1966.
- [10] L.E. Kinsler, A.R, Frey, A.B. Coppens, and J.V. Sanders, Fundamentals of Acoustics, John Wiley & Sons, 2000.
- <span id="page-25-7"></span>[11] M.G. Krein and H. Langer, On some mathematical principles in the linear theory of damped oscillations of continua I. Integral Equations Operator Theory, 1 (1978) 364–399.
- <span id="page-25-11"></span><span id="page-25-5"></span>[12] L.D. Landau and E.M. Lifshitz, Fluid Mechanics, Pergamon Press, 1982.
- [13] P. Monk, Finite Element Methods for Maxwell's Equations, Oxford Clarendon Press, 2003.
- <span id="page-26-1"></span><span id="page-26-0"></span>[14] P.M. Morse, Vibration and Sound, Acoustical Society of America through the American Institute of Physics, 1995.
- [15] R. Ohayon and C. Soize, Structural Acoustic and Vibration. Mechanical Models Variational Formulations and Discretization, Academic Press, New York, 1998.
- <span id="page-26-3"></span>[16] M. Petzoldt, Regularity and error estimates for elliptic problems and discontinuous coefficients, Ph.D. Thesis, Berlin, Freie Univ. (2001).
- [17] M. Petzoldt, Regularity result for the Laplace interface problems in two dimensions, Z. Anal. Anw., 20 (2001) 431–455.
- <span id="page-26-4"></span>[18] M. Petzoldt, A posteriori error estimates for elliptic equations with discontinuous coefficients, Adv. Comp. Math., 16 (2002) 47–75.
- <span id="page-26-2"></span>[19] A.D. Pierce, Acoustics: An Introduction to its Physical Principles and Applications, Acoustical Society of America through the American Institute of Physics, 1994.
- <span id="page-26-5"></span>[20] M. Reed and B. Simon, Methods of Modern Mathematical Physics. IV. Analysis of Operators Academic Press, New York, 1978.

CI<sup>2</sup>MA, DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE.

E-mail address: flepe@ing-mat.udec.cl

Departamento de Matematicas, Facultad de Ciencias, Universidad de Oviedo, Calvo ´ SOTELO S/N, OVIEDO, SPAIN.

E-mail address: salim@uniovi.es

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD DEL BÍO-BÍO, CASILLA 5-C, CONCEPCIÓN, CHILE AND CENTRO DE INVESTIGACIÓN EN INGENIERÍA MATEMÁTICA  $(Cl^2MA)$ , UNIVERSIDAD DE CONCEPCIÓN, CONCEPCIÓN, CHILE.

E-mail address: dmora@ubiobio.cl

CI<sup>2</sup>MA, DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C, CONCEPCIÓN, CHILE.

E-mail address: rodolfo@ing-mat.udec.cl