Yet another generalization of Postnikov's hook length formula for binary trees

Guo-Niu HAN

ABSTRACT. — We discover another one-parameter generalization of Postnikov's hook length formula for binary trees. The particularity of our formula is that the hook length h_v appears as an exponent. As an application, we derive another simple hook length formula for binary trees when the underlying parameter takes the value 1/2.

1. Introduction

Consider the set $\mathcal{B}(n)$ of all binary trees with n vertices. For each vertex v of $T \in \mathcal{B}(n)$ the hook length of v, denoted by h_v , or just h for short, is the number of descendants of v (including v). The following hook length formula for binary trees

(1)
$$\sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \left(1 + \frac{1}{h_v} \right) = \frac{2^n}{n!} (n+1)^{n-1}$$

was discovered by Postnikov [Po04]. Further combinatorial proofs and extensions have been proposed by several authors [CY08, GS06, MY07, Se08]. In particular, Lascoux conjectured the following one-parameter generalization

(2)
$$\sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \left(x + \frac{1}{h_v} \right) = \frac{1}{(n+1)!} \prod_{k=0}^{n-1} ((n+1+k)x + n + 1 - k),$$

which was subsequently proved by Du-Liu [DL08]. The latter generalization appears to be very natural, because the *left-hand side* of (2) can be obtained from the left-hand side of (1) by replacing 1 by x.

It is also natural to look for an extension of (1) by introducing a new variable z in the *right-hand side*, namely by replacing $2^n(n+1)^{n-1}/n!$ by $2^n z(n+z)^{n-1}/n!$ It so happens that the corresponding left-hand side is also a sum on binary trees, but this time the hook length h_v appears as an exponent. The purpose of this Note is to prove the following Theorem.

Theorem 1. For each positive integer n we have

(3)
$$\sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \frac{(z+h)^{h-1}}{h(2z+h-1)^{h-2}} = \frac{2^n z}{n!} (n+z)^{n-1}.$$

With z = 1 in (3) we recover Postnikov's identity (1). The following corollary is derived from our identity (3) by taking z = 1/2.

Corollary 2. For each positive integer n we have

(4)
$$\sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \left(1 + \frac{1}{2h}\right)^{h-1} = \frac{(2n+1)^{n-1}}{n!}.$$

2. Proof of the Theorem

Let us take an example before proving the Theorem. There are five binary trees with n = 3 vertices:



The hook lengths of T_1, T_2, T_3, T_4 are all the same 1, 2, 3; but the hook lengths of T_5 are 1, 1, 3. The left-hand side of (3) is then equal to

$$4 \times \frac{1}{(2z)^{-1}} \cdot \frac{(z+2)^1}{2} \cdot \frac{(z+3)^2}{3(2z+1)} + \frac{1}{(2z)^{-1}} \cdot \frac{1}{(2z)^{-1}} \cdot \frac{(z+3)^2}{3(2z+1)} = \frac{2^3 z(z+3)^2}{3!}$$

Let y(x) be a formal power series in x such that

(5)
$$y(x) = e^{xy(x)}.$$

By the Lagrange inversion formula $y(x)^z$ has the following explicit expansion:

(6)
$$y(x)^{z} = \sum_{n \ge 0} z(n+z)^{n-1} \frac{x^{n}}{n!}.$$

Since $y^{2z} = (y^z)^2$ we have

(7)
$$\sum_{n\geq 0} 2z(n+2z)^{n-1} \frac{x^n}{n!} = \left(\sum_{n\geq 0} z(n+z)^{n-1} \frac{x^n}{n!}\right)^2.$$

Comparing the coefficients of x^n on both sides of (7) yields the following Lemma.

Lemma 3. We have

(8)
$$\frac{2z(n+2z)^{n-1}}{n!} = \sum_{k=0}^{n} \frac{z(k+z)^{k-1}}{k!} \times \frac{z(n-k+z)^{n-k-1}}{(n-k)!}.$$

Proof of the Theorem. Let

$$P(n) = \sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \frac{(z+h)^{h-1}}{h(2z+h-1)^{h-2}}$$

With each binary tree $T \in \mathcal{B}(n)$ $(n \geq 1)$ we can associate a triplet (T', T'', u), where $T' \in \mathcal{B}(k)$ $(0 \leq k \leq n-1)$, $T'' \in \mathcal{B}(n-1-k)$ and u is a vertex of hook length $h_u = n$. Hence

(9)
$$P(n) = \sum_{k=0}^{n-1} P(k)P(n-1-k) \times \frac{(z+n)^{n-1}}{n(2z+n-1)^{n-2}}.$$

It is routine to verify that $P(n) = 2^n z(z+n)^{n-1}/n!$ for n = 1, 2, 3. Suppose that $P(k) = 2^k z(z+k)^{k-1}/k!$ for $k \le n-1$. From identity (9) and Lemma 3 we have

$$P(n) = \sum_{k=0}^{n-1} \frac{2^k z(z+k)^{k-1}}{k!} \times \frac{2^{n-k-1} z(z+n-k-1)^{n-k-2}}{(n-k-1)!} \times \frac{(z+n)^{n-1}}{n(2z+n-1)^{n-2}} = \frac{2^n z}{n!} (z+n)^{n-1}.$$

By induction, formula (3) is true for any positive integer n.

3. Concluding and Remarks

The right-hand sides of (3) and (4) have been studied by other authors [GS06, DL08, MY07], but our formula has the following two major differences: (i) the hook length h_v appears as an exponent; (ii) the underlying set remains the set of binary trees, whereas in the above mentioned papers the summation has been changed to the set of *m*-ary trees or plane forests. It is interesting to compare Corollary 2 with the following results obtained by Du and Liu [DL08]. Note that the right-hand sides of formulas (4), (10) and (11) are all identical!

Proposition 4. For each positive integer n we have

(10)
$$\sum_{T \in \mathcal{T}(n)} \prod_{v \in I(T)} \left(\frac{2}{3} + \frac{1}{3h}\right) = \frac{(2n+1)^{n-1}}{n!},$$

where $\mathcal{T}(n)$ is the set of all 3-ary trees with n internal vertices and I(T) is the set of all internal vertices of T.

Proposition 5. For each positive integer n we have

(11)
$$\sum_{T \in \mathcal{F}(n)} \prod_{v \in (T)} \left(2 - \frac{1}{h}\right) = \frac{(2n+1)^{n-1}}{n!},$$

where $\mathcal{F}(n)$ is the set of all plane forests with n vertices.

References

- [CY08] Chen, William Y.C.; Yang, Laura L.M., On Postnikov's hook length formula for binary trees, European Journal of Combinatorics, in press, 2008.
- [DL08] Du, Rosena R. X.; Liu, Fu, (k, m)-Catalan Numbers and Hook Length Polynomials for Plane Trees, European J. Combin, 28 (2007), pp. 1312–1321.
- [GS06] Gessel, Ira M.; Seo, Seunghyun, A refinement of Cayley's formula for trees, arXiv:math.CO/0507497.
- [MY07] Moon, J. W.; Yang, Laura L. M., Postnikov identities and Seo's formulas, Bull. Inst. Combin. Appl., 49 (2007), pp. 21–31.
- [Po04] Postnikov, Alexander, Permutohedra, associahedra, and beyond, arXiv:math. CO/0507163, 2004.

I.R.M.A. UMR 7501 Université Louis Pasteur et CNRS, 7, rue René-Descartes F-67084 Strasbourg, France guoniu@math.u-strasbg.fr