

The resolvent order: a unification of the orders by Zarantonello, by Loewner, and by Moreau

Sedi Bartz*, Heinz H. Bauschke[†] and Xianfu Wang[‡]

June 28, 2016

Abstract

We introduce and investigate the resolvent order, which is a binary relation on the set of firmly nonexpansive mappings. It unifies well-known orders introduced by Loewner (for positive semidefinite matrices) and by Zarantonello (for projectors onto convex cones). A connection with Moreau's order of convex functions is also presented. We also construct partial orders on (quotient sets of) proximal mappings and convex functions. Various examples illustrate our results.

2010 Mathematics Subject Classification: Primary 06A06, 26B25, 47H05, 47H09, Secondary 15B57, 47L07, 52A41, 90C25.

Keywords: Baillon–Haddad theorem, convex cone, convex function, envelope, firmly nonexpansive mapping, Loewner order, Löwner order, maximally monotone operator, Moreau envelope, Moreau's order, positive semidefinite matrix, projection mapping, projector, proximal mapping, proximity operator, resolvent, resolvent order, Zarantonello's order.

1 Introduction

In this paper, we assume that

X is a real Hilbert space, (1)

with inner product $\langle \cdot, \cdot \rangle$ and induced norm $\|\cdot\|$. We denote the set of all functions from X to $\mathbb{R} \cup \{+\infty\}$ that are *convex*¹, lower semicontinuous and proper by $\Gamma_0(X)$. Let $A: X \rightrightarrows X$ be a

*Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: sedi.bartz@ubc.ca.

[†]Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: heinz.bauschke@ubc.ca.

[‡]Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: shawn.wang@ubc.ca.

¹We assume the reader is familiar with basic convex analysis; see, e.g., [12, 13, 17, 21, 18, 4].

set-valued operator, i.e., $(\forall x \in X) Ax \subseteq X$ and denote the graph of A by $\text{gra } A$. Recall that A is monotone if

$$(\forall (x, x^*) \in \text{gra } A)(\forall (y, y^*) \in \text{gra } A) \quad \langle x - y, x^* - y^* \rangle \geq 0 \quad (2)$$

and that A is *maximally monotone* if it is monotone and cannot be extended without destroying monotonicity. The notion of maximal monotonicity has proven to be useful in modern optimization and nonlinear analysis; see, e.g., [4, 5, 6, 8, 18, 19, 20, 24, 25]. We denote the set of maximally monotone operators on X by $\mathcal{M}(X)$. This set includes subdifferential operators of functions in $\Gamma_0(X)$ as well as all square matrices with symmetric parts that are positive semidefinite. Furthermore, we denote by $\mathcal{F}(X)$ the set of all mappings $T: X \rightarrow X$ that are *firmly nonexpansive*², i.e.,

$$(\forall x \in X)(\forall y \in X) \quad \|Tx - Ty\|^2 \leq \langle x - y, Tx - Ty \rangle. \quad (3)$$

Thanks to the work of Minty [15] (see also [9]), we can identify a maximally monotone operator A from $\mathcal{M}(X)$ with its *resolvent* in $\mathcal{F}(X)$ via

$$J_A := (\text{Id} + A)^{-1}. \quad (4)$$

Here $\text{Id} = \nabla q = \partial q$ is the identity operator on X , where $q: x \mapsto \frac{1}{2}\|x\|^2$. If we focus instead on the important subset of subdifferential operators in $\mathcal{M}(X)$, then we recover Moreau's [16] *proximal mapping* (or proximity operator)

$$P_f := J_{\partial f} = (\text{Id} + \partial f)^{-1}, \quad (5)$$

where $f \in \Gamma_0(X)$ and $\partial f \in \mathcal{M}(X)$ is the subdifferential operator of f . The set of proximal mappings, which we write as $\mathcal{P}(X)$, can also be described as follows. Given $f \in \Gamma_0(X)$, let $\text{env}(f) := q \square f$ be the (Moreau) *envelope* of f , where \square denotes infimal convolution. The set of all envelopes is written as $M_0(X)$. Then

$$\nabla \text{env}(f^*) = P_f = J_{\partial f} = (\text{Id} + \partial f)^{-1}, \quad (6)$$

where $f^* \in \Gamma_0(X)$ is the Fenchel conjugate of f . (Thus, we can loosely write $M_0(X) = \Gamma_0(X) \square q$ and $\nabla M_0(X) = \mathcal{P}(X)$.)

Having set up the necessary notation, we can now describe the goal and the organization of this paper.

The goal of this paper is to introduce a new order³ on $\mathcal{F}(X)$ which we call the resolvent order. It induces orders on $\mathcal{P}(X)$, $\mathcal{M}(X)$, $\Gamma_0(X)$, and $M_0(X)$ which will allow us to unify and connect to several well known orders from linear and nonlinear analysis, namely to the orders by Zarantonello, by Loewner, and by Moreau. We provide several examples and also present a partial order on (a quotient set of) the set of proximal mappings $\mathcal{P}(X)$ and on (a quotient set of) the set of convex functions $\Gamma_0(X)$.

The remainder of the paper is organized as follows. In Section 2, we present various results that make the proofs of the main results more structured. The resolvent order on $\mathcal{F}(X)$ is defined in

² Note that by the Cauchy–Schwarz inequality, every firmly nonexpansive mapping is *nonexpansive*, i.e., Lipschitz continuous with constant 1.

³To keep the language in this paper from being overly technical, we will refer to an “order” as a binary relation that is at least reflexive.

Section 3, where we also provide basic properties and characterizations for $\mathcal{P}(X)$. In fact, transitivity of the order is established for $\mathcal{P}(X)$. In Section 4, we discuss partitions of the identity and show that transitivity fails for $\mathcal{F}(X)$. In Sections 5, 6, and 7, we connect the resolvent order to the orders by Zarantonello, by Loewner, and by Moreau, respectively. New orders on $\mathcal{M}(X)$ and $\Gamma_0(X)$ are introduced in Section 8. These orders are not *partial orders*. A quotient construction is presented in Section 9 which results in partial orders on $\mathcal{P}(X)$ and on $\Gamma_0(X)$.

The notation we employ is standard and follows, e.g., [4].

2 Auxiliary results

In this section, we collect various results that will be useful later.

Fact 2.1. *Let $T: X \rightarrow X$. Then the following are equivalent:*

- (i) T is firmly nonexpansive, i.e., $(\forall x \in X)(\forall y \in X) \langle Tx - Ty, x - y \rangle \geq \|Tx - Ty\|^2$.
- (ii) $(\forall x \in X)(\forall y \in X) \|Tx - Ty\|^2 + \|(x - Tx) - (y - Ty)\|^2 \leq \|x - y\|^2$.
- (iii) $\text{Id} - T$ is firmly nonexpansive.
- (iv) $2T - \text{Id}$ is nonexpansive.

Proof. See, e.g., [4], [10], or [11]. ■

Corollary 2.2. *The sets $\mathcal{F}(X)$ and $\mathcal{P}(X)$ are convex. If $\lambda \in [0, 1]$, then $\lambda \mathcal{F}(X) \subseteq \mathcal{F}(X)$ and $\lambda \mathcal{P}(X) \subseteq \mathcal{P}(X)$.*

Proof. For $\mathcal{F}(X)$, the convexity follows using the last item from Fact 2.1 (see also [2, Corollary 1.8]). For the convexity of $\mathcal{P}(X)$, see [16]. To obtain the inclusions, it suffices to note that $0 \in \mathcal{P}(X)$. ■

Lemma 2.3. *Let T_1 and T_2 be firmly nonexpansive on X . Then $T_2 - T_1$ is nonexpansive.*

Proof. By Fact 2.1, we can write each $T_i = (\text{Id} + N_i)/2$, where N_i is nonexpansive. It follows that $T_2 - T_1 = (N_2 - N_1)/2$ is nonexpansive. ■

Fact 2.4. *Let $f \in \Gamma_0(X)$ and set $h := f^* - \mathfrak{q}$. Then the following are equivalent:*

- (i) f is Fréchet differentiable on X and ∇f is nonexpansive.
- (ii) f is Fréchet differentiable on X and ∇f is firmly nonexpansive.
- (iii) $\mathfrak{q} - f$ is convex
- (iv) $f^* - \mathfrak{q}$ is convex.
- (v) $h \in \Gamma_0(X)$ and $f = \text{env}(h^*)$.
- (vi) $h \in \Gamma_0(X)$ and $\nabla f = P_h$.

Proof. See [4, Theorem 18.17], and also [1], [3], [16]. ■

Corollary 2.5. *Any linear combination of proximal mappings that is monotone and nonexpansive is actually a proximal mapping.*

Proof. Let h_1, \dots, h_n be in $\Gamma_0(X)$ such that $P := \sum_{i=1}^n \alpha_i P_{h_i}$ is nonexpansive and monotone. Set $(\forall i \in \{1, \dots, n\}) f_i := \text{env}(h_i^*) = h_i^* \square q = (h_i + q)^*$, $f := \sum_{i=1}^n \alpha_i f_i$ and $h := f^* - q$. Then $P = \nabla f$ and f is thus convex and Fréchet differentiable. By Fact 2.4, $h \in \Gamma_0(X)$ and $P = \nabla f = P_h$. ■

Corollary 2.6. *Let P_1 and P_2 be proximal mappings. Then $P_1 - P_2$ is a proximal mapping if and only if $P_1 - P_2$ is monotone.*

Proof. “ \Rightarrow ”: Clear. “ \Leftarrow ”: By Lemma 2.3, $P_1 - P_2$ is nonexpansive. Now apply Corollary 2.5. ■

Fact 2.7. *Let $T: X \rightarrow X$ be linear and self-adjoint. Then T is firmly nonexpansive if and only if T is monotone and nonexpansive, in which case T is a proximal mapping*

Proof. See [4, Corollary 18.15] and also [1]. ■

3 The resolvent order

From the point of view of monotone operator theory, the set of firmly nonexpansive mappings is the same as the set of resolvents. This motivates the language in the following definition.

Definition 3.1. (resolvent order) *We define on $\mathcal{F}(X)$ a binary relation via*

$$T_1 \preceq T_2 \Leftrightarrow T_2 - T_1 \in \mathcal{F}(X). \quad (7)$$

Let us collect some basic properties.

Lemma 3.2. *Let T, T_0, T_1 be in $\mathcal{F}(X)$. The binary relation \preceq satisfies the following:*

- (i) **(reflexivity)** $T \preceq T$.
- (ii) **(existence of least and greatest element)** $0 \preceq T \preceq \text{Id}$.
- (iii) **(order reversal)** $T_0 \preceq T_1 \Leftrightarrow \text{Id} - T_1 \preceq \text{Id} - T_0$.
- (iv) $T_0 \preceq T_1 \Leftrightarrow (\forall \lambda \in [0, 1]) T_0 \preceq (1 - \lambda)T_0 + \lambda T_1 \preceq T_1$.

Proof. (i): $T - T = 0$ is firmly nonexpansive. (ii): $T - 0 = T$ is firmly nonexpansive as is $\text{Id} - T$ by Fact 2.1. (iii): Indeed, $T_1 - T_0 = (\text{Id} - T_0) - (\text{Id} - T_1)$. (iv): Suppose that $T_0 \preceq T_1$, i.e., $T_1 - T_0 \in \mathcal{F}(X)$. Let $\lambda \in [0, 1]$. Then $T_1 - ((1 - \lambda)T_0 + \lambda T_1) = (1 - \lambda)(T_1 - T_0)$ and $(1 - \lambda)T_0 + \lambda T_1 - T_0 = \lambda(T_1 - T_0)$ both of which are firmly nonexpansive by Corollary 2.2. The converse implication is trivial. ■

The following two observations are easily verified.

Example 3.3. (lack of symmetry) Suppose that $X \neq \{0\}$. Then $0 \preceq \text{Id}$ but $\text{Id} \not\preceq 0$.

Example 3.4. (lack of antisymmetry) Suppose that x_1 and x_2 are two distinct vectors in X , and set $(\forall x \in X) T_1(x) := x_1$ and $T_2(x) := x_2$. Then $T_1 \preceq T_2$ and $T_2 \preceq T_1$ yet $T_1 \neq T_2$.

In Section 9, we will present a quotient construction that makes the binary relation antisymmetric.

We now turn to proximal mappings which allows us to obtain stronger conclusions.

Theorem 3.5. Let f and g be in $\Gamma_0(X)$. Then the following are equivalent:

- (i) $P_f \preceq P_g$, i.e., $P_g - P_f \in \mathcal{F}(X)$.
- (ii) $P_g - P_f \in \mathcal{P}(X)$.
- (iii) $\text{env}(g^*) - \text{env}(f^*) \in \mathbf{M}_0(X)$.
- (iv) $\text{env}(f) - \text{env}(g) \in \mathbf{M}_0(X)$.

Proof. “(i) \Leftarrow (ii)”: Clear. “(i) \Rightarrow (ii)”: Since $P_g - P_f$ is firmly nonexpansive it is also monotone. Now apply Corollary 2.6. “(ii) \Rightarrow (iii)”: Integrate. “(ii) \Leftarrow (iii)”: Differentiate. “(iii) \Leftarrow (iv)”: This is clear since $\text{env}(g^*) - \text{env}(f^*) = (q - \text{env}(g)) - (q - \text{env}(f)) = \text{env}(f) - \text{env}(g)$. ■

Theorem 3.6. (transitivity for proximal mappings) Let f, g, h be in $\Gamma_0(X)$ such that $P_f \preceq P_g$ and $P_g \preceq P_h$. Then $P_f \preceq P_h$.

Proof. By the hypothesis and Theorem 3.5, there exist a and b in $\Gamma_0(X)$ such that

$$P_g - P_f = P_a \quad \text{and} \quad P_h - P_g = P_b. \quad (8)$$

Adding yields $P_h - P_f = P_a + P_b$. On the other hand, $P_a + P_b$ is monotone because P_a and P_b are monotone. Altogether, we deduce from Corollary 2.6, that $P_h - P_f$ is a proximal mapping. ■

Corollary 3.7. (proximal mappings are directed) $(\mathcal{P}(X), \preceq)$ is a directed set.

Proof. The reflexivity of \preceq was observed in Lemma 3.2(i) while the transitivity of \preceq is a consequence of Theorem 3.6. Finally, if P_1 and P_2 are in $\mathcal{P}(X)$, then $P_1 \preceq \text{Id}$ and $P_2 \preceq \text{Id}$ by Lemma 3.2(ii). ■

We conclude this section with an example.

Example 3.8. Denote the unit ball centered at 0 of radius 1 in X by C , and set $T := \text{Id} - P_C$. Then $(\forall n \in \mathbb{N}) T^n = \text{Id} - P_{nC} \in \mathcal{P}(X)$ and $T^n - T^{n+1} = P_{(n+1)C} - P_{nC} \in \mathcal{P}(X)$. Consequently,

$$(\forall n \in \mathbb{N}) \quad 0 \preceq T^{n+1} \preceq T^n \preceq \dots \preceq T \preceq T^0 = \text{Id}. \quad (9)$$

Proof. The identity for T^n is easily verified by mathematical induction and discussing cases. To verify that $T^n - T^{n+1}$, observe that by Corollary 2.6 it suffices to show that $T^n - T^{n+1}$ is monotone. In turn, this is achieved by discussing cases and invoking the Cauchy–Schwarz inequality. ■

4 Partitions of the identity and the partial sum property

In this section, we discuss partial sums of firmly nonexpansive mappings arising in partitions of the identity. Somewhat surprisingly, we also show that the transitivity result for proximal mappings (Theorem 3.6) *fails* for firmly nonexpansive mappings (see Example 4.5 below).

We start with a positive result.

Lemma 4.1. *Let T_1, T_2, T_3 be in $\mathcal{F}(X)$ such that $T_1 + T_2 + T_3 = \text{Id}$. Then $T_1 + T_2$ is firmly nonexpansive.*

Proof. Since $T_1 + T_2 = \text{Id} - T_3$, this follows from Fact 2.1. ■

For proximal mappings we are able to extend Lemma 4.1 from 3 to any number of operators:

Theorem 4.2. (partial sum property for proximal mappings) *Let $n \in \{1, 2, \dots\}$, let P_1, \dots, P_n be in $\mathcal{P}(X)$ such that $P_1 + P_2 + \dots + P_n = \text{Id}$, and let $m \in \{1, \dots, n\}$. Then $P_1 + \dots + P_m \in \mathcal{P}(X)$.*

Proof. There exist functions f_1, \dots, f_n in $\Gamma_0(X)$ such that for each i , $\nabla \text{env}(f_i^*) = P_i$, and $\text{env}(f_1^*) + \dots + \text{env}(f_n^*) = q$. It follows that

$$q - (\text{env}(f_1^*) + \dots + \text{env}(f_m^*)) = \text{env}(f_{m+1}^*) + \dots + \text{env}(f_n^*) \quad (10)$$

is convex. By Fact 2.4, $\nabla(\text{env}(f_1^*) + \dots + \text{env}(f_m^*)) = P_1 + \dots + P_m$ is a proximal mapping. ■

Surprisingly, the counterpart of Theorem 4.2 for firmly nonexpansive mappings is false as the next two results show.

Lemma 4.3. *In $X = \mathbb{R}^2$, let $n \in \{2, 3, \dots\}$, let $\theta \in]\arccos(1/\sqrt{2}), \arccos(1/\sqrt{2n})]$, set $\alpha := 1/(2n \cos(\theta))$, and denote by R_θ be the counterclockwise rotator by θ . Then the following hold:*

- (i) αR_θ and $\alpha R_{-\theta}$ are firmly nonexpansive.
- (ii) $n\alpha R_\theta$ and $n\alpha R_{-\theta}$ are not firmly nonexpansive.
- (iii) $n\alpha R_\theta + n\alpha R_{-\theta} = \text{Id}$.

Proof. Observe that $(\forall x \in X) \langle x, R_\theta x \rangle = \cos(\theta) \|x\|^2 = \cos(\theta) \|R_\theta x\|^2$. It follows that

$$(\forall \alpha \in \mathbb{R}_+) \quad \alpha R_\theta \in \mathcal{F}(X) \quad \Leftrightarrow \quad \alpha \in [0, \cos(\theta)]. \quad (11)$$

Let $\alpha \in \mathbb{R}_{++}$ satisfy

$$n\alpha R_\theta + n\alpha R_{-\theta} = n\alpha (R_\theta + R_{-\theta}) = n\alpha \begin{pmatrix} 2\cos(\theta) & 0 \\ 0 & 2\cos(\theta) \end{pmatrix} = \text{Id}; \quad (12)$$

equivalently,

$$\alpha := \frac{1}{2n \cos(\theta)}. \quad (13)$$

Combining (11) and (13) yields

$$\alpha R_\theta \in \mathcal{F}(X) \Leftrightarrow \frac{1}{2n} \leq \cos^2(\theta). \quad (14)$$

On the other hand, $(n\alpha)R_\theta \notin \mathcal{F}(X) \Leftrightarrow n\alpha > \cos(\theta) \Leftrightarrow (2\cos(\theta))^{-1} > \cos(\theta)$. Altogether,

$$[\alpha R_\theta \in \mathcal{F}(X) \text{ and } n\alpha R_\theta \notin \mathcal{F}(X)] \Leftrightarrow \cos(\theta) < \frac{1}{2\cos(\theta)} \leq n\cos(\theta) \quad (15a)$$

$$\Leftrightarrow \cos^2(\theta) < \frac{1}{2} \leq n\cos^2(\theta) \quad (15b)$$

$$\Leftrightarrow \cos(\theta) < \frac{1}{\sqrt{2}} \leq \sqrt{n}\cos(\theta) \quad (15c)$$

$$\Leftrightarrow \frac{1}{\sqrt{2n}} \leq \cos(\theta) < \frac{1}{\sqrt{2}}. \quad (15d)$$

Note that (15) has no solution for $n = 1$; however, (15) has solutions for every $n \geq 2$. Because $R_{-\theta} = R_\theta^*$, the result follows with [4, Corollary 4.3] or by arguing along the same lines as above for $R_{-\theta}$. ■

We now obtain the following direct consequence of Lemma 4.3:

Example 4.4. (partial sum property fails for general firmly nonexpansive mappings) *Let $n \in \{2, 3, \dots\}$, and let θ , α , and $R_{\pm\theta}$ be as in Lemma 4.3. Furthermore, set $T_1 := \dots = T_n := \alpha R_\theta$ and $T_{n+1} := \dots = T_{2n} := \alpha R_{-\theta}$. Then each T_i is firmly nonexpansive, $T_1 + \dots + T_{2n} = \text{Id}$, yet $T_1 + \dots + T_n$ is not firmly nonexpansive.*

We conclude this section with another negative result.

Example 4.5. (lack of transitivity for firmly nonexpansive mappings) *Suppose that $X = \mathbb{R}^2$, and set $R := \alpha R_\theta$ and $S := \alpha R_{-\theta}$, where θ and α are as in Example 4.4 for $n = 2$. Then*

$$R \text{ and } S \text{ are firmly nonexpansive,} \quad (16a)$$

$$2R \text{ and } 2S \text{ are not firmly nonexpansive,} \quad (16b)$$

$$2R + 2S = \text{Id.} \quad (16c)$$

Now set

$$T_1 := S, \quad T_2 := R + S, \quad T_3 := 2R + S. \quad (17)$$

Then $T_1 \in \mathcal{F}(X)$ by (16a). Next, (16a) and (16c) imply that $T_3 = \text{Id} - S \in \mathcal{F}(X)$. Since $\mathcal{F}(X)$ is convex (Corollary 2.2), it follows that $T_2 = (T_1 + T_3)/2 \in \mathcal{F}(X)$. Because $T_2 - T_1 = T_3 - T_2 = R \in \mathcal{F}(X)$ by (16a), we have

$$T_1 \preceq T_2 \text{ and } T_2 \preceq T_3. \quad (18)$$

On the other hand, $T_3 - T_1 = 2R \notin \mathcal{F}(X)$ by (16b). Thus,

$$T_1 \not\preceq T_3. \quad (19)$$

Altogether, we deduce that

$$(\mathcal{F}(X), \preceq) \text{ is not transitive.} \quad (20)$$

5 Compatibility with Zarantonello's partial order

Zarantonello introduced in [22, 23] a partial ordering of the set of projectors onto nonempty closed convex cones contained in X via

$$P_C \preceq_Z P_D \Leftrightarrow P_C P_D = P_C. \quad (21)$$

He established various nice properties which we collect in the following result.

Fact 5.1. (Zarantonello) *Let C and D be nonempty closed convex cones in X . Then the following hold:*

- (i) $P_C \preceq_Z P_D \Leftrightarrow P_D - P_C$ is a projector, in which case⁴ $P_D - P_C = P_{D \cap C^\ominus}$.
- (ii) $P_C \preceq_Z P_D \Leftrightarrow [P_C P_D = P_D P_C \text{ and } (\forall x \in X) \langle x, P_C x \rangle \leq \langle x, P_D x \rangle]$.
- (iii) $P_C \preceq_Z P_D \Rightarrow C \subseteq D$.
- (iv) $P_C \preceq_Z P_D \Rightarrow P_C, P_D, P_{C^\ominus}, P_{D^\ominus}$ pairwise commute with their products being the projectors onto the intersection of their ranges ($P_C P_D = P_C P_D = P_{C \cap D}$, etc.).
- (v) Suppose that C and D are subspaces. Then $P_C \preceq_Z P_D \Leftrightarrow C \subseteq D$.

Proof. (i)–(iv): See [22, Lemma 5.12] and [23, page 347]. (v): If $C \subseteq D$, then $D = C \oplus (D \cap C^\perp)$, which implies that $P_D - P_C = P_{D \cap C^\perp}$ is a projector and $P_C \preceq_Z P_D$ by (i). The other implication is (iii). ■

Remark 5.2. *Generalizing (21) and hoping that Fact 5.1(i) holds by just replacing projectors by proximal mappings will not work: indeed, $P_f - P_f = 0 = P_{\{0\}}$ is a proximal map and a projector yet $P_f P_f \neq P_f$.*

Next, let us show that Zarantonello's order is compatible with the order from Definition 3.1:

Lemma 5.3. (compatibility with Zarantonello's order) *Let C and D be nonempty closed convex cones in X . Then $P_C \preceq_Z P_D \Leftrightarrow P_C \preceq P_D$.*

Proof. “ \Rightarrow ”: Assume that $P_C \preceq_Z P_D$. By Fact 5.1(i), $P_D - P_C$ is a projector, hence a proximal mapping and thus firmly nonexpansive. Therefore, $P_C \preceq P_D$. “ \Leftarrow ”: Assume that $P_C \preceq P_D$, i.e., $P_D - P_C$ is firmly nonexpansive. By Theorem 3.5, $P_D - P_C$ is a proximal mapping. Hence, for

$$f := \frac{1}{2}d_C^2 - \frac{1}{2}d_D^2, \quad (22)$$

there exists $h \in \Gamma_0(X)$ such that

$$\nabla f = (\text{Id} - P_C) - (\text{Id} - P_D) = P_D - P_C = P_h. \quad (23)$$

Since P_h is monotone, the function f is convex and so $f \in \Gamma_0(X)$. Now $\nabla f = P_h = (\text{Id} + \partial h)^{-1} \Rightarrow \partial f^* = (\nabla f)^{-1} = \text{Id} + \partial h = \partial(q + h)$. Thus, after integrating and noting that (23) is invariant under adding constants to h , we may and do assume that

$$h = f^* - q. \quad (24)$$

⁴Here $C^\ominus := \{x \in X \mid \sup \langle C, x \rangle = 0\}$ is the polar cone of C .

Furthermore, combining [4, Example 13.3(ii) and Example 13.24(iii)] yields

$$\left(\frac{1}{2}d_C^2\right)^* = q + \iota_{C^\ominus}. \quad (25)$$

Let us now compute f^* at $u \in X$. By [4, Proposition 14.19],

$$f^*(u) = \sup_{v \in \text{dom}((1/2)d_D^2)^*} \left(\left(\frac{1}{2}d_C^2\right)^*(u+v) - \left(\frac{1}{2}d_D^2\right)^*(v) \right) \quad (26a)$$

$$= \sup_{v \in D^\ominus} (q(u+v) + \iota_{C^\ominus}(u+v) - q(v)) \quad (26b)$$

$$= q(u) + \sup_{v \in D^\ominus} (\langle u, v \rangle + \iota_{C^\ominus}(u+v)). \quad (26c)$$

Two cases are now conceivable.

Case 1: $(\exists v \in D^\ominus) u+v \notin C^\ominus$.

Then $f^*(u) = +\infty$ and hence $f^*(u) - q(u) = +\infty$.

Case 2: $u + D^\ominus \subseteq C^\ominus$.

Then

$$f^*(u) - q(u) = \sup_{v \in D^\ominus} \langle u, v \rangle = \iota_{D^\ominus}^*(u) = \iota_{D^\ominus \ominus}(u) = \iota_D(u) \in \{0, +\infty\}. \quad (27)$$

Altogether, $h = f^* - q$ takes only values in $\{0, +\infty\}$, i.e., h is an *indicator function*. Thus, P_h must be a projector⁵. Therefore, by Fact 5.1(i), $P_C \preceq_Z P_D$. \blacksquare

6 Compatibility with the Loewner order via resolvents

In this section, we assume that

$$X = \mathbb{S}^n := \{A \in \mathbb{R}^{n \times n} \mid A = A^*\} \quad (28)$$

is the finite-dimensional Hilbert space of all real symmetric matrices of size $n \times n$ with the inner product $\langle A, B \rangle$ being the trace of AB . We shall focus on the closed convex cone of *positive semidefinite matrices*:

$$\mathbb{S}_+^n := \{A \in \mathbb{S}^n \mid (\forall x \in \mathbb{R}^n) \langle x, Ax \rangle \geq 0\} = \{A \in \mathbb{S}^n \mid A \text{ is monotone}\}. \quad (29)$$

Let A and B be in \mathbb{S}_+^n . The classical *Loewner* (or Löwner) *order* [14] states

$$B \preceq_L A \quad :\Leftrightarrow \quad A - B \in \mathbb{S}_+^n, \text{ i.e., } A - B \text{ is monotone.} \quad (30)$$

⁵We may obtain additional information as follows. Suppose first, as in *Case 2*, that $u + D^\ominus \subseteq C^\ominus$. This case must occur since h is proper. (In passing, note that this precisely states that u is in the so-called *star-difference* $C^\ominus \star D^\ominus$; see [12].) Since $0 \in D^\ominus$, it is clear that $u \in C^\ominus$. On the other hand, since we are working with *cones*, we have $(\forall \varepsilon > 0) \varepsilon u + D^\ominus = \varepsilon(u + D^\ominus) \subseteq \varepsilon C^\ominus = C^\ominus$. Letting $\varepsilon \rightarrow 0^+$, we deduce that $D^\ominus \subseteq C^\ominus$ and thus $C \subseteq D$ as is also guaranteed by Fact 5.1(i). If conversely $u \in C^\ominus$, then $u + D^\ominus \subseteq u + C^\ominus \subseteq C^\ominus$. Altogether, we have shown that u is as in *Case 2* if and only if $u \in C^\ominus$. Therefore, $h = \iota_{D \cap C^\ominus}$, which is consistent with Fact 5.1(i).

Passing to resolvents, we have

$$B \preceq_L A \Leftrightarrow \text{Id} + B \preceq_L \text{Id} + A \Leftrightarrow (\text{Id} + A)^{-1} \preceq_L (\text{Id} + B)^{-1} \Leftrightarrow J_A \preceq_L J_B. \quad (31)$$

The question now arises whether the Loewner order for resolvents is compatible with our order from Definition 3.1. Clearly,

$$J_A \preceq J_B \Rightarrow J_B - J_A \text{ is monotone} \Leftrightarrow J_A \preceq_L J_B. \quad (32)$$

Conversely, assume that $J_A \preceq_L J_B$, i.e., $J_B - J_A$ is monotone. On the other hand, $J_B - J_A$ is nonexpansive by Lemma 2.3. Altogether, by Fact 2.7, $J_B - J_A$ is firmly nonexpansive, i.e., $J_A \preceq J_B$. In summary,

$$B \preceq_L A \Leftrightarrow J_A \preceq_L J_B \Leftrightarrow J_A \preceq J_B, \quad (33)$$

which shows that *the Loewner order and our order are compatible*. (In passing, we note that the comments in this section have extensions to self-adjoint operators on Hilbert space.)

7 A connection to Moreau's order

In his seminal work [16], Moreau introduced an order of $\Gamma_0(X)$ via

$$g \preceq_M f \quad :\Leftrightarrow \quad (\exists h \in \Gamma_0(X)) \quad f = g + h. \quad (34)$$

In fact, using this notation, we can write the equivalence of (iii) and (iv) in Fact 2.4, which was first observed by Moreau [16], more succinctly as $f \preceq_M q \Leftrightarrow q \preceq_M f^*$. To make the connection with our order, let us take f and g from $\Gamma_0(X)$. Using Corollary 2.6 and Theorem 3.5, we have the following equivalences:

$$\text{env}(g) \preceq_M \text{env}(f) \Leftrightarrow \text{env}(f) - \text{env}(g) \text{ is convex} \Leftrightarrow \text{env}(g^*) - \text{env}(f^*) \text{ is convex} \quad (35a)$$

$$\Leftrightarrow P_g - P_f \text{ is monotone} \Leftrightarrow \text{env}(f) - \text{env}(g) \in M_0(X) \quad (35b)$$

$$\Leftrightarrow P_g - P_f \in \mathcal{P}(X) \Leftrightarrow P_f \preceq P_g. \quad (35c)$$

Hence, *our order is compatible with Moreau's order* when restricted to $M_0(X)$, the set of envelope functions on X .

8 Ordering monotone operators and convex functions

The classical bijection between the maximally monotone operators on X and the firmly nonexpansive mappings on X , introduced by Minty [15] (see also [9]), is

$$\mathcal{M}(X) \rightarrow \mathcal{F}(X): A \mapsto J_A = (\text{Id} + A)^{-1}. \quad (36)$$

We thus define a new binary relation on $\mathcal{M}(X)$ via

$$B \preceq A \quad :\Leftrightarrow \quad J_A \preceq J_B. \quad (37)$$

For instance, we have $N_X = 0 \preceq A \preceq N_{\{0\}}$ by Lemma 3.2(ii) while Lemma 3.2(iv) gives a statement relating to the resolvent average introduced in [2]. The results in Section 6 show that when $X = \mathbb{R}^n$ and we consider \mathcal{S}_+^n , which is a subset of $\mathcal{M}(X)$, then these notions are compatible (see (33)).

Furthermore, we can use this binary relation on $\mathcal{M}(X)$ to define a new binary relation on $\Gamma_0(X)$ via

$$g \preceq f \quad :\Leftrightarrow \quad \partial g \preceq \partial f. \quad (38)$$

With these definitions and using (35), we have the equivalences

$$g \preceq f \Leftrightarrow \partial g \preceq \partial f \Leftrightarrow J_{\partial f} \preceq J_{\partial g} \Leftrightarrow P_f \preceq P_g \Leftrightarrow \text{env}(g) \preceq_M \text{env}(f) \quad (39)$$

which show that our binary relation on $\Gamma_0(X)$ plays along nicely with Moreau's order.

Let us present another example. Let A and B be in \mathcal{S}_+^n and define the quadratic forms

$$(\forall x \in \mathbb{R}^n) \quad q_A(x) := \frac{1}{2} \langle x, Ax \rangle \quad \text{and} \quad q_B(x) := \frac{1}{2} \langle x, Bx \rangle. \quad (40)$$

Then $\nabla q_A = A$ and $\nabla q_B = B$. Using (31), we see that

$$q_B \leq q_A \text{ (pointwise)} \Leftrightarrow 0 \leq q_{A-B} \Leftrightarrow A - B \in \mathcal{S}_+^n \Leftrightarrow B \preceq_L A \Leftrightarrow J_A \preceq J_B \quad (41a)$$

$$\Leftrightarrow B \preceq A \Leftrightarrow \nabla q_B \preceq \nabla q_A \Leftrightarrow q_B \preceq q_A \Leftrightarrow P_{q_A} \preceq P_{q_B} \quad (41b)$$

$$\Leftrightarrow \text{env}(q_B) \preceq_M \text{env}(q_A). \quad (41c)$$

This nicely illustrates the connections between the various orders considered in this paper.

9 New partial orders

In our final section, we introduce a quotient space construction which remedies the lack of anti-symmetry observed in Example 3.4.

We start with a simple but useful result.

Lemma 9.1. *Let $T \in \mathcal{F}(X)$ be such that $-T \in \mathcal{F}(X)$. Then T is a constant mapping.*

Proof. Take x and y from X . Then

$$\|Tx - Ty\|^2 \leq \langle Tx - Ty, x - y \rangle \quad \text{and} \quad \|(-T)x - (-T)y\|^2 \leq -\langle Tx - Ty, x - y \rangle. \quad (42)$$

Thus,

$$0 \leq \|Tx - Ty\|^2 \quad (43a)$$

$$\leq \min \{ \langle Tx - Ty, x - y \rangle, -\langle Tx - Ty, x - y \rangle \} \quad (43b)$$

$$= -|\langle Tx - Ty, x - y \rangle| \quad (43c)$$

$$\leq 0. \quad (43d)$$

Therefore, $Tx - Ty = 0$. ■

We now define a binary relation on $\mathcal{F}(X)$ via

$$T_1 \sim T_2 \quad :\Leftrightarrow \quad T_2 - T_1 \text{ is a constant mapping.} \quad (44)$$

It is straightforward to check that $(\mathcal{F}(X), \sim)$ is an *equivalence relation*. Denote the corresponding *quotient set* by

$$[\mathcal{F}(X)] := \mathcal{F}(X) / \sim. \quad (45)$$

Now define a binary relation on $[\mathcal{F}(X)]$ by

$$[T_1] \preceq [T_2] \quad :\Leftrightarrow \quad T_1 \preceq T_2. \quad (46)$$

Then $([\mathcal{F}(X)], \preceq)$ is *reflexive*; moreover, by Lemma 9.1, $([\mathcal{F}(X)], \preceq)$ is *antisymmetric*. If we restrict to proximal mappings, then $([\mathcal{P}(X)], \preceq)$ is also *transitive* by Theorem 3.6. In summary,

$$([\mathcal{P}(X)], \preceq) \text{ is a partially ordered set.} \quad (47)$$

Finally, let us investigate (44) from the view point of maximally monotone operators via (36).

Lemma 9.2. *Let A and B be in $\mathcal{M}(X)$ and let $c \in X$. Then the following are equivalent:*

- (i) $(\forall x \in X) J_B x = c + J_A x$.
- (ii) $(\forall x \in X) Bx = -c + A(x - c)$.
- (iii) $\text{gra } B = (c, -c) + \text{gra } A$.

Proof. “(i) \Rightarrow (iii)”: By assumption, $(\forall x \in X) (J_B x, x - J_B x) = (c, -c) + (J_A x, x - J_A x)$. Using the Minty parametrization (see [4, (23.18)]) of the graph, we see that $\text{gra } B = (c, -c) + \text{gra } A$. The other implications are proved similarly. ■

In view of Lemma 9.2, the equivalence relation (44) in $\mathcal{F}(X)$ gives rise to the following equivalence relation on $\mathcal{M}(X)$:

$$A \sim B \quad :\Leftrightarrow \quad (\exists c \in X)(\forall x \in X) Bx = -c + A(x - c). \quad (48)$$

In turn, we can “integrate” (48) to obtain the following equivalence relation on $\Gamma_0(X)$:

$$f \sim g \quad :\Leftrightarrow \quad (\exists c \in X)(\exists \gamma \in \mathbb{R})(\forall x \in X) g(x) = f(x - c) - \langle c, x \rangle + \gamma. \quad (49)$$

The last equivalence relation induces a quotient set $[\Gamma_0(X)] := \Gamma_0(X) / \sim$. Interpreting (47) in this setting, we obtain the following result.

Theorem 9.3. *Equip the quotient set $[\Gamma_0(X)]$ with the binary relation*

$$[g] \preceq [f] \quad :\Leftrightarrow \quad [P_f] \preceq [P_g] \Leftrightarrow P_f \preceq P_g. \quad (50)$$

Then

$$([\Gamma_0(X)], \preceq) \text{ is a partially ordered set.} \quad (51)$$

Acknowledgments

Sedi Bartz was supported by a postdoctoral fellowship of the Pacific Institute for the Mathematical Sciences and by NSERC grants of Heinz Bauschke and Xianfu Wang. Heinz Bauschke was partially supported by the Canada Research Chair program and by the Natural Sciences and Engineering Research Council of Canada. Xianfu Wang was partially supported by the Natural Sciences and Engineering Research Council of Canada.

References

- [1] J.-B. Baillon and G. Haddad, Quelques propriétés des opérateurs angle-bornés et n -cycliquement monotones, *Israel Journal of Mathematics* 26 (1977), 137–150.
- [2] S. Bartz, H.H. Bauschke, S.M. Moffat, and X. Wang, The resolvent average of monotone operators: dominant and recessive properties, *SIAM Journal on Optimization* 26 (2016), 602–634.
- [3] H.H. Bauschke and P.L. Combettes, The Baillon–Haddad theorem revisited, *Journal of Convex Analysis* 17 (2010), 781–787.
- [4] H.H. Bauschke and P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, Springer, 2011.
- [5] J.M. Borwein, Fifty years of maximal monotonicity, *Optimization Letters* 4 (2010), 473–490.
- [6] H. Brezis, *Opérateurs Maximaux Monotones et Semi-Groupes de Contractions dans les Espaces de Hilbert*, North-Holland/Elsevier, 1973.
- [7] R.E. Bruck and S. Reich, Nonexpansive projections and resolvents of accretive operators in Banach spaces, *Houston Journal of Mathematics* 3 (1977), 459–470.
- [8] R.S. Burachik and A.N. Iusem, *Set-Valued Mappings and Enlargements of Monotone Operators*, Springer, 2008.
- [9] J. Eckstein and D.P. Bertsekas, On the Douglas–Rachford splitting method and the proximal point algorithm for maximal monotone operators, *Mathematical Programming* 55 (1992), 293–318.
- [10] K. Goebel and W.A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge University Press, 1990.
- [11] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Marcel Dekker, 1984.
- [12] J.-B. Hiriart-Urruty and C. Lemaréchal, *Convex Analysis and Minimization Algorithms I*, Springer, 1993.
- [13] J.-B. Hiriart-Urruty and C. Lemaréchal, *Convex Analysis and Minimization Algorithms II*, Springer, 1993.
- [14] K. Löwner, Über monotone Matrixfunktionen, *Mathematische Zeitschrift* 38 (1934), 177–216.
- [15] G.J. Minty, Monotone (nonlinear) operators in Hilbert spaces, *Duke Mathematical Journal* 29 (1962), 341–346.
- [16] J.-J. Moreau, Proximité et dualité dans un espace hilbertien *Bulletin de la Société Mathématique de France* 93 (1965), 273–299.
- [17] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, 1970.
- [18] R.T. Rockafellar and R. J-B Wets, *Variational Analysis*, Springer, corrected 3rd printing, 2009.
- [19] S. Simons, *Minimax and Monotonicity*, Springer, 1998.
- [20] S. Simons, *From Hahn-Banach to Monotonicity*, Springer, 2008.
- [21] C. Zălinescu, *Convex Analysis in General Vector Spaces*, World Scientific Publishing, 2002.
- [22] E.H. Zarantonello, Projections on convex sets in Hilbert space and spectral theory I: Projections on convex sets, in *Contributions to Nonlinear Functional Analysis*, E.H. Zarantonello (editor), Mathematics Research Center Publication 27, Academic Press, New York-London, 1971, pages 237–341.
- [23] E.H. Zarantonello, Projections on convex sets in Hilbert space and spectral theory II: Spectral theory, in *Contributions to Nonlinear Functional Analysis*, E.H. Zarantonello (editor), Mathematics Research Center Publication 27, Academic Press, New York-London, 1971, pages 343–424.
- [24] E. Zeidler, *Nonlinear Functional Analysis and Its Applications II/A: Linear Monotone Operators*, Springer, 1990.
- [25] E. Zeidler, *Nonlinear Functional Analysis and Its Applications II/B: Nonlinear Monotone Operators*, Springer, 1990.