

RAMIFIED OPTIMAL TRANSPORTATION WITH PAYOFF ON THE BOUNDARY

QINGLAN XIA

Department of Mathematics
University of California at Davis
Davis, CA 95616, USA

SHAOFENG XU

School of Economics
Renmin University of China
Beijing, 100872, China

ABSTRACT. This paper studies a variant of ramified/branched optimal transportation problems. Given the distributions of production capacities and market sizes, a firm looks for an allocation of productions over factories, a distribution of sales across markets, and a transport path that delivers the product to maximize its profit. Mathematically, given any two measures μ and ν on X , and a payoff function h , the planner wants to minimize $\mathbf{M}_\alpha(T) - \int_X hd(\partial T)$ among all transport paths T from $\tilde{\mu}$ to $\tilde{\nu}$ with $\tilde{\mu} \leq \mu$ and $\tilde{\nu} \leq \nu$, where \mathbf{M}_α is the standard cost functional used in ramified transportation. After proving the existence result, we provide a characterization of the boundary measures of the optimal solution. They turn out to be the original measures restricted on some Borel subsets up to a Delta mass on each connected component. Our analysis further finds that as the boundary payoff increases, the corresponding solution of the current problem converges to an optimal transport path, which is the solution of the standard ramified transportation.

1. INTRODUCTION

1.1. The ROTPB problem. Transportation is an important force shaping the spatial distribution of economic activities. Consider a firm that produces and sells a product in various regions. Given the locations and capacities of these regions and the associated production costs and sale prices of the product, the firm looks for a distribution of productions over factories, a distribution of sales across markets, and a transport path that delivers the product to maximize its profit. The firm's optimal plan over productions and sales depends on its choice of transport path, and vice versa. The interactions between location and transport choices, however, often render these problems difficult to analyze.

In this paper, we address some of these interactions in the framework of the ramified optimal transportation. More precisely, we consider the following resource allocation problem: Let μ and ν be two Radon measures on a convex compact subset X of the Euclidean space \mathbb{R}^m , \mathbf{M}_α be the standard cost functional used in ramified transportation [28] for $\alpha \in [0, 1)$ and h be a continuous function on the support of the signed measure $\nu - \mu$. We consider the problem:

2010 *Mathematics Subject Classification.* 49Q20, 49Q10, 90B10.

Key words and phrases. branch transportation, economies of scale, optimal transport path, branching structure, rectifiable current, partial transportation.

Problem (ROTPB(μ, ν)). *Minimize*

$$(1.1) \quad \mathbf{E}_\alpha^h(T) := \mathbf{M}_\alpha(T) - \int_X hd(\partial T)$$

among all rectifiable 1-current T with $\partial T \preceq \nu - \mu$ as signed measures.¹

In the context of the above example, measures μ and ν represent, respectively, the distributions of production capacities and market sizes. The function h represents the payoff associated with moving mass from μ to ν , and it captures the production cost of the product over μ and its sale price over ν . The firm aims to maximize its profit defined as sale revenues minus costs involved in transportation and production. We call this problem as *Ramified optimal transportation with payoff on the boundary* (ROTPB).

1.2. Background. This paper is related to the literature of optimal transport problems which concerns efficient mass transportation. These problems are studied early on by Monge and Kantorovich, and has been extensively analyzed in recent years. Classical references can be found in the books [22, 23] by Villani, [18] by Santambrogio, and the user's guide [1] by Ambrosio and Gigli. Our paper is most closely related to the ramified optimal transportation (ROT) (also called branched transportation) literature, which models branching transport structures thanks to the efficiency in group transportation. In contrast to the Monge-Kantorovich problems where the transportation cost is solely determined by a transport map, the cost in ramified transport problems is determined by the actual transport path. The Eulerian formulation of the ROT problem is proposed by the first author in [24], with related motivations, set-up and applications surveyed in [28]. An equivalent Lagrangian formulation of the problem is established by Maddalena, Morel, and Solimini in [13]. One may refer to [2] for detailed discussions of the research in this direction. Some interesting recent developments on ROT can be found for instance in [3, 5, 7, 12, 17, 19].

Our paper differentiates itself from the existing ROT literature in two main regards. First, in the literature both measures μ and ν are fixed and of equal mass, and the problem only involves finding a cost-minimizing transport path. By contrast, the planner in this paper optimizes over all possible combinations of $(\tilde{\mu}, \tilde{\nu})$ with $\tilde{\mu} \leq \mu$, $\tilde{\nu} \leq \nu$ and $|\tilde{\mu}| = |\tilde{\nu}|$. Similar kind of optimal partial mass transport has been studied for instance by Caffarelli and McCann [4] and also Figalli [11] for the scenario of Monge-Kantorovich problems with a particular attention to the quadratic cost. Second, the planner faces a reward for relocating mass at the boundary, and thus the solution relies on the payoff function h . This element has been absent in the literature up to our best knowledge.

1.3. Main results. Our main results include three parts: the existence theorem (Theorem 3.1), the characterization theorems (Theorem 1.1, Theorem 4.18), and the approximation theorem (Theorem 5.4).

We first prove the existence of an \mathbf{E}_α^h -minimizer T^* for the ROTPB(μ, ν) problem in Theorem 3.1. This optimal solution T^* is an α -optimal transport path of finite \mathbf{M}_α cost from μ^* to ν^* for some measures $\mu^* \leq \mu$ and $\nu^* \leq \nu$. As such, T^* automatically inherits many nice geometric properties of optimal transport paths as described previously in [28]. We next characterize the optimal allocation measures μ^* and ν^* . In the finite atomic case, we show

¹The notation \preceq is introduced in (2.4).

Theorem 1.1. *Suppose μ and ν are two finite atomic measures on X , $0 < \alpha < 1$, and $T^* \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem. Let $\{K_k : k = 1, 2, \dots, \ell\}$ be the set of the connected components of the support of T^* . Then, for each $k = 1, 2, \dots, \ell$, it holds that*

$$(1.2) \quad \mu^* \llcorner K_k = \mu \llcorner K_k - m_k \delta_{p_k} \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k - n_k \delta_{q_k},$$

for some points $p_k \in K_k \cap \text{spt}(\mu^*)$ and $q_k \in K_k \cap \text{spt}(\nu^*)$ with

$$m_k := \max\{\mu(K_k) - \nu(K_k), 0\} \text{ and } n_k := \max\{\nu(K_k) - \mu(K_k), 0\}.$$

As a result, we have the decomposition

$$(1.3) \quad \mu^* = \mu \llcorner A - \mathbf{a} \text{ and } \nu^* = \nu \llcorner B - \mathbf{b},$$

for $A = \text{spt}(\mu^*)$, $B = \text{spt}(\nu^*)$, and

$$(1.4) \quad \mathbf{a} = \sum_{k=1}^{\ell} m_k \delta_{p_k}, \quad \mathbf{b} = \sum_{k=1}^{\ell} n_k \delta_{q_k}.$$

Note that in equation (1.2), at least one of m_k and n_k is zero for each k . The equation says that on each connected component K_k , all existing resources in the optimal allocation source measure μ^* will be used up, and all demands in the optimal allocation destination measure ν^* will be met with at most one exception at either a source node or a destination node. There are three scenarios:

- In the balanced case where $\mu(K_k) = \nu(K_k)$, then

$$\mu^* \llcorner K_k = \mu \llcorner K_k \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k.$$

All source and destination nodes are fully in use.

- In the over-supply case where $\mu(K_k) > \nu(K_k)$, then

$$\mu^* \llcorner K_k = \mu \llcorner K_k - (\mu(K_k) - \nu(K_k)) \delta_{p_k} \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k.$$

All source nodes excluding the one at p_k and all destination nodes are fully in use.

- In the over-demand case where $\mu(K_k) < \nu(K_k)$, then

$$\mu^* \llcorner K_k = \mu \llcorner K_k \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k - (\nu(K_k) - \mu(K_k)) \delta_{q_k}.$$

All source nodes and all destination nodes except for the one at q_k are fully in use.

In Theorem 4.18, we further extend the results of Theorem 1.1 to general cases.

The third part of the main results highlights an important implication of the current study for solving an optimal transport path. We consider a version of ROTPB problems, where the measures μ and ν are disjointly supported and the payoff function h_c takes a constant value $2c$ on the support of ν , and vanishes on the support of μ . In the early example, the parameter c represents (half of) the gap between the sale price and the production cost, and it effectively determines the payoff from relocating a unit of mass. Intuitively, the larger the payoff, the more incentive the planner has to relocate mass from sources to destinations. When the payoff is sufficiently large, it is in the best interest of the planner to move as much mass as possible. We prove in Theorem 5.4 that an optimal transport path, which solves the standard ramified transportation problem, can be obtained as a limit of the solutions to a sequence of ROTPB problems associated with a series of increasing boundary payoff. This finding thus provides a novel perspective for approximating an optimal transport path.

2. PRELIMINARIES

2.1. Basic notations in geometric measure theory. We first recall some terminology about rectifiable currents as in [10] or [20].

Let $\Omega \subseteq \mathbb{R}^m$ be an open domain and for any integer $k \geq 0$ let $\mathcal{D}^k(\Omega)$ be the set of all C^∞ differential k -forms in Ω with compact support with the usual Fréchet topology [10]. A k -dimensional current S in Ω is a continuous linear functional on $\mathcal{D}^k(\Omega)$. Denote $\mathcal{D}_k(\Omega)$ as the set of all k -dimensional currents in Ω . The *mass* of a current $T \in \mathcal{D}_k(\Omega)$ is defined by

$$\mathbf{M}(T) := \sup\{|T(\omega)| : \|\omega\| \leq 1, \omega \in \mathcal{D}^k(\Omega)\}.$$

Motivated by the Stokes' theorem, the *boundary* of a current $S \in \mathcal{D}_k(\Omega)$ for $k \geq 1$ is the current ∂S in $\mathcal{D}_{k-1}(\Omega)$ defined by

$$\partial S(\omega) := S(d\omega)$$

for any $\omega \in \mathcal{D}^{k-1}(\Omega)$. A current $T \in \mathcal{D}_k(\Omega)$ is called *normal* if $\mathbf{M}(T) + \mathbf{M}(\partial T) < +\infty$. A sequence of currents $\{S_i\}$ in $\mathcal{D}_k(\Omega)$ is said to be weakly convergent to another current $S \in \mathcal{D}_k(\Omega)$, denoted by $S_i \rightharpoonup S$, if

$$S_i(\omega) \rightarrow S(\omega)$$

for any $\omega \in \mathcal{D}^k(\Omega)$.

As in [20], a subset $M \subseteq \mathbb{R}^m$ is called (countably) k -rectifiable if $M = \bigcup_{i=0}^{\infty} M_i$, where $\mathcal{H}^k(M_0) = 0$ under the k -dimensional Hausdorff measure \mathcal{H}^k and each M_i , for $i = 1, 2, \dots$, is a subset of an k -dimensional C^1 submanifold in \mathbb{R}^m . A *rectifiable k -current* S is a k -dimensional current coming from an oriented k -rectifiable set with multiplicities. More precisely, $S \in \mathcal{D}_k(\Omega)$ is a *rectifiable k -current* if it can be expressed as

$$S(\omega) = \int_M \langle \omega(x), \xi(x) \rangle \theta(x) d\mathcal{H}^k(x), \quad \forall \omega \in \mathcal{D}^k(\Omega)$$

where

- M is an \mathcal{H}^k measurable and k -rectifiable subset of Ω .
- θ is an $\mathcal{H}^k \llcorner M$ integrable positive function and is called the multiplicity function of S .
- $\xi : M \rightarrow \Lambda_k(\mathbb{R}^m)$ is an \mathcal{H}^k measurable unit tangent vector field on M and is called the orientation of S .

The rectifiable current S described as above is often denoted by

$$S = \underline{\underline{\tau}}(M, \theta, \xi).$$

In this case, the mass of S is expressed as

$$\mathbf{M}(S) = \int_M \theta(x) d\mathcal{H}^k(x).$$

Since θ is $\mathcal{H}^k \llcorner M$ integrable, each rectifiable current S here is assumed to have finite mass.

2.2. Basic notations in ramified optimal transportation. Let X be a convex compact subset of the Euclidean space \mathbb{R}^m . The ramified optimal transport problem (also called branched optimal transportation problem in the literature) considers the following Plateau-type problem:

Problem (ROT). *Given two (positive) measures μ^+ and μ^- on X of equal mass and $\alpha < 1$, minimize*

$$\mathbf{M}_\alpha(T) := \int_M \theta^\alpha d\mathcal{H}^1.$$

among all rectifiable 1-current $T = \underline{\tau}(M, \theta, \xi)$ in \mathbb{R}^m with $\partial T = \mu^- - \mu^+$ in the sense of distributions.

Each rectifiable 1-current $T = \underline{\tau}(M, \theta, \xi)$ such that $\partial T = \mu^- - \mu^+$ is called a *transport path* from μ^+ to μ^- . Let

$$\text{Path}(\mu^+, \mu^-) = \{T \text{ is a rectifiable 1-current} : \partial T = \mu^- - \mu^+\}$$

be the collection of all transport paths from μ^+ to μ^- .

For the ROT problem, the existence of an \mathbf{M}_α -minimizer in $\text{Path}(\mu^+, \mu^-)$ is shown in [24]. Each \mathbf{M}_α -minimizer is called an α -optimal transport path. One shall note that for some combinations of exponent α and pair of measures μ^\pm , it is possible the \mathbf{M}_α cost of any transport path $T \in \text{Path}(\mu^+, \mu^-)$ is infinite, and thus the existence of a solution to the ROT problem is trivial in that case.

When $1 - \frac{1}{m} < \alpha < 1$, it is shown in [24] that for any pair of measures μ^\pm of equal (finite) mass there exists an α -optimal transport path of finite \mathbf{M}_α -cost from μ^+ to μ^- . Moreover, a distance is defined by setting

$$(2.1) \quad d_\alpha(\mu^+, \mu^-) := \min\{\mathbf{M}_\alpha(T) : \partial T = \mu^- - \mu^+\}$$

between μ^+ and μ^- . By [24, Theorem 3.1], it holds that

$$(2.2) \quad d_\alpha(\mu^+, \mu^-) \leq C_{m,\alpha} \text{diam}(X) \|\mu^+\|^\alpha,$$

where the constant

$$(2.3) \quad C_{m,\alpha} = \frac{\sqrt{m}}{2(2^{1-m(1-\alpha)} - 1)}.$$

In general, the existence of finite cost α -optimal transport path between μ^+ and μ^- depends on the dimensional information of the measures (see [8], [26]). In [26], the d_α -metric is defined on the space of finite atomic probability measures for any real number $\alpha < 1$.

The following notations are also employed in the analysis:

- Let μ and ν be two (positive) measures on X . We say $\mu \leq \nu$ if $\nu - \mu$ is still a (positive) measure on X . In this case, we say that μ is *feasible* relative to ν .
- Let $\mu_1 = \mu_1^+ - \mu_1^-$ and $\mu_2 = \mu_2^+ - \mu_2^-$ be the Jordan decompositions of two signed measures. We say

$$(2.4) \quad \mu_1 \preceq \mu_2$$

if $\mu_1^+ \leq \mu_2^+$ and $\mu_1^- \leq \mu_2^-$.

- For any signed measure $\bar{\mu}$, let $\text{spt}(\bar{\mu})$ denote its support, $\|\bar{\mu}\|$ denote its total variation, and $|\bar{\mu}|$ denote its total variation measure.

For each rectifiable 1-current T , its boundary ∂T can be viewed as a signed measure.

3. THE ROTPB PROBLEM

This section analyzes the spatial resource allocation problem ROTPB(μ, ν) as stated in Introduction. For simplicity, when both the parameter α and the function h are clear from the context, we simply write \mathbf{E}_α^h given in (1.1) as \mathbf{E} .

The ROTPB(μ, ν) problem is indeed a double-minimizing problem

$$\min \left\{ \min \left\{ \mathbf{E}_\alpha^h(T) : \partial T = \tilde{\nu} - \tilde{\mu} \right\} : \tilde{\mu} \leq \mu, \tilde{\nu} \leq \nu \text{ with } \|\tilde{\mu}\| = \|\tilde{\nu}\| \right\}.$$

For each fixed $\tilde{\mu} \leq \mu, \tilde{\nu} \leq \nu$ with $\|\tilde{\mu}\| = \|\tilde{\nu}\|$, the inner minimization problem

$$\min \left\{ \mathbf{E}_\alpha^h(T) = \mathbf{M}_\alpha(T) - \int_X h d(\partial T) : \partial T = \tilde{\nu} - \tilde{\mu} \right\}$$

can be re-written as

$$(3.1) \quad \min \left\{ \mathbf{M}_\alpha(T) : \partial T = \tilde{\nu} - \tilde{\mu} \right\} - \int_X h d\tilde{\nu} + \int_X h d\tilde{\mu}.$$

Thus, under the d_α metric as given in (2.1), the ROTPB(μ, ν) problem can also be expressed as: Minimize

$$\mathbf{E}(\tilde{\mu}, \tilde{\nu}) := d_\alpha(\tilde{\mu}, \tilde{\nu}) - \int_X h d\tilde{\nu} + \int_X h d\tilde{\mu}$$

among all feasible measures $\tilde{\mu} \leq \mu$ and $\tilde{\nu} \leq \nu$ with $\|\tilde{\mu}\| = \|\tilde{\nu}\|$.

From the perspective of the firm in the example given in Introduction, the ROTPB(μ, ν) problem can be interpreted as follows. Given the distributions of production capacities (μ) and market sizes (ν), the firm chooses an operation plan $\tilde{\mu} \leq \mu$ and $\tilde{\nu} \leq \nu$ to minimize the total costs incurred in production ($\int_X h d\tilde{\mu}$) and transportation ($d_\alpha(\tilde{\mu}, \tilde{\nu})$) net the sale revenue ($\int_X h d\tilde{\nu}$).

We now state the existence theorem for the ROTPB(μ, ν) problem.

Theorem 3.1 (Existence). *Let μ and ν be two Radon measures on X , $0 \leq \alpha < 1$ and h be a continuous function on the support of the signed measure $\nu - \mu$. Then there exists a rectifiable 1-current T^* of finite \mathbf{M}_α cost that minimizes*

$$\mathbf{E}_\alpha^h(T) := \mathbf{M}_\alpha(T) - \int_X h d(\partial T)$$

among all rectifiable 1-current T with $\partial T \preceq \nu - \mu$ as signed measures.

Remark 3.2. By the Jordan decomposition theorem, for any signed measure $\bar{\mu}$, there exists a unique positive measures μ^+ and μ^- such that $\bar{\mu} = \mu^+ - \mu^-$ and $\mu^+ \perp \mu^-$. Thus, without loss of generality, we may assume that μ and ν are mutually singular when studying the ROTPB(μ, ν) problem.

Proof. We prove this result by using the direct method of calculus of variations. Let $\{T_i\}$ be any \mathbf{E} -minimizing sequence of rectifiable 1-currents. That is,

$$\lim_{i \rightarrow \infty} \mathbf{E}(T_i) = \inf \left\{ \mathbf{E}(T) : \partial T \preceq \nu - \mu \right\},$$

and $\partial T_i \preceq \nu - \mu$ for each i . With no loss of generality, we may assume $\mathbf{E}(T_i) \leq \mathbf{E}(0) = 0$. Thus,

$$(3.2) \quad \mathbf{M}_\alpha(T_i) = \mathbf{E}(T_i) + \int_X h d(\partial T_i) \leq \int_X h d(\partial T_i) \leq \int_X |h| d(|\nu - \mu|) < \infty$$

as h is continuous on the compact set $\text{spt}(\nu - \mu)$, the support of $\nu - \mu$. Now suppose $T_i \in \text{Path}(\tilde{\mu}_i, \tilde{\nu}_i)$. Since $\mathbf{M}_\alpha(T_i)$ is finite, there exists an \mathbf{M}_α -minimizer with finite cost for the minimization problem

$$\min \{ \mathbf{M}_\alpha(T) : \partial T = \partial T_i \}.$$

Note this minimizer is also an \mathbf{E} -minimizer for the inner minimization problem (3.1) with $\tilde{\mu} = \tilde{\mu}_i$ and $\tilde{\nu} = \tilde{\nu}_i$. Without loss of generality, we may assume that T_i is such a minimizer, which is an α -optimal transport path of finite cost. By (3.2), the sequence $\{ \mathbf{M}_\alpha(T_i) \}$ is bounded. Employing Lemma 3.3 below shows that the sequence $\{ \mathbf{M}(T_i) \}$ is also bounded. As a result, we get a sequence of normal 1-currents $\{ T_i \}$ with equi-bounded mass and boundary mass. By the compactness of normal 1-currents ([10]), and taking a subsequence if necessary, we may assume that the sequence $\{ T_i \}$ converges to a normal 1-current T^* with respect to flat convergence. Since \mathbf{M}_α is lower semi-continuous with respect to flat convergence ([6, 14]), we have

$$\mathbf{M}_\alpha(T^*) \leq \liminf_{i \rightarrow \infty} \mathbf{M}_\alpha(T_i) < \infty.$$

According to the rectifiability theorem (e.g., Theorem 2.7 in [25]), finite mass and finite \mathbf{M}_α mass together imply that T^* is also 1-rectifiable. Since $\{ T_i \}$ converges to T^* in flat convergence, the sequence $\{ \partial T_i \}$ is weak-* convergent to ∂T^* as signed measures.

Since h is continuous on the support $\text{spt}(\nu - \mu)$, $\text{spt}(\partial T_i) \subseteq \text{spt}(\nu - \mu)$, and ∂T_i is weak-* convergent to ∂T^* , we have

$$\int_X h d(\partial T) = \lim_{i \rightarrow \infty} \int_X h d(\partial T_i).$$

As a result,

$$\mathbf{E}(T^*) = \mathbf{M}_\alpha(T^*) - \int_X h d(\partial T^*) \leq \liminf_{i \rightarrow \infty} \{ \mathbf{M}_\alpha(T_i) - \int_X h d(\partial T_i) \} = \lim_{i \rightarrow \infty} \mathbf{E}(T_i).$$

When each $\partial T_i \leq \nu - \mu$, its limit $\partial T^* \leq \nu - \mu$ holds as well. This shows that T^* is a solution to the ROTPB(μ, ν) problem. \square

The proof of the theorem takes advantage of the following lemma:

Lemma 3.3. *Suppose T is an α -optimal transport path with $\mathbf{M}_\alpha(T) < \infty$, then*

$$(3.3) \quad \mathbf{M}(T) \leq \left(\frac{\mathbf{M}(\partial T)}{2} \right)^{1-\alpha} \mathbf{M}_\alpha(T).$$

Proof. Suppose $T = \underline{T}(M, \theta, \xi)$ is an α -optimal transport path from μ^+ to μ^- , where $\partial T = \mu^- - \mu^+$ is the Jordan decomposition of ∂T as a signed measure. Since T is an α -optimal transport path of finite cost, it follows (from (4.13) for instance) that $\theta(x) \leq \mu^+(X) = \frac{1}{2} \mathbf{M}(\partial T)$ for \mathcal{H}^1 - a.e. $x \in M$. Thus,

$$\begin{aligned} \mathbf{M}(T) &= \int_M \theta(x) d\mathcal{H}^1(x) = \int_M \theta(x)^\alpha \theta(x)^{1-\alpha} d\mathcal{H}^1(x) \\ &\leq \int_M \theta(x)^\alpha (\mu^+(X))^{1-\alpha} d\mathcal{H}^1(x) = \left(\frac{\mathbf{M}(\partial T)}{2} \right)^{1-\alpha} \mathbf{M}_\alpha(T). \end{aligned}$$

\square

In the rest of the analysis, we assume that μ and ν are mutually singular, and h is continuous on the support of $\nu - \mu$.

Proposition 3.4. *If $\min\{h(x) : x \in \text{spt}(\mu)\} \geq \max\{h(x) : x \in \text{spt}(\nu)\}$, then $T^* = 0$ is the unique solution to the ROTPB(μ, ν) problem.*

Proof. Suppose T^* is a solution to the ROTPB(μ, ν) problem with $\partial T^* = \tilde{\nu} - \tilde{\mu}$. Since $\tilde{\mu}$ and $\tilde{\nu}$ have the same mass,

$$\begin{aligned} \mathbf{E}(T^*) &= \mathbf{M}_\alpha(T^*) - \int_X h d\tilde{\nu} + \int_X h d\tilde{\mu} \\ &\geq \mathbf{M}_\alpha(T^*) - \int_X \max\{h(x) : x \in \text{spt}(\nu)\} d\tilde{\nu} + \int_X \min\{h(x) : x \in \text{spt}(\mu)\} d\tilde{\mu} \\ &= \mathbf{M}_\alpha(T^*) + (\min\{h(x) : x \in \text{spt}(\mu)\} - \max\{h(x) : x \in \text{spt}(\nu)\}) \tilde{\mu}(X) \geq 0 \end{aligned}$$

where the equality holds if and only if $T^* = 0$. \square

The condition in the proposition implies that it is impossible to obtain positive net payoff from relocating mass, needless to mention the incurred transportation cost. It is thus in the best interest of the planner to not move any mass at all. This proposition illustrates the role of boundary payoff played in the problem, which we will further examine in Section 5.

Suppose that the ROTPB(μ, ν) problem has a solution $T^* \in \text{Path}(\mu^*, \nu^*)$. Then, T^* is inherently an α -optimal transport path in $\text{Path}(\mu^*, \nu^*)$ with finite \mathbf{M}_α cost. Thus, T^* itself exhibits some nice regularity properties (acyclic, uniform upper-bound on the degree of vertices, uniform lower-bound on the angles between edges at each vertex, boundary and interior regularity, etc) as stated in [28] for being \mathbf{M}_α optimal.

4. PROPERTIES OF THE OPTIMAL ALLOCATION MEASURES

This section is devoted to characterizing the optimal allocation measures μ^* and ν^* . Let

$$\mathcal{E}(\mu, \nu) := \min \left\{ d_\alpha(\tilde{\mu}, \tilde{\nu}) - \int_X h d\tilde{\nu} + \int_X h d\tilde{\mu} \mid \tilde{\mu} \leq \mu \text{ and } \tilde{\nu} \leq \nu \text{ with } \|\tilde{\mu}\| = \|\tilde{\nu}\| \right\}$$

denote the minimum value of the ROTPB(μ, ν) problem. We first observe some basic properties of \mathcal{E} .

Proposition 4.1. *Suppose $0 \leq \tilde{\mu} \leq \mu$ and $0 \leq \tilde{\nu} \leq \nu$. Then,*

$$(4.1) \quad 0 \geq \mathcal{E}(\tilde{\mu}, \tilde{\nu}) \geq \mathcal{E}(\mu, \nu).$$

In particular, if $\mathcal{E}(\mu, \nu) = 0$, then for all $(\tilde{\mu}, \tilde{\nu})$ with $0 \leq \tilde{\mu} \leq \mu$ and $0 \leq \tilde{\nu} \leq \nu$, it holds that $\mathcal{E}(\tilde{\mu}, \tilde{\nu}) = 0$.

Proof. The results follow from the definition of $\mathcal{E}(\mu, \nu)$. \square

Here, $\mathcal{E}(\mu, \nu)$ is non-positive and monotonic since $-\mathcal{E}(\mu, \nu)$ represents the overall possible profit generated for the planner from the pair (μ, ν) . When $\mathcal{E}(\mu, \nu) = 0$, there is no way to generate a non-zero $\mathcal{E}(\tilde{\mu}, \tilde{\nu})$ from some part $(\tilde{\mu}, \tilde{\nu})$ of (μ, ν) .

Proposition 4.2. *Suppose for each $i = 1, 2$, T_i^* is a solution to the ROTPB(μ_i, ν_i) problem, and T_{1+2}^* is a solution to the ROTPB($\mu_1 + \mu_2, \nu_1 + \nu_2$) problem, then*

$$(4.2) \quad \mathbf{E}(T_{1+2}^*) \leq \mathbf{E}(T_1^*) + \mathbf{E}(T_2^*).$$

This proposition implies that

$$(4.3) \quad \mathcal{E}(\mu_1 + \mu_2, \nu_1 + \nu_2) \leq \mathcal{E}(\mu_1, \nu_1) + \mathcal{E}(\mu_2, \nu_2).$$

Proof. By assumption, for each $i = 1, 2$, $T_i^* \in \text{Path}(\mu_i^*, \nu_i^*)$ with $\mu_i^* \leq \mu_i$ and $\nu_i^* \leq \nu_i$. Then, $T_1^* + T_2^* \in \text{Path}(\mu_1^* + \mu_2^*, \nu_1^* + \nu_2^*)$ with $\mu_1^* + \mu_2^* \leq \mu_1 + \mu_2$ and $\nu_1^* + \nu_2^* \leq \nu_1 + \nu_2$. Since T_{1+2}^* is a solution to the ROTPB($\mu_1 + \mu_2, \nu_1 + \nu_2$) problem, we have

$$\begin{aligned} \mathbf{E}(T_{1+2}^*) &\leq \mathbf{E}(T_1^* + T_2^*) = \mathbf{M}_\alpha(T_1^* + T_2^*) - \int_X hd(\partial T_1^* + \partial T_2^*) \\ &\leq \mathbf{M}_\alpha(T_1^*) + \mathbf{M}_\alpha(T_2^*) - \int_X hd(\partial T_1^*) - \int_X hd(\partial T_2^*) \\ &= \mathbf{E}(T_1^*) + \mathbf{E}(T_2^*). \end{aligned}$$

□

Following from the above proof, if the equality in (4.2) holds, then

$$\mathbf{M}_\alpha(T_1^* + T_2^*) = \mathbf{M}_\alpha(T_1^*) + \mathbf{M}_\alpha(T_2^*).$$

Suppose $T_i = \underline{\tau}(M_i, \theta_i, \xi_i)$ with $\theta_i(x) > 0$ for \mathcal{H}^1 -a.e. $x \in M_i$ with $i = 1, 2$. Since $\alpha < 1$,

$$\begin{aligned} &\mathbf{M}_\alpha(T_1^* + T_2^*) - \mathbf{M}_\alpha(T_1^*) - \mathbf{M}_\alpha(T_2^*) \\ &\leq \int_{M_1 \cap M_2} (\theta_1(x) + \theta_2(x))^\alpha - \theta_1(x)^\alpha - \theta_2(x)^\alpha d\mathcal{H}^1(x) \leq 0, \end{aligned}$$

where the equalities hold only if $\mathcal{H}^1(M_1 \cap M_2) = 0$.

We now give a necessary condition on the solution to the ROTPB(μ, ν) problem.

Corollary 4.3. *Suppose that $T^* \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem. Then $\mathcal{E}(\mu - \mu^*, \nu - \nu^*) = 0$.*

Proof. Since $T^* \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem, $\mathcal{E}(\mu, \nu) = \mathcal{E}(\mu^*, \nu^*)$. By (4.1) and (4.3),

$$0 \geq \mathcal{E}(\mu - \mu^*, \nu - \nu^*) \geq \mathcal{E}(\mu, \nu) - \mathcal{E}(\mu^*, \nu^*) = 0.$$

Therefore, $\mathcal{E}(\mu - \mu^*, \nu - \nu^*) = 0$. □

The lemma says that the mass left unmoved by the solution would not generate further gains for the planner.

Proposition 4.4. *Suppose that the ROTPB(μ, ν) problem has a non-zero solution $T^* \in \text{Path}(\mu^*, \nu^*)$ and $\alpha < 1$. Then there exists no real number $\sigma > 1$ such that $\sigma\mu^* \leq \mu$ and $\sigma\nu^* \leq \nu$.*

Proof. Otherwise, assume that there exists a real number $\sigma > 1$ such that $\sigma\mu^* \leq \mu$ and $\sigma\nu^* \leq \nu$. We consider the function

$$g(\lambda) := \mathbf{E}(\lambda\mu^*, \lambda\nu^*) = \lambda^\alpha d_\alpha(\mu^*, \nu^*) - \lambda \int_X |h| d\nu^* + \lambda \int_X |h| d\mu^*$$

for $\lambda \in [0, \sigma]$. Since $T^* \in \text{Path}(\mu^*, \nu^*)$ is a non-zero solution to the ROTPB(μ, ν) problem, $d_\alpha(\mu^*, \nu^*) = \mathbf{M}_\alpha(T^*) > 0$. Thus, given $\alpha < 1$,

$$\begin{aligned} g'(1) &= \alpha d_\alpha(\mu^*, \nu^*) - \int_X |h| d\nu^* + \int_X |h| d\mu^* \\ &< d_\alpha(\mu^*, \nu^*) - \int_X |h| d\nu^* + \int_X |h| d\mu^* \\ &= \mathbf{E}(T^*) \leq \mathbf{E}(0) = 0. \end{aligned}$$

As a result, there exists a $\lambda^* \in (1, \sigma)$ such that $g(\lambda^*) < g(1)$. Because $\sigma\mu^* \leq \mu$ and $\sigma\nu^* \leq \nu$, we also have $\lambda^*\mu^* \leq \sigma\mu^* \leq \mu$ and $\lambda^*\nu^* \leq \sigma\nu^* \leq \nu$. Hence $\mathbf{E}(\lambda^*\mu^*, \lambda^*\nu^*) = g(\lambda^*) < g(1) = \mathbf{E}(\mu^*, \nu^*)$, which contradicts with T^* being a solution to the ROTPB(μ, ν) problem. \square

At a solution to the ROTPB(μ, ν) problem, the planner might only move out a portion of the mass held at one source or ship in mass less than registered at a single destination. However, the above proposition shows that this can not happen at all the involved sources and destinations. Otherwise, an improvement can be achieved by a proportional increase of the transported mass at these locations. This is because the resulting marginal payoff from moving more mass outweighs the marginal transportation cost thanks to the transport economy of scale when $\alpha < 1$.

The remainder of this section focuses on characterizing the optimal allocation measures μ^* and ν^* , with the main result stated in Theorem 4.18. We first set up some technical bases.

Definition 4.5. Let $T = \underline{\tau}(M, \theta, \xi)$ and $S = \underline{\tau}(N, \rho, \eta)$ be two rectifiable 1-currents. We say S is on T if $\mathcal{H}^1(N \setminus M) = 0$, and $\rho(x) \leq \theta(x)$ for \mathcal{H}^1 almost all $x \in N$.

Note that when $S = \underline{\tau}(N, \rho, \eta)$ is on $T = \underline{\tau}(M, \theta, \xi)$, then $\xi(x) = \pm\eta(x)$ for \mathcal{H}^1 almost all $x \in N$, since two rectifiable sets have the same tangent a.e. on their intersection.

Theorem 4.6. *Suppose that $T^* \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem, and $0 < \alpha < 1$. If there exists a rectifiable 1-current S on T^* with*

$$\partial(T^* + S) \preceq \nu - \mu \text{ and } \partial(T^* - S) \preceq \nu - \mu,$$

then $S = 0$.

Proof. Assume that $S = \underline{\tau}(N, \rho, \eta)$ is a non-zero rectifiable 1-current on $T^* = \underline{\tau}(W, \theta, \xi)$. One may assume that $N = W$ by extending $\rho(x) = 0$ and $\eta(x) = \xi(x)$ for $x \in W \setminus N$. Since T^* is a solution to the ROTPB(μ, ν) problem and $\partial(T^* \pm S) \preceq \nu - \mu$, the function $g(t) := \mathbf{E}(T^* + tS)$ defined on the interval $[-1, 1]$ achieves its minimum value at $t = 0$. Nevertheless,

$$\begin{aligned} g(t) &= \mathbf{E}(T^* + tS) = \mathbf{M}_\alpha(T^* + tS) - \int_X hd(\partial(T^* + tS)) \\ &= \mathbf{E}(T^*) + \int_W |\theta(x) + t\rho(x)\langle \xi(x), \eta(x) \rangle|^\alpha - \theta(x)^\alpha d\mathcal{H}^1(x) - t \int_X hd(\partial S). \end{aligned}$$

Here, the value of the inner product $\langle \xi(x), \eta(x) \rangle = \pm 1$ for $\mathcal{H}^1 - a.e. x \in W$. Then,

$$g''(0) = \alpha(\alpha - 1) \int_W \theta(x)^{\alpha-2} \rho(x)^2 d\mathcal{H}^1(x) < 0,$$

since $0 < \alpha < 1$ and S is non-zero. This says that g can not achieve a local minimum at $t = 0$, a contradiction. \square

4.1. Finite atomic case. In the context of finite atomic measures, Theorem 4.6 has important implications for the structure of the optimal transport path T^* as demonstrated by the following results.

Proposition 4.7. *Suppose both*

$$\mu = \sum_{i=1}^{\ell} a_i \delta_{x_i} \text{ and } \nu = \sum_{j=1}^n b_j \delta_{y_j}$$

are two finite atomic measures on X , $0 < \alpha < 1$, and $T^* \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem. Also, let

$$P := \text{spt}(\mu - \mu^*) \cup \text{spt}(\nu - \nu^*)$$

denote the union of the supports of the measures $\mu - \mu^*$ and $\nu - \nu^*$. Then each connected component of the support of T^* contains at most one element of P .

Proof. Without loss of generality, we may assume that the support of T^* is connected, and we want to show that the set

$$(4.4) \quad \begin{aligned} P &= \{x_i : \mu^*(\{x_i\}) < \mu(\{x_i\})\} \cup \{y_j : \nu^*(\{y_j\}) < \nu(\{y_j\})\} \\ &= \{p \in \{x_1, \dots, x_\ell, y_1, \dots, y_n\} : (\nu - \nu^*)\{p\} + (\mu - \mu^*)\{p\} > 0\} \end{aligned}$$

contains at most one element. Assume that P has at least two distinct elements p_1 and p_2 . Also, we may assume that $(\mu - \mu^*)\{p_1\} > 0$ and $(\mu - \mu^*)\{p_2\} > 0$ (the proofs for the other cases are similar). Since T^* is acyclic (see [28, Propositions 2.1 and 2.2]), there exists a unique oriented curve γ on the support of T^* from p_1 to p_2 , and set $S = \sigma I_\gamma$ with

$$\sigma = \min(\{\theta(x) : x \in \gamma\}, (\mu - \mu^*)\{p_1\}, (\mu - \mu^*)\{p_2\}) > 0,$$

and I_γ being the rectifiable 1-current associated with γ (see (4.10) for the precise definition). Then, S is non-zero and on T in the sense of Definition 4.5. Moreover, by the choice of σ ,

$$\mu^* \pm \sigma(\delta_{p_2} - \delta_{p_1}) \leq \mu.$$

Thus,

$$\partial(T \pm S) = \nu^* - \mu^* \pm \sigma(\delta_{p_2} - \delta_{p_1}) \preceq \nu - \mu.$$

According to Theorem 4.6, S must be zero, a contradiction. \square

The set P in Proposition 4.7 represents the collection of boundary nodes on which the amount of mass involved in the optimal transport path T^* is smaller than its counterpart specified initially. The proof hinges on the fact that if a connected component of the support of T^* contains two elements in P , one would be able to cut cost by reallocating the mass transported along T^* , which however is precluded by Theorem 4.6.

According to Proposition 4.4, in the finite atomic case, there exists at least one point p on the support of μ^* or one point q on the support of ν^* , such that either

$$(4.5) \quad \mu^*(\{p\}) = \mu(\{p\}) \text{ or } \nu^*(\{q\}) = \nu(\{q\}).$$

Proposition 4.7 says that with at most one exception on each connected component, equation (4.5) holds for all points p or q on the supports of μ^* or ν^* , respectively. Consequently, with the help of Proposition 4.7 and Corollary 4.3, we may prove Theorem 1.1 as follows.

Proof of Theorem 1.1. By Proposition 4.7, each K_k contains at most one element of the set P . Thus, one of the following two cases holds:

Case 1:

$$\mu^* \llcorner K_k = \mu \llcorner K_k - m_k \delta_{p_k} \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k$$

for some point $p_k \in K_k \cap \text{spt}(\mu^*)$ and some real number $m_k \geq 0$.

Case 2:

$$\mu^* \llcorner K_k = \mu \llcorner K_k \text{ and } \nu^* \llcorner K_k = \nu \llcorner K_k - n_k \delta_{q_k}$$

for some point $q_k \in K_k \cap \text{spt}(\nu^*)$ and some real number $n_k \geq 0$.

In the first case,

$$\mu^*(K_k) = \mu(K_k) - m_k \text{ and } \nu^*(K_k) = \nu(K_k).$$

Since $\mu(K_k) \geq \mu^*(K_k) = \nu^*(K_k) = \nu(K_k)$, it follows that

$$m_k = \mu(K_k) - \mu^*(K_k) = \mu(K_k) - \nu(K_k) = \max\{\mu(K_k) - \nu(K_k), 0\}.$$

Analogously, in the second case, we pick

$$n_k = \max\{\nu(K_k) - \mu(K_k), 0\}$$

as desired. \square

If the measure of mass at each source node is sufficiently large, all source nodes would fall into the set P , yielding a natural partition of the transport path T^* as stated in the following corollary. In this case, destination nodes can be classified by the source node from which they receive the mass. Under a symmetric condition, a similar decomposition exists for destination nodes.

Corollary 4.8. *Suppose both*

$$\mu = \sum_{i=1}^{\ell} a_i \delta_{x_i} \text{ and } \nu = \sum_{j=1}^n b_j \delta_{y_j}$$

are (positive) finite atomic measures on X , and T^* is a solution to the ROTPB(μ, ν) problem.

(a) *If*

$$(4.6) \quad \min_{1 \leq i \leq \ell} a_i \geq \sum_{j=1}^n b_j,$$

then T^* can be decomposed as $T^* = T_1 + T_2 + \cdots + T_\ell$, where for each $i = 1, \dots, \ell$, T_i is an α -optimal transport path from a single source located at x_i .

(b) *Similarly, if*

$$(4.7) \quad \min_{1 \leq j \leq n} b_j \geq \sum_{i=1}^{\ell} a_i,$$

then T^* can be decomposed as $T^* = T_1 + T_2 + \cdots + T_n$, where for each $j = 1, \dots, n$, T_j is an α -optimal transport path to a single destination located at y_j .

Proof. We only need to prove case (a) as (b) follows from a symmetric argument. To do so, it is sufficient to show that each connected component of the support of T^* contains only one source point in $\{x_1, x_2, \dots, x_\ell\}$. We prove it by contradiction. Assume that there exists a connected component of the support of T^* that contains at least two sources, say x_1 and x_2 . Then

$$\mu^*({x_1}) > 0 \text{ and } \mu^*({x_2}) > 0.$$

As a result,

$$\mu^*({x_1}) < \mu^*({x_1}) + \mu^*({x_2}) \leq \|\mu^*\| = \|\nu^*\| \leq \sum_{j=1}^n b_j \leq a_1 = \mu({x_1}),$$

by (4.6). This shows that x_1 belongs to the set P in (4.4). Similar argument leads to $x_2 \in P$. This contradicts Proposition 4.7. Let $\{K_i : i = 1, 2, \dots, \ell\}$ be

the connected components of the support of T^* , and set $T_i = T \llcorner K_i$. Since T is α -optimal, and $\{K_i\}$ are pairwise disjoint, each T_i is also α -optimal. \square

4.2. General case. In what follows, we generalize the results of Theorem 1.1 for μ and ν being any two Radon measures, not necessarily finite atomic. To do so, we adopt a Lagrangian approach, and follow some notations used in [7].

By Theorem 3.1, the ROTPB(μ, ν) problem has a solution

$$(4.8) \quad T^* = \underline{\underline{T}}(W, \varphi, \zeta) \in \text{Path}(\mu^*, \nu^*).$$

We denote by Γ the space of 1-Lipschitz curves $\gamma : [0, \infty) \rightarrow \mathbb{R}^m$, which are eventually constant (and hence of finite length). For $\gamma \in \Gamma$, we denote the values

$$t_0(\gamma) := \sup\{t : \gamma \text{ is constant on } [0, t]\}$$

and

$$t_\infty(\gamma) := \inf\{t : \gamma \text{ is constant on } [t, \infty)\},$$

and denote $\gamma(\infty) := \lim_{t \rightarrow \infty} \gamma(t)$. Given $\gamma \in \Gamma$, the projections of γ onto its starting and stopping points are

$$(4.9) \quad p_0(\gamma) := \gamma(0) \text{ and } p_\infty(\gamma) := \gamma(\infty).$$

We say that a curve $\gamma \in \Gamma$ is *simple* if $\gamma(s) \neq \gamma(t)$ for every $t_0(\gamma) \leq s < t \leq t_\infty(\gamma)$. In particular, γ is non-constant in any non-trivial sub-interval $[s, t] \subseteq [t_0(\gamma), t_\infty(\gamma)]$.

For each simple curve $\gamma \in \Gamma$, we may canonically associate it with the rectifiable 1-current

$$(4.10) \quad I_\gamma := \underline{\underline{T}} \left(\text{Im}(\gamma), \frac{\gamma'}{|\gamma'|}, 1 \right),$$

where $\text{Im}(\gamma)$ denotes the image of the curve γ in \mathbb{R}^m . It is easy to check that $\mathbf{M}(I_\gamma) = \mathcal{H}^1(\text{Im}(\gamma))$ and $\partial I_\gamma = \delta_{\gamma(\infty)} - \delta_{\gamma(0)}$; since γ is simple, if it is also non-constant, then $\gamma(\infty) \neq \gamma(0)$ and $\mathbf{M}(\partial I_\gamma) = 2$.

A normal current $T \in \mathcal{D}_1(\mathbb{R}^m)$ is said *acyclic* if there exists no non-trivial current S such that $\partial S = 0$ and $\mathbf{M}(T) = \mathbf{M}(T - S) + \mathbf{M}(S)$.

Now, we recall a fundamental result of Smirnov in [21], which establishes that every acyclic normal 1-current can be written as a weighted average of simple Lipschitz curves in the following sense.

Definition 4.9. Let T be a normal 1-current in \mathbb{R}^m represented as a vector-valued measure $\vec{T} \llcorner |T|$, and let η be a finite positive measure on Γ such that

$$(4.11) \quad T = \int_\Gamma I_\gamma d\eta(\gamma)$$

in the sense that for every smooth compactly supported 1-form $\omega \in \mathcal{D}^1(\mathbb{R}^m)$, it holds that

$$(4.12) \quad T(\omega) = \int_\Gamma I_\gamma(\omega) d\eta(\gamma).$$

We say that η is a *good decomposition* of T if η is supported on non-constant, simple curves and satisfies the following equalities:

- (a) $\mathbf{M}(T) = \int_\Gamma \mathbf{M}(I_\gamma) d\eta(\gamma) = \int_\Gamma \mathcal{H}^1(\text{Im}(\gamma)) d\eta(\gamma)$;
- (b) $\mathbf{M}(\partial T) = \int_\Gamma \mathbf{M}(\partial I_\gamma) d\eta(\gamma) = 2\eta(\Gamma)$.

It has been shown in [15, Theorem 10.1] that optimal transport paths T^* with finite \mathbf{M}_α cost are acyclic, and hence they admit such a good decomposition.

In the next result, we collect some useful properties of good decompositions, whose proof can be found in [5, Proposition 3.6].

Theorem 4.10. *(Existence and properties of good decompositions)[16, Theorem 5.1] and [5, Proposition 3.6]. Let T be an α -optimal transport path from μ^- to μ^+ with finite \mathbf{M}_α cost. Then T is acyclic and there is a Borel finite measure η on Γ such that η is a good decomposition of T . Moreover, if η is a good decomposition of T , the following statements hold:*

- $\mu^- = \int_\Gamma \delta_{\gamma(0)} d\eta(\gamma), \mu^+ = \int_\Gamma \delta_{\gamma(\infty)} d\eta(\gamma).$
- If $T = \underline{\underline{\tau}}(M, \theta, \xi)$ is rectifiable, then

$$(4.13) \quad \theta(x) = \eta(\{\gamma \in \Gamma : x \in \text{Im}(\gamma)\})$$

for \mathcal{H}^1 -a.e. $x \in M$.

- For every $\tilde{\eta} \leq \eta$, the representation

$$\tilde{T} = \int_\Gamma I_\gamma d\tilde{\eta}(\gamma)$$

is a good decomposition of \tilde{T} . Moreover, if $T = \underline{\underline{\tau}}(M, \theta, \xi)$ is rectifiable, then \tilde{T} can be written as $\tilde{T} = \underline{\underline{\tau}}(M, \tilde{\theta}, \xi)$ with $\tilde{\theta}(x) \leq \min\{\theta(x), \tilde{\eta}(\Gamma)\}$ for \mathcal{H}^1 -a.e. $x \in M$.

We now introduce the following notations. Let $T^* = \underline{\underline{\tau}}(W, \varphi, \zeta) \in \text{Path}(\mu^*, \nu^*)$ be given as in (4.8), and η be a good decomposition of T^* . Denote

$$\tilde{\mu} = \mu - \mu^* \text{ and } \tilde{\nu} = \nu - \nu^*.$$

For any $x \in W$, let us denote by $\Gamma(x)$ the set of simple curves $\gamma \in \Gamma$ such that $x \in \text{Im}(\gamma)$. By equation (4.13), $\varphi(x) = \eta(\Gamma(x))$ for \mathcal{H}^1 -a.e. $x \in W$.

Proposition 4.11. *For any $x \in W$ with $\varphi(x) = \eta(\Gamma(x)) > 0$, denote*

$$\bar{\mu}_x = (p_0)_\# (\eta \llcorner \Gamma(x)) \text{ and } \bar{\nu}_x = (p_\infty)_\# (\eta \llcorner \Gamma(x))$$

where p_0 and p_∞ are projections given in (4.9). Let

$$\tilde{\mu} = \tilde{\mu}_x^{ac} + \tilde{\mu}_x^s, \tilde{\mu}_x^{ac} \ll \bar{\mu}_x, \tilde{\mu}_x^s \perp \bar{\mu}_x$$

be the Lebesgue-Radon-Nikodým decomposition of $\tilde{\mu}$ with respect to $\bar{\mu}_x$, and

$$\tilde{\nu} = \tilde{\nu}_x^{ac} + \tilde{\nu}_x^s, \tilde{\nu}_x^{ac} \ll \bar{\nu}_x, \tilde{\nu}_x^s \perp \bar{\nu}_x$$

be the Lebesgue-Radon-Nikodým decomposition of $\tilde{\nu}$ with respect to $\bar{\nu}_x$. Then

- (a) There exist $p_x \in W$, $q_x \in W$, $m_x \geq 0$, and $n_x \geq 0$ such that

$$\tilde{\mu}_x^{ac} = m_x \delta_{p_x} \text{ and } \tilde{\nu}_x^{ac} = n_x \delta_{q_x}.$$

- (b) If $m_x > 0$, then $\bar{\mu}_x(\{p_x\}) > 0$ and there exists a Lipschitz curve γ_x^- from p_x to x such that $\varphi(y) \geq \bar{\mu}_x(p_x)$ for \mathcal{H}^1 -a.e. $y \in \text{Im}(\gamma_x^-)$.
- (c) If $n_x > 0$, then $\bar{\nu}_x(\{q_x\}) > 0$ and there exists a Lipschitz curve γ_x^+ from x to q_x such that $\varphi(y) \geq \bar{\nu}_x(q_x)$ for \mathcal{H}^1 -a.e. $y \in \text{Im}(\gamma_x^+)$.
- (d) At least one of m_x and n_x is zero.

Proof. Note that

$$\bar{\mu}_x(X) = (p_0)_\#(\eta \llcorner \Gamma(x))(X) = (\eta \llcorner \Gamma(x))(p_0^{-1}(X)) = \eta(\Gamma(x)) = \varphi(x) > 0,$$

and $\bar{\mu}_x \leq (p_0)_\# \eta = \mu^*$. For the sake of contradiction, we assume that $\tilde{\mu}_x^{ac}$ is non-zero and not a multiple of a Dirac mass. Then there exists a Borel measurable set R such that

$$\tilde{\mu}_x^{ac}(R) > 0 \text{ and } \tilde{\mu}_x^{ac}(X \setminus R) > 0.$$

Since $\tilde{\mu}_x^{ac} \ll \bar{\mu}_x$, there exists a non-negative function $g \in \mathcal{L}^1(X, \bar{\mu}_x)$ such that $\tilde{\mu}_x^{ac} = g \cdot \bar{\mu}_x$. We define

$$\mu_0 := \min\{g(\cdot), 1\} \cdot \bar{\mu}_x.$$

Still we have $\mu_0(R) > 0$ and $\mu_0(X \setminus R) > 0$. Without loss of generality, we may assume that $0 < \mu_0(R) \leq \mu_0(X \setminus R)$. Setting

$$\mu_1 = \mu_0 \llcorner R \text{ and } \mu_2 = \frac{\mu_0(R)}{\mu_0(X \setminus R)} (\mu_0 \llcorner (X \setminus R))$$

yields two positive measures μ_1 and μ_2 such that μ_1 is concentrated on R and μ_2 is concentrated on $X \setminus R$ with equal mass. Moreover, since both μ_1 and μ_2 are absolutely continuous with respect to $\bar{\mu}_x$, there exist two non-negative $\bar{\mu}_x$ -integrable functions ρ_1 and ρ_2 such that

$$\mu_1 = \rho_1 \bar{\mu}_x \text{ and } \mu_2 = \rho_2 \bar{\mu}_x.$$

Let $\rho = \rho_1 - \rho_2$. Note that $|\rho| \leq 1$. From the construction of μ_1 and μ_2 , we have

$$\bar{\mu}_x(\rho) = \int_X \rho d\bar{\mu}_x = \int_X \rho_1 d\bar{\mu}_x - \int_X \rho_2 d\bar{\mu}_x = 0,$$

and

$$\int_X |\rho| d\bar{\mu}_x = 2 \int_X \rho_1 d\bar{\mu}_x = 2\tilde{\mu}_x^{ac}(R) > 0.$$

Now, for any $\gamma \in \Gamma(x)$, let γ^- be the part of γ from $\gamma(0)$ to x . Define

$$S := \int_{\Gamma(x)} \rho(\gamma(0)) I_{\gamma^-} d\eta(\gamma).$$

Then for any smooth function f with a compact support, we have

$$\begin{aligned} \partial S(f) &= S(df) \\ &= \int_{\Gamma(x)} \rho(\gamma(0)) I_{\gamma^-} (df) d\eta(\gamma) \\ &= \int_{\Gamma(x)} \rho(\gamma(0)) \partial(I_{\gamma^-})(f) d\eta(\gamma) \\ &= \int_{\Gamma(x)} \rho(\gamma(0)) (f(x) - f(\gamma(0))) d\eta(\gamma) \\ &= \int_{\Gamma(x)} \rho(\gamma(0)) f(x) d\eta(\gamma) - \int_{\Gamma(x)} \rho(\gamma(0)) f(\gamma(0)) d\eta(\gamma) \\ &= f(x) \bar{\mu}_x(\rho) - (\rho \bar{\mu}_x)(f) = -(\rho \bar{\mu}_x)(f). \end{aligned}$$

Therefore, $\partial S = -\rho \bar{\mu}_x \neq 0$.

Since T^* is rectifiable, by construction S is also rectifiable. We may write it as $S = \underline{\llcorner}(M_S, \theta_S, \xi_S)$ for some $M_S \subseteq W$. At \mathcal{H}^1 -a.e. $y \in M_S$,

$$\theta_S(y) \leq \int_{\Gamma(x) \cap \Gamma(y)} |\rho(\gamma(0))| d\eta(\gamma) \leq \int_{\Gamma(x) \cap \Gamma(y)} 1 d\eta(\gamma) \leq \int_{\Gamma(y)} 1 d\eta(\gamma) = \varphi(y).$$

This shows that S is on T^* in the sense of Definition 4.5.

We now show that $\partial(T^* \pm S) \leq \nu - \mu$. Given

$$\partial(T^* \pm S) = \nu^* - \mu^* \pm (\rho \bar{\mu}_x) = \nu^* - (\mu^* \mp (\rho \bar{\mu}_x)),$$

it is sufficient to show that $\mu^* \mp (\rho \bar{\mu}_x) \leq \mu$ as (positive) measures, which is the case provided that $\bar{\mu}_x \leq \mu^*$ and $|\rho| \leq 1$. Also,

$$\mu - (\mu^* \mp (\rho \bar{\mu}_x)) = \mu - \mu^* \pm (\rho \bar{\mu}_x) = \tilde{\mu} \pm (\mu_1 - \mu_2)$$

are positive measures because $\mu_1 \leq \bar{\mu}_x^{ac} \leq \tilde{\mu}$ and similarly $\mu_2 \leq \tilde{\mu}$. As a result, $\partial(T^* \pm S) \leq \nu - \mu$. By Theorem 4.6, S is zero which contradicts $\partial S \neq 0$. Therefore, $\bar{\mu}_x^{ac}$ must be in the form of $m_x \delta_{p_x}$ for some $m_x \geq 0$ and $p_x \in W$. Similarly, we have $\bar{\nu}_x^{ac} = n_x \delta_{q_x}$ for some $n_x \geq 0$ and $q_x \in W$. This proves part (a).

Now assume that $m_x > 0$. Since $\bar{\mu}_x^{ac} = m_x \delta_{p_x} \ll \bar{\mu}_x$, we have $\bar{\mu}_x(\{p_x\}) > 0$. That is,

$$0 < (p_0)_\# (\eta|_{\Gamma(x)}) (\{p_x\}) = \eta(\{\gamma \in \Gamma(x) : \gamma(0) = p_x\}).$$

Because T^* is acyclic and η is a good decomposition of T^* , for η -a.e. $\gamma \in \Gamma(x)$ with $\gamma(0) = p_x$, the image $Im(\gamma)$ of γ shares a common Lipschitz curve γ_x^- in W from p_x to x . For \mathcal{H}^1 -a.e. y on $Im(\gamma_x^-)$,

$$\varphi(y) = \eta(\{\gamma \in \Gamma(y)\}) \geq \eta(\{\gamma \in \Gamma(x) : \gamma(0) = p_x\}) = \bar{\mu}_x(\{p_x\}).$$

This proves part (b). Similar arguments lead to part (c).

Suppose by contradiction that both $m_x > 0$ and $n_x > 0$. Then, by parts (b) and (c), we consider the rectifiable 1-current

$$S_x := \sigma \left(I_{\gamma_x^-} + I_{\gamma_x^+} \right),$$

for $\sigma = \min\{\bar{\mu}_x(\{p_x\}), \bar{\nu}_x(\{q_x\}), m_x, n_x\} > 0$. Clearly, S_x is on T^* , non-zero, and $\partial(T^* \pm S_x) \leq \nu - \mu$. This contradicts Theorem 4.6. Therefore, at least one of m_x and n_x is zero. \square

To derive the generalized version of Theorem 1.1, we introduce the concept of path-connectivity on rectifiable 1-currents as follows.

Definition 4.12. Let $T = \underline{T}(M, \theta, \xi)$ be a rectifiable 1-current. For any two points $x_1, x_2 \in X$, we say x_1 and x_2 are T -path-connected if there exists a Lipschitz curve $\gamma : [0, 1] \rightarrow X$ such that $\gamma(0) = x_1, \gamma(1) = x_2, \mathcal{H}^1(Im(\gamma) \setminus M) = 0$, and there exists a number $c > 0$ such that $\theta(z) \geq c$ for \mathcal{H}^1 -a.e. $z \in Im(\gamma) \cap M$.

The T -path-connectivity defines an equivalence relation on X . A rectifiable 1-current $T = \underline{T}(M, \theta, \xi)$ is called path-connected if every two points on M are T -path-connected.

For any T -path-connected component M' of M , we consider the restriction $T \llcorner M'$ of T on M' . $T \llcorner M'$ is zero if M' contains only one point. In this case, we say that the component M' is degenerate. When M' is non-degenerate, i.e., it contains at least two distinct points x_1 and x_2 , we have

$$\mathbf{M}(T \llcorner M') = \int_{M'} \theta d\mathcal{H}^1 \geq c \mathcal{H}^1(Im(\gamma)) > 0$$

using the notations given in Definition 4.12. Since $\mathbf{M}(T) < \infty$, M has at most countably many non-degenerate T -path-connected components.

Observe that non-degenerate components may fail to exist even if $\mathbf{M}(T) > 0$. For instance, let C (e.g., a fat-Cantor set) be a nowhere dense subset of $[0, 1]$ with $0 < \mathcal{H}^1(C) < 1$. Then, for $S = \underline{T}(C, \chi_C, 1)$, each S -path-connected component is

degenerate. Luckily, the following lemma indicates that each non-zero α -optimal transport path has at least one non-degenerate path-connected component.

Lemma 4.13. *Let $T = \underline{\underline{T}}(M, \theta, \xi)$ be a non-zero α -optimal transport path for some $0 < \alpha < 1$. Then*

$$T = \sum_{i \in J} T \llcorner M_i,$$

where $\{M_i : i \in J\}$ are the collection of all non-degenerate T -path-connected components of M , and J is a non-empty countable set.

To prove Lemma 4.13, we first recall the notation of superlevel set as introduced in [27]: For any $\lambda > 0$, the λ -superlevel set of a rectifiable current $T = \underline{\underline{T}}(M, \theta, \xi)$ is the set

$$M_\lambda := \{p \in M : \theta(p) \geq \lambda\}.$$

Lemma 4.14. ([27, Proposition 4.3]) *Let $T = \underline{\underline{T}}(M, \theta, \xi)$ be any α -optimal transport path. Then for any $\sigma_1 > \sigma_2 > 0$ and any $\bar{p} \in M_{\sigma_1}$, there exists an open ball neighborhood $B_r(p)$ of p such that*

$$(4.14) \quad M_{\sigma_1} \cap B_r(p) \subseteq \text{the support of } Q_p \subseteq M_{\sigma_2} \cap B_r(p),$$

where $Q_p = \sum_{i=1}^K m_i \Gamma_i$ is a bi-Lipschitz chain.

Here, as stated in [27, Corollary 4.2], each Γ_i is a bi-Lipschitz curve from p . These bi-Lipschitz curves Γ_i are pairwise disjoint except at their common endpoint p , and K is a universal constant.

Proof of Lemma 4.13: Let $\{M_i : i \in J\}$ be the collection of all non-degenerate T -path-connected components of M , where J is countable. For any $p \in M$ with $\theta(p) > 0$, let $\sigma_1 = \theta_p$ and $\sigma_2 = \frac{1}{2}\sigma_1$. By (4.14), any point on the support of the bi-Lipschitz curve Q_p is T -path-connected with p . Hence, p belongs to some non-degenerate T -path-connected component M_i for some $i \in J$. As a result, we decompose $M_+ = \{x \in M : \theta(x) > 0\}$ as the disjoint union of M_i with $i \in J$. Thus, $T = \sum_{i \in J} T \llcorner M_i$. \square

We now go back to the study of T^* , which is also an α -optimal transport path. Consequently, one can write

$$T^* = \sum_{i \in J} T^* \llcorner W_i,$$

where $\{W_i : i \in J\}$ are the collection of all non-degenerate T^* -path-connected components of W , and J is a non-empty countable set if T^* is non-zero.

Lemma 4.15. *Suppose x_1 and x_2 belong to the same non-degenerate T^* -path-connected component of W . Then, at most one of $\{m_{x_1}, n_{x_1}, m_{x_2}, n_{x_2}\}$ is non-zero.*

Proof. Otherwise, let us just assume $m_{x_1} > 0$ and $n_{x_2} > 0$, with the proof for other cases following similarly. Since x_1 and x_2 are path-connected on T^* , there exists a Lipschitz curve $\gamma_{x_1}^{x_2}$ from x_1 to x_2 such that $\mathcal{H}^1(\text{Im}(\gamma_{x_1}^{x_2}) \setminus W) = 0$, and there exists a number $c > 0$ such that $\varphi(z) \geq c$ for \mathcal{H}^1 -a.e. $z \in \text{Im}(\gamma_{x_1}^{x_2})$. Now, we consider the rectifiable 1-current

$$S := \sigma \left(I_{\gamma_{x_1}^-} + I_{\gamma_{x_1}^{x_2}} + I_{\gamma_{x_2}^+} \right),$$

for $\sigma = \min\{\bar{\mu}_{x_1}(\{p_{x_1}\}), \bar{\nu}_{x_2}(\{q_{x_2}\}), m_{x_1}, n_{x_2}, c\} > 0$. Clearly, S is on T^* , non-zero, and $\partial(T^* \pm S_x) \leq \nu - \mu$. This contradicts Theorem 4.6. \square

For each $i \in J$, if there exists one point $x \in W_i$ such that $m_x > 0$, then by part(b) of Proposition 4.11, the associated point p_x is also T^* -path-connected with x and hence $p_x \in W_i$. By Lemma 4.15, $m_x \delta_{p_x}$ is independent of the choice of $x \in W_i$, and thus can be represented by $m_i \delta_{p_i}$. If $m_x = 0$ for all $x \in W_i$, we simply pick p_i to be any fixed point in W_i and set $m_i = 0$. Analogously, we denote $n_x \delta_{q_x}$ by $n_i \delta_{q_i}$ for each $i \in J$. As a result, we arrive at two atomic measures

$$(4.15) \quad \mathbf{a} = \sum_{i \in J} m_i \delta_{p_i} \text{ and } \mathbf{b} = \sum_{i \in J} n_i \delta_{q_i},$$

where either $m_i = 0$ or $n_i = 0$ for each $i \in J$.

Lemma 4.16. *It holds that*

$$(\tilde{\mu} - \mathbf{a}) \perp \mu^*, \mathbf{a} \ll \mu^* \text{ and } (\tilde{\nu} - \mathbf{b}) \perp \nu^*, \mathbf{b} \ll \nu^*.$$

Proof. Let $\hat{\mu} = \tilde{\mu} - \mathbf{a}$. Then for each $x \in W$ with $\varphi(x) > 0$, by Proposition 4.11,

$$\hat{\mu} \perp \bar{\mu}_x.$$

Thus, there exists a $\hat{\mu}$ -negligible set A_x such that $\bar{\mu}_x(A_x) = \bar{\mu}_x(X)$. Observe that one may pick countably many points $\{x_k : \varphi(x_k) > 0\}_{k=1}^{\infty} \subset W$ so that for η -a.e. $\gamma \in \Gamma$, γ passes at least one of these points. One way to select these points is taking a countable dense subset of the 1-rectifiable set $W_+ := \{x \in W : \varphi(x) > 0\}$. Now for each k ,

$$\eta(\Gamma(x_k)) = \bar{\mu}_{x_k}(X) = \bar{\mu}_{x_k}(A_{x_k}) = \eta(\{\gamma : \gamma(0) \in A_{x_k}\}),$$

and thus

$$\eta(\Gamma) = \eta\left(\bigcup_k \Gamma(x_k)\right) = \eta\left(\{\gamma \in \Gamma : \gamma(0) \in \bigcup_k A_{x_k}\}\right).$$

As a result,

$$\mu^*(X) = \mu^*\left(\bigcup_k A_{x_k}\right) \text{ and } \hat{\mu}\left(\bigcup_k A_{x_k}\right) = \sum_k \hat{\mu}(A_{x_k}) = 0.$$

Therefore, $\mu^* \perp \hat{\mu}$ as desired. For each $i \in J$, assume $m_i \delta_{p_i} = m_x \delta_{p_x}$, then

$$m_i \delta_{p_i} \ll \bar{\mu}_x \ll \mu^*.$$

Thus, $\mathbf{a} \ll \mu^*$. Similarly, we have $(\tilde{\nu} - \mathbf{b}) \perp \nu^*$ and $\mathbf{b} \ll \nu^*$. \square

Lemma 4.17. *For any two (positive) measures μ_1 and μ_2 . Let $\lambda = \mu_1 + \mu_2$. If $\mu_1 \perp \mu_2$, then there exists a λ -measurable set A such that $\mu_1 = \lambda \llcorner A$ and $\mu_2 = \lambda \llcorner (X \setminus A)$.*

Proof. Since $\lambda = \mu_1 + \mu_2$, $\mu_1 \ll \lambda$ and $\mu_2 \ll \lambda$. By the Radon-Nikodým theorem, there exists a non-negative λ -measurable function f such that $\mu_1 = f\lambda$ and $\mu_2 = (1-f)\lambda$. Given μ_2 is a positive measure, it follows that $0 \leq f(x) \leq 1$ for λ -a.e. x . Let $K = \{x : 0 < f(x) < 1\}$. We claim that $\lambda(K) = 0$. Indeed, assume $\lambda(K) > 0$. Then, $\mu_1(K) = \int_K f(x) d\lambda(x) > 0$ and similarly $\mu_2(K) = \int_K (1-f(x)) d\lambda(x) > 0$. This contradicts $\mu_1 \perp \mu_2$. Therefore, $\lambda(K) = 0$. Setting $A = \{x : f(x) = 1\}$ yields $\mu_1 = \lambda \llcorner A$ and $\mu_2 = \lambda \llcorner (X \setminus A)$. \square

Combining the preceding results leads to the following theorem, which is a generalized version of Theorem 1.1.

Theorem 4.18. *Suppose μ and ν are two Radon measures on X , $0 < \alpha < 1$, and $T^* = \underline{T}(W, \varphi, \zeta) \in \text{Path}(\mu^*, \nu^*)$ is a solution to the ROTPB(μ, ν) problem. Then*

- (1) There exist two atomic measures \mathbf{a} and \mathbf{b} , a μ -measurable set A , and a ν -measurable set B such that

$$\mu \llcorner A = \mu^* + \mathbf{a} \text{ and } \nu \llcorner B = \nu^* + \mathbf{b}.$$

- (2) Let $\{W_i : i \in J\}$ be the collection of all non-degenerate T^* -path-connected components of W . Then, for each $i \in J$, there exist $m_i \geq 0$ and $n_i \geq 0$ with either $m_i = 0$ or $n_i = 0$; and two points $p_i, q_i \in W_i$ such that

$$\mathbf{a} = \sum_{i \in J} m_i \delta_{p_i} \text{ and } \mathbf{b} = \sum_{i \in J} n_i \delta_{q_i}.$$

The theorem says that on each path-connected component, at most one atom is not fully in use. In particular, when both μ and ν are atom-free² measures, it follows that

$$\mu^* = \mu \llcorner A \text{ and } \nu^* = \nu \llcorner B.$$

Proof. The atomic measures \mathbf{a} and \mathbf{b} are obtained by (4.15). By Lemma 4.16, $(\mu - \mathbf{a} - \mu^*) \perp \mu^*$. By Lemma 4.17, there exists a $(\mu - \mathbf{a})$ -measurable set A such that $\mu^* = (\mu - \mathbf{a}) \llcorner A$. Since μ^* concentrates on A and $\mathbf{a} \ll \mu^*$, we have $\mathbf{a} \llcorner A = \mathbf{a}$ and A is also μ -measurable. Thus, $\mu \llcorner A = \mu^* + \mathbf{a}$. Similarly, we have $\nu \llcorner B = \nu^* + \mathbf{b}$ for some ν -measurable set B . \square

In light of the theorem, on locations involving mass transportation, the measure \mathbf{a} , which represents the mass left unmoved by the solution T^* , must be atomic, so is measure \mathbf{b} which summarizes the distribution of excess demand at destinations. This is because if not the planner can exploit further gains by relocating the mass moved along the path T^* due to the efficiency in group transportation.

5. THE IMPACT OF BOUNDARY PAYOFF

An important deviation of the ROTPB problem from the literature is the dependence of its solution on the boundary payoff as exemplified by Proposition 3.4. To gain further insights, in what follows we examine the implications of the payoff function h for the problem. For the sake of expositional tractability, we assume that μ and ν are disjointly supported (i.e., $spt(\mu) \cap spt(\nu) = \emptyset$) and the function h takes the form

$$(5.1) \quad h(x) = \begin{cases} c_\mu, & \text{if } x \in spt(\mu) \\ c_\nu, & \text{if } x \in spt(\nu) \end{cases}$$

where c_μ and c_ν are constants. In this case, for any $T \in Path(\tilde{\mu}, \tilde{\nu})$,

$$\mathbf{E}_\alpha^h(T) = \mathbf{M}_\alpha(T) - \int_X c_\nu d\tilde{\nu} + \int_X c_\mu d\tilde{\mu} = \mathbf{M}_\alpha(T) - 2c\|\tilde{\mu}\| = \mathbf{M}_\alpha(T) - c\mathbf{M}(\partial T),$$

where $c = \frac{c_\nu - c_\mu}{2}$. The corresponding ROTPB(μ, ν) problem in this case becomes: Minimize

$$(5.2) \quad \mathbf{E}_\alpha^c(T) := \mathbf{M}_\alpha(T) - c\mathbf{M}(\partial T)$$

among all transport paths T with $\partial T \preceq \nu - \mu$. Without loss of generality, we may assume that $c_\nu = 2c$ and $c_\mu = 0$ in equation (5.1).

For each c , by Theorem 3.1, the ROTPB(μ, ν) problem has a solution T_c^* that minimizes \mathbf{E}_α^c . When $c \leq 0$, by Proposition 3.4, the problem has a unique solution $T_c^* = 0$. Thus, in the following context, we only need to investigate T_c^* for $c > 0$.

²A measure μ is called atom-free if $\mu(\{p\}) = 0$ for every $p \in X$.

Proposition 5.1. *Suppose μ and ν are two disjointly supported measures on X of equal mass, and $T_c^* \in \text{Path}(\mu_c^*, \nu_c^*)$ is a solution to the ROTPB(μ, ν) problem associated with $c > 0$. Then, for any transport path $T \in \text{Path}(\mu, \nu)$,*

$$\mathbf{M}_\alpha(T) - \mathbf{M}_\alpha(T_c^*) \geq c(\mathbf{M}(\partial T) - \mathbf{M}(\partial T_c^*)) \geq 0,$$

and hence $\mathbf{M}_\alpha(T_c^*) \leq d_\alpha(\mu, \nu)$. Moreover, define

$$(5.3) \quad s(\mu, \nu) := \inf \{ \|\mu - \tilde{\mu}\| + \|\nu - \tilde{\nu}\| : \tilde{\mu} \leq \mu, \tilde{\nu} \leq \nu, \tilde{\nu} - \tilde{\mu} \neq \nu - \mu \}.$$

If $s(\mu, \nu) > 0$ and $c > \frac{d_\alpha(\mu, \nu)}{s(\mu, \nu)}$, then T_c^* is an optimal transport path in $\text{Path}(\mu, \nu)$.

Proof. Indeed, for any transport path $T \in \text{Path}(\mu, \nu)$,

$$\begin{aligned} \mathbf{M}_\alpha(T) - \mathbf{M}_\alpha(T_c^*) &= (\mathbf{E}_\alpha^c(T) + c\mathbf{M}(\partial T)) - (\mathbf{E}_\alpha^c(T_c^*) + c\mathbf{M}(\partial T_c^*)) \\ &= (\mathbf{E}_\alpha^c(T) - \mathbf{E}_\alpha^c(T_c^*)) + c(\mathbf{M}(\partial T) - \mathbf{M}(\partial T_c^*)) \\ &\geq c(\mathbf{M}(\partial T) - \mathbf{M}(\partial T_c^*)) \geq 0. \end{aligned}$$

Also, when $s(\mu, \nu) > 0$ and $c > \frac{d_\alpha(\mu, \nu)}{s(\mu, \nu)}$, assume that T_c^* is not an optimal transport path in $\text{Path}(\mu, \nu)$. Since T_c^* is a solution to the ROTPB(μ, ν) problem, it is an optimal transport path in $\text{Path}(\mu_c^*, \nu_c^*)$. Thus, $\mu_c^* \neq \mu$ and $\nu_c^* \neq \nu$. Now, for any optimal transport path T in $\text{Path}(\mu, \nu)$, it follows that

$$\begin{aligned} d_\alpha(\mu, \nu) &\geq \mathbf{M}_\alpha(T) - \mathbf{M}_\alpha(T_c^*) \geq c(\mathbf{M}(\partial T) - \mathbf{M}(\partial T_c^*)) \\ &= c(\|\mu - \mu_c^*\| + \|\nu - \nu_c^*\|) \geq cs(\mu, \nu), \end{aligned}$$

a contradiction with the choice of c . \square

Proposition 5.1 shows that the transportation cost $\mathbf{M}_\alpha(T_c^*)$ associated with the solution T_c^* is bounded from above. More interestingly, when the parameter c , a measure of the profitability for relocating mass, is sufficiently large, T_c^* represents an optimal way of transporting mass from μ to ν . The intuition is that since the transportation cost is bounded, a large enough c would induce the planner to move as much mass as possible. This argument can be further validated by the following proposition, which derives an upper bound as well as the decay rate for the amount of mass left unmoved by T_c^* .

Proposition 5.2. *Suppose $\|\mu\| = \|\nu\|$, $c > 0$, $1 - \frac{1}{m} < \alpha < 1$ and $T_c^* \in \text{Path}(\mu_c^*, \nu_c^*)$ denotes the solution to the ROTPB(μ, ν) problem. Then*

$$(5.4) \quad \|\mu - \mu_c^*\| = \|\nu - \nu_c^*\| \leq \left(\frac{C_{m,\alpha} \text{diam}(X)}{2c} \right)^{\frac{1}{1-\alpha}},$$

where $C_{m,\alpha}$ is the constant given in (2.3).

Proof. Let $\tilde{T} \in \text{Path}(\mu - \mu_c^*, \nu - \nu_c^*)$ be an α -optimal transport path, and denote $T = T_c^* + \tilde{T} \in \text{Path}(\mu, \nu)$. By (2.2),

$$\begin{aligned} 0 &\leq \mathbf{E}_\alpha^c(T) - \mathbf{E}_\alpha^c(T_c^*) \\ &= (\mathbf{M}_\alpha(T) - c\mathbf{M}(\partial T)) - (\mathbf{M}_\alpha(T_c^*) - c\mathbf{M}(\partial T_c^*)) \\ &= (\mathbf{M}_\alpha(T) - \mathbf{M}_\alpha(T_c^*)) - c(\mathbf{M}(\partial T) - \mathbf{M}(\partial T_c^*)) \\ &\leq \mathbf{M}_\alpha(\tilde{T}) - c(\|\mu - \mu_c^*\| + \|\nu - \nu_c^*\|) \\ &\leq C_{m,\alpha} \text{diam}(X) \|\mu - \mu_c^*\|^\alpha - 2c\|\mu - \mu_c^*\|, \end{aligned}$$

which leads to inequality (5.4). \square

The next proposition characterizes the monotonicity properties of the solution. Intuitively, as c rises, the planner tends to move more mass between sources and destinations, resulting in larger transportation costs.

Proposition 5.3. *Suppose $\|\mu\| = \|\nu\|$, $c > 0$, $1 - \frac{1}{m} < \alpha < 1$ and $T_c^* \in \text{Path}(\mu_c^*, \nu_c^*)$ denotes the solution to the ROTPB(μ, ν) problem. Then, as a function of $c \in \mathbb{R}$,*

- (1) $\mathbf{E}_\alpha^c(T_c^*)$ is decreasing;
- (2) $\mathbf{M}_\alpha(T_c^*)$ is increasing with $\lim_{c \rightarrow \infty} \mathbf{M}_\alpha(T_c^*) = d_\alpha(\mu, \nu)$;
- (3) $\mathbf{M}(\partial T_c^*)$ is increasing with $\lim_{c \rightarrow \infty} \partial T_c^* = \nu - \mu$.

Proof. Indeed, for any $c_1 < c_2$,

$$\mathbf{E}_\alpha^{c_1}(T_{c_1}^*) = \mathbf{M}_\alpha(T_{c_1}^*) - c_1 \mathbf{M}(\partial T_{c_1}^*) \geq \mathbf{M}_\alpha(T_{c_1}^*) - c_2 \mathbf{M}(\partial T_{c_1}^*) = \mathbf{E}_\alpha^{c_2}(T_{c_1}^*) \geq \mathbf{E}_\alpha^{c_2}(T_{c_2}^*).$$

Also, the inequalities $\mathbf{E}_\alpha^{c_1}(T_{c_1}^*) \leq \mathbf{E}_\alpha^{c_1}(T_{c_2}^*)$ and $\mathbf{E}_\alpha^{c_2}(T_{c_2}^*) \leq \mathbf{E}_\alpha^{c_2}(T_{c_1}^*)$ imply that

$$\begin{aligned} \mathbf{M}_\alpha(T_{c_1}^*) - c_1 \mathbf{M}(\partial T_{c_1}^*) &\leq \mathbf{M}_\alpha(T_{c_2}^*) - c_1 \mathbf{M}(\partial T_{c_2}^*) \\ \mathbf{M}_\alpha(T_{c_2}^*) - c_2 \mathbf{M}(\partial T_{c_2}^*) &\leq \mathbf{M}_\alpha(T_{c_1}^*) - c_2 \mathbf{M}(\partial T_{c_1}^*). \end{aligned}$$

Rewriting them gives

$$c_2 (\mathbf{M}(\partial T_{c_2}^*) - \mathbf{M}(\partial T_{c_1}^*)) \geq \mathbf{M}_\alpha(T_{c_2}^*) - \mathbf{M}_\alpha(T_{c_1}^*) \geq c_1 (\mathbf{M}(\partial T_{c_2}^*) - \mathbf{M}(\partial T_{c_1}^*)).$$

Since $c_1 < c_2$, we have $\mathbf{M}(\partial T_{c_2}^*) \geq \mathbf{M}(\partial T_{c_1}^*)$ and $\mathbf{M}_\alpha(T_{c_2}^*) \geq \mathbf{M}_\alpha(T_{c_1}^*)$. This shows that both $\mathbf{M}_\alpha(T_c^*)$ and $\mathbf{M}(\partial T_c^*)$ are increasing functions of c .

Moreover, by inequality (5.4), $\lim_{c \rightarrow \infty} \partial T_c^* = \lim_{c \rightarrow \infty} \nu_c^* - \mu_c^* = \nu - \mu$. Since d_α is a distance between measures of equal mass, by Proposition 5.1,

$$\begin{aligned} 0 &\leq d_\alpha(\mu, \nu) - \lim_{c \rightarrow \infty} \mathbf{M}_\alpha(T_c^*) \\ &= d_\alpha(\mu, \nu) - \lim_{c \rightarrow \infty} d_\alpha(\mu_c^*, \nu_c^*) \\ &\leq \lim_{c \rightarrow \infty} (d_\alpha(\mu, \mu_c^*) + d_\alpha(\nu, \nu_c^*)) = 0. \end{aligned}$$

Thus, $d_\alpha(\mu, \nu) = \lim_{c \rightarrow \infty} \mathbf{M}_\alpha(T_c^*)$. \square

Theorem 5.4. *Suppose μ and ν are two disjointly supported measures on X of equal mass, $1 - \frac{1}{m} < \alpha < 1$, and let $T_c^* \in \text{Path}(\mu_c^*, \nu_c^*)$ denote the solution to the ROTPB(μ, ν) problem corresponding to parameter c . If for some sequence $\{c_n\}$ converging to ∞ , the associated sequence $\{T_{c_n}^*\}$ is subsequentially convergent to T as rectifiable normal 1-currents with respect to flat convergence, then T is an α -optimal transport path from μ to ν .*

Proof. By the lower semi-continuity of \mathbf{M}_α and Proposition 5.3,

$$\mathbf{M}_\alpha(T) \leq \liminf_{n \rightarrow \infty} \mathbf{M}_\alpha(T_{c_n}^*) = d_\alpha(\mu, \nu).$$

Since $\partial T = \nu - \mu$, T itself is also a transport path from μ to ν , and it holds that $d_\alpha(\mu, \nu) \leq \mathbf{M}_\alpha(T)$. As a result, T is an optimal transport path. \square

Remark 5.5. Theorem 5.4 provides a novel perspective for approximating an optimal transport path. In light of this theorem, one can solve a sequence of ROTPB problems associated with a monotonically increasing series of $\{c_n\}$, and then use the limit of their solutions to obtain the desired path. For small values of c_n , the path $T_{c_n}^*$ is typically of simple structure and thus relatively easy to solve. As c_n rises, the planner would start moving more mass through transport paths of increasing

complexity, which eventually converge to an optimal transport path from μ to ν . We leave exploration along this line to future research.

REFERENCES

- [1] L. Ambrosio, N. Gigli. *A User's Guide to Optimal Transport*. In: *Modelling and Optimisation of Flows on Networks*. Lecture Notes in Mathematics, vol 2062. (2013) Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-32160-3-1>.
- [2] M. Bernot, V. Caselles, and J.-M. Morel. *Optimal transportation networks. Models and theory*. Lecture Notes in Mathematics, 1955. Springer, Berlin, 2009.
- [3] A. Brancolini and B. Wirth. *General transport problems with branched minimizers as functionals of 1-currents with prescribed boundary*. Calc. Var. Partial Differential Equations, 57(3):Art. 82, 39, 2018.
- [4] L.A. Caffarelli and R.J. McCann, *Free boundaries in optimal transport and Monge–Ampère obstacle problems*. Ann. of Math. 171 (2010) 673–730.
- [5] M. Colombo, A. De Rosa, and A. Marchese. *Improved stability of optimal traffic paths*. Calc. Var. Partial Differential Equations, 57:28, 2018.
- [6] M. Colombo, A. De Rosa, A. Marchese, and S. Stuvard. *On the lower semicontinuous envelope of functionals defined on polyhedral chains*. Nonlinear Analysis, Volume 163, November 2017, Pages 201-215
- [7] M. Colombo, A. De Rosa, and A. Marchese. *On the well-posedness of branched transportation*. Comm. Pure Appl. Math. 2020.
- [8] G. Devillanova and S. Solimini, *On the dimension of an irrigable measure* Rend. Sem. Mat. Univ. Padova, 117 (2007), pp. 1–49.
- [9] D. P. Thierry, R. Hardt. *Size minimization and approximating problems*, Calc. Var. Partial Differential Equations 17 (2003), 405-442.
- [10] H. Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153, Springer-Verlag New York Inc., New York 1969.
- [11] A. Figalli *The optimal partial transport problem* Arch. Ration. Mech. Anal. 195 (2010) 533–560
- [12] A. Marchese, A. Massaccesi, S. Stuvard, R. Tione. *A multi-material transport problem with arbitrary marginals*. arXiv:1807.10969v3.
- [13] F. Maddalena, S. Solimini, and J.M. Morel. *A variational model of irrigation patterns*. Interfaces Free Bound., 5(4):391–415, 2003.
- [14] A. Marchese and B. Wirth. *Approximation of rectifiable 1-currents and weak-* relaxation of the h-mass*. Journal of Mathematical Analysis and Applications, Volume 479, Issue 2: 2268-2283, 2019.
- [15] E. Paolini and E. Stepanov. *Optimal transportation networks as at chains*. Interfaces Free Bound., 8(4):393-436,2006.
- [16] E. Paolini and E. Stepanov. *Decomposition of acyclic normal currents in a metric space*. J. Funct. Anal., 263 (11):3358-3390, 2012.
- [17] P. Pegon, F. Santambrogio, and Q. Xia. *A fractal shape optimization problem in branched transport*. J. Math. Pures Appl. (9), 123:244–269, 2019.
- [18] F. Santambrogio. *Optimal Transport for Applied Mathematicians- Calculus of Variations, PDEs, and Modeling*, Progress in Nonlinear Differential Equations and Their Applications, Birkhäuser, 2015.
- [19] B. Schiffman. *α -balanced and efficient branched transport systems and their landscape functions*. PhD thesis. University of California at Davis, 2020.
- [20] L. Simon. *Lectures on geometric measure theory*, volume 3 of Proceedings of the Centre for Mathematical Analysis, Australian National University. Australian National University Centre for Mathematical Analysis, Canberra, 1983.
- [21] S. K. Smirnov. *Decomposition of solenoidal vector charges into elementary solenoids, and the structure of normal one-dimensional flows*. Algebra i Analiz, 5:206-238, 1993.
- [22] C. Villani. *Topics in mass transportation*. AMS Graduate Studies in Math. 58 (2003).
- [23] C. Villani. *Optimal transport: old and new*. Grundlehren der mathematischen Wissenschaften, (2009)
- [24] Q. Xia, *Optimal paths related to transport problems*, Communications in Contemporary Mathematics, 5 (2003), 251-279.
- [25] Q. Xia, *Interior Regularity of Optimal Transport Paths*, Calculus of Variations and Partial Differential Equations, 20, No. 3 (2004), 283–299.

- [26] Q. Xia, A. Vershynina. *On the transport dimension of measures*, SIAM J. MATH. ANAL., Vol. 41, No. 6, (2010) pp. 2407–2430.
- [27] Q. Xia, *Boundary regularity of optimal transport paths*, Adv. Calc. Var. Volume 4, Issue 2, (2011), 153–174.
- [28] Q. Xia, *Motivations, ideas and applications of ramified optimal transportation*, ESAIM Math. Model. Numer. Anal. Vol. 49 Number 6 (2015), 1791–1832.
Email address: qlxia@math.ucdavis.edu
Email address: xushaofeng@ruc.edu.cn