

ALMOST HADAMARD MATRICES: GENERAL THEORY AND EXAMPLES

TEODOR BANICA, ION NECHITA, AND KAROL ŻYCZKOWSKI

ABSTRACT. We develop a general theory of “almost Hadamard matrices”. These are by definition the matrices $H \in M_N(\mathbb{R})$ having the property that $U = H/\sqrt{N}$ is orthogonal, and is a local maximum of the 1-norm on $O(N)$. Our study includes a detailed discussion of the circulant case ($H_{ij} = \gamma_{j-i}$) and of the two-entry case ($H_{ij} \in \{x, y\}$), with the construction of several families of examples, and some 1-norm computations.

INTRODUCTION

An Hadamard matrix is a square matrix having ± 1 entries, whose rows are pairwise orthogonal. The simplest example, appearing at $N = 2$, is the Walsh matrix:

$$H_2 = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

At $N = 3$ we cannot have examples, due to the orthogonality condition, which forces N to be even. At $N = 4$ now, we have several examples, for instance $H_4 = H_2 \otimes H_2$:

$$H_4 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}$$

For higher values of N , the construction of Hadamard matrices is quite a tricky problem. First, by permuting rows and columns or by multiplying them by -1 , we can always assume that the first 3 rows of our matrix look as follows:

$$H = \begin{pmatrix} 1 \dots 1 & 1 \dots 1 & 1 \dots 1 & 1 \dots 1 \\ 1 \dots 1 & 1 \dots 1 & -1 \dots -1 & -1 \dots -1 \\ 1 \dots 1 & -1 \dots -1 & 1 \dots 1 & -1 \dots -1 \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

Now if we denote by x, y, z, t the sizes of the 4 columns, the orthogonality conditions between the first 3 rows give $x = y = z = t$, so $N = x + y + z + t$ is a multiple of 4.

2000 *Mathematics Subject Classification.* 05B20 (15B10).

Key words and phrases. Hadamard matrix, Orthogonal group.

A similar analysis with 4 rows instead of 3, or any other kind of abstract or concrete consideration doesn't give any further restriction on N , and we have:

Hadamard Conjecture (HC). *There is at least one Hadamard matrix of size $N \times N$, for any $N \in 4\mathbb{N}$.*

This conjecture, going back to the 19th century, is probably one of the most beautiful statements in combinatorics, and in mathematics in general. The numeric verification so far goes up to $N = 664$, see [24]. For a general presentation of the problem, see [28].

At the level of concrete examples, the only ones which are simple to describe are the tensor powers of the Walsh matrix, having size 2^k . For some other examples, see [20].

Yet another good problem, simple-looking as well, concerns the circulant case. Given a vector $\gamma \in (\pm 1)^N$, one can ask whether the matrix $H \in M_N(\pm 1)$ defined by $H_{ij} = \gamma_{j-i}$ is Hadamard or not. Here is a solution to the problem, appearing at $N = 4$:

$$K_4 = \begin{pmatrix} -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \end{pmatrix}$$

More generally, any vector $\gamma \in (\pm 1)^4$ satisfying $\sum \gamma_i = \pm 1$ is a solution to the problem. The following conjecture, going back to [34], states that there are no other solutions:

Circulant Hadamard Conjecture (CHC). *There is no circulant Hadamard matrix of size $N \times N$, for any $N \neq 4$.*

The fact that such a simple-looking problem is still open might seem quite surprising. If we denote by $S \subset \{1, \dots, N\}$ the set of positions of the -1 entries of γ , the Hadamard matrix condition is simply $|S \cap (S + k)| = |S| - N/4$, for any $k \neq 0$, taken modulo N . Thus, the above conjecture simply states that at $N \neq 4$, such a set S cannot exist!

Summarizing, the Hadamard matrices are very easy to introduce, and they quickly lead to some difficult and interesting combinatorial problems. Regarding now the other motivations for studying such matrices, these are quite varied:

- (1) The Hadamard matrices were first studied by Sylvester [36], who was seemingly attracted by their plastic beauty: just replace the ± 1 entries by black and white tiles, and admire the symmetries and dissymmetries of the resulting design!
- (2) More concretely now, the Hadamard matrices can be used for various coding purposes, and have several applications to engineering, and quantum physics. For instance the Walsh matrices $H_2^{\otimes k}$ are used in the Olivia MFSK radio protocol.

Most applications of the Hadamard matrices, however, come from their generalizations. A “complex Hadamard” matrix is a matrix $H \in M_N(\mathbb{C})$ all whose entries are on the unit circle, and whose rows are pairwise orthogonal. The basic example is $\tilde{F}_N = \sqrt{N}F_N$, where

F_N is the matrix of the Fourier transform over \mathbb{Z}_N . That is, with $\omega = e^{2\pi i/N}$:

$$\tilde{F}_N = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{N-1} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(N-1)} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \omega^{N-1} & \omega^{2(N-1)} & \dots & \omega^{(N-1)^2} \end{pmatrix}$$

As a first observation, the existence of this matrix prevents the existence of a “complex version” of the HC. However, when trying to construct complex Hadamard matrices by using roots of unity of a given order, a wide, subtle, and quite poorly understood generalization of the HC problematics appears. See e.g. [4], [12], [26], [27].

As for the motivations and applications, these partly come from pure mathematics, cf. e.g. [3], [17], [23], [32], and partly come from quantum physics, cf. e.g. [8], [37].

Let us go back now to the real case. Since the determinant of N vectors is maximized when these vectors are chosen pairwise orthogonal, we have the following result:

Theorem A. *For a matrix $H \in M_N(\pm 1)$ we have $|\det H| \leq N^{N/2}$, with equality if and only if H is Hadamard.*

This result, due to Hadamard himself [18], has led to a number of interesting problems, and to the general development of the theory of Hadamard matrices. See [20].

As already mentioned, in order for an Hadamard matrix to exist, its size N must be a multiple of 4. For numbers of type $N = 4n + k$ with $k = 1, 2, 3$, several “real” generalizations of the Hadamard matrices have been constructed. The idea is usually to consider matrices $H \in M_N(\pm 1)$, whose rows are as orthogonal as they can be:

Definition A. *A “quasi-Hadamard” matrix is a square matrix $H \in M_N(\pm 1)$ which is as orthogonal as possible, e.g. which maximizes the quantity $|\det H|$.*

This definition is of course a bit vague, but the main idea is there. For a detailed discussion of the different notions here, we refer to the articles [1], [25], [31].

Yet another interpretation of the Hadamard matrices comes from the Cauchy-Schwarz inequality. Since for $U \in O(N)$ we have $\|U\|_2 = \sqrt{N}$, we obtain:

Theorem B. *For a matrix $U \in O(N)$ we have $\|U\|_1 \leq N\sqrt{N}$, with equality if and only if $H = \sqrt{N}U$ is Hadamard.*

This result, first pointed out in [5], shows that the matrices of type $H = \sqrt{N}U$, with $U \in O(N)$ being a maximizer of the 1-norm on $O(N)$, can be thought of as being some kind of “analytic generalizations” of the Hadamard matrices. Note that such matrices exist for any N , in particular for $N = 4n + k$ with $k = 1, 2, 3$.

As an example, the maximum of the 1-norm on $O(3)$ can be shown to be the number 5, coming from the matrix $U_3 = K_3/\sqrt{3}$ and its various conjugates, where:

$$K_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$$

This result, proved in [5] by using the Euler-Rodrigues formula, is actually something quite accidental. In general, the integration on $O(N)$ is quite a subtle business, and the maximum of the 1-norm is quite difficult to approach. See [6], [7], [15].

These integration problems make the above type of matrices quite hard to investigate. Instead of looking directly at them, we will rather enlarge the attention to the matrices of type $H = \sqrt{N}U$, where $U \in O(N)$ is a local maximizer of the 1-norm on $O(N)$. Indeed, according to the Hessian computations in [5], these latter matrices are characterized by the fact that all their entries are nonzero, and $SU^t > 0$, where $S_{ij} = \text{sgn}(U_{ij})$.

Summarizing, Theorem B suggests the following definition:

Definition B. An “almost Hadamard” matrix is a square matrix $H \in M_N(\mathbb{R})$ such that $U = H/\sqrt{N}$ is orthogonal, and is a local maximum of the 1-norm on $O(N)$.

The basic examples are of course the Hadamard matrices. We have as well the following $N \times N$ matrix, with $N \in \mathbb{N}$ arbitrary, which generalizes the above matrices K_3, K_4 :

$$K_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 2 - N & 2 & \dots & 2 & 2 \\ 2 & 2 - N & \dots & \dots & 2 \\ \dots & \dots & \dots & \dots & \dots \\ 2 & \dots & \dots & \dots & \dots \\ 2 & 2 & \dots & \dots & 2 - N \end{pmatrix}$$

A lot of other interesting examples exist, as we will show in this paper. Here is for instance a remarkable matrix, having order $N \in 2\mathbb{N} + 1$, and circulant structure:

$$L_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 1 & -\cos^{-1} \frac{\pi}{N} & \cos^{-1} \frac{2\pi}{N} & \dots & \cos^{-1} \frac{(N-1)\pi}{N} \\ \cos^{-1} \frac{(N-1)\pi}{N} & 1 & -\cos^{-1} \frac{\pi}{N} & \dots & -\cos^{-1} \frac{(N-2)\pi}{N} \\ \dots & \dots & \dots & \dots & \dots \\ -\cos^{-1} \frac{\pi}{N} & \cos^{-1} \frac{2\pi}{N} & -\cos^{-1} \frac{3\pi}{N} & \dots & 1 \end{pmatrix}$$

Yet another series, with $N = q^2 + q + 1$, where $q = p^k$ is a prime power, comes from the adjacency matrix of the projective plane over \mathbb{F}_q . Here is for instance the matrix

associated to the Fano plane ($q = 2$), where $x = 2 - 4\sqrt{2}$, $y = 2 + 3\sqrt{2}$:

$$I_7 = \frac{1}{2\sqrt{7}} \begin{pmatrix} x & x & y & y & y & x & y \\ y & x & x & y & y & y & x \\ x & y & x & x & y & y & y \\ y & x & y & x & x & y & y \\ y & y & x & y & x & x & y \\ y & y & y & x & y & x & x \\ x & y & y & y & x & y & x \end{pmatrix}$$

The aim of the present paper is to provide a systematic study of such matrices, with the construction of a number of non-trivial examples, and with the development of some general theory as well. Our motivation comes from two kinds of problems:

- (1) The Hadamard matrix problematics. The world of Hadamard matrices is extremely rigid, and we think that our study of almost Hadamard matrices, where there is much more freedom, can help. As an example, there are several non-trivial classes of circulant almost Hadamard matrices at any $N \in \mathbb{N}$, and trying to understand them might end up in shedding some new light on the CHC.
- (2) Generalizations of Hadamard matrices. The Hadamard matrices have applications in a number of areas of physics and engineering, notably in coding theory, and in various branches of quantum physics. One problem, however, is that these matrices exist only at $N = 4n$. At $N = 4n + k$ with $k = 1, 2, 3$ some generalizations would be needed, and we believe that our almost Hadamard matrices can help.

The paper is organized as follows: 1 is a preliminary section, in 2-3 we investigate two special cases, namely the circulant case and the two-entry case, and 4 contains a list of examples. The final section, 5, contains a few concluding remarks.

Acknowledgements. We would like to thank Guillaume Aubrun for several useful discussions. The work of T.B. was supported by the ANR grant ‘‘Granma’’. I.N. acknowledges financial support from the ANR project OSvsQPI 2011 BS01 008 01 and from a CNRS PEPS grant. The work of K.Z. was supported by the grant N N202 261938, financed by the Polish Ministry of Science and Higher Education.

1. PRELIMINARIES

We consider in this paper various square matrices $M \in M_N(\mathbb{C})$. The indices of our matrices will usually range in the set $\{0, 1, \dots, N - 1\}$.

Definition 1.1. *We use the following special $N \times N$ matrices:*

- (1) 1_N is the identity matrix.
- (2) J_N is the ‘‘flat’’ matrix, having all entries equal to $1/N$.
- (3) F_N is the Fourier matrix, given by $(F_N)_{ij} = \omega^{ij}/\sqrt{N}$, with $\omega = e^{2\pi i/N}$.

We denote by D the generic diagonal matrices, by U the generic orthogonal or unitary matrices, and by H the generic Hadamard matrices, and their generalizations.

Our starting point is the following observation, from [5]:

Proposition 1.2. *For $U \in O(N)$ we have $\|U\|_1 \leq N\sqrt{N}$, with equality if and only if $H = \sqrt{N}U$ is Hadamard.*

Proof. The first assertion follows from the Cauchy-Schwarz inequality:

$$\sum_{ij} |U_{ij}| \leq \left(\sum_{ij} 1^2 \right)^{1/2} \left(\sum_{ij} |U_{ij}|^2 \right)^{1/2} = N\sqrt{N}$$

For having equality the numbers $|U_{ij}|$ must be equal, and since the sum of squares of these numbers is N , we must have $|U_{ij}| = 1/\sqrt{N}$, which gives the result. \square

As already mentioned in the introduction, the study of maximizers of the 1-norm on $O(N)$ is a quite difficult task. However, we have here the following result, from [5]:

Theorem 1.3. *For $U \in O(N)$, the following are equivalent:*

- (1) U is a local maximizer of the 1-norm on $O(N)$.
- (2) $U_{ij} \neq 0$, and $SU^t > 0$, where $S_{ij} = \text{sgn}(U_{ij})$.

Proof. As already mentioned, this result is from [5]. Here is the idea of the proof:

Let us first prove that if U is a local maximizer of the 1-norm, then $U_{ij} \neq 0$. For this purpose, let U_1, \dots, U_N be the columns of U , and let us perform a rotation of U_1, U_2 :

$$\begin{pmatrix} U_1^t \\ U_2^t \end{pmatrix} = \begin{pmatrix} \cos t \cdot U_1 - \sin t \cdot U_2 \\ \sin t \cdot U_1 + \cos t \cdot U_2 \end{pmatrix}$$

In order to compute the 1-norm, let us permute the columns of U , in such a way that the first two rows look as follows, with $X_k \neq 0, Y_k \neq 0, A_k C_k > 0, B_k D_k < 0$:

$$\begin{pmatrix} U_1 \\ U_2 \end{pmatrix} = \begin{pmatrix} 0 & 0 & Y & A & B \\ 0 & X & 0 & C & D \end{pmatrix}$$

If we agree that the lower-case letters denote the 1-norms of the corresponding upper-case vectors, and we let $K = u_3 + \dots + u_N$, then for $t > 0$ small we have:

$$\begin{aligned} \|U^t\|_1 &= \|\cos t \cdot U_1 - \sin t \cdot U_2\|_1 + \|\sin t \cdot U_1 + \cos t \cdot U_2\|_1 + K \\ &= (\cos t + \sin t)(x + y + a + d) + (\cos t - \sin t)(b + c) + K \end{aligned}$$

Now since U locally maximizes the 1-norm on $O(N)$, the derivative of this quantity must be negative in the limit $t \rightarrow 0$. So, we obtain the following inequality:

$$(x + y + a + d) - (b + c) \leq 0$$

Consider now the matrix obtained by interchanging U_1, U_2 . Since this matrix must be as well a local maximizer of the 1-norm on $O(N)$, we obtain:

$$(x + y + b + c) - (a + d) \leq 0$$

We deduce that $x + y = 0$, so $x = y = 0$, and the 0 entries of U_1, U_2 must appear at the same positions. By permuting the rows of U the same must hold for any two rows U_i, U_j . Now since U cannot have zero columns, all its entries must be nonzero, as claimed.

It remains to prove that if $U_{ij} \neq 0$, then U is a local maximizer of $F(U) = \|U\|_1$ if and only if $SU^t > 0$, where $S_{ij} = \text{sgn}(U_{ij})$. For this purpose, we differentiate F :

$$dF = \sum_{ij} S_{ij} dU_{ij}$$

We know that $O(N)$ consists of the zeroes of the polynomials $A_{ij} = \sum_k U_{ik}U_{jk} - \delta_{ij}$. So, U is a critical point of F if and only if $dF \in \text{span}(dA_{ij})$. Now since $A_{ij} = A_{ji}$, this is the same as asking for a symmetric matrix M such that $dF = \sum_{ij} M_{ij} dA_{ij}$. But:

$$\sum_{ij} M_{ij} dA_{ij} = \sum_{ijk} M_{ij} (U_{ik} dU_{jk} + U_{jk} dU_{ik}) = 2 \sum_{lk} (MU)_{lk} dU_{lk}$$

Thus the critical point condition reads $S = 2MU$, so the matrix $M = SU^t/2$ must be symmetric. Now the Hessian of F applied to a vector $X = UY$, with $Y \in O(N)$, is:

$$\text{Hess}(F)(X) = \frac{1}{2} \text{Tr}(X^t \cdot SU^t \cdot X) = \frac{1}{2} \text{Tr}(Y^t \cdot U^t S \cdot Y)$$

Thus the Hessian of F is positive definite when $U^t S$ is positive definite, which is the same as saying that $U(U^t S)U^t = SU^t$ is positive definite, and we are done. \square

The above result gives rise to the following definition:

Definition 1.4. A square matrix $H \in M_N(\mathbb{R}^*)$ is called “almost Hadamard” if $U = H/\sqrt{N}$ is orthogonal, and the following equivalent conditions are satisfied:

- (1) U is a local maximizer of the 1-norm on $O(N)$.
- (2) $U_{ij} \neq 0$, and with $S_{ij} = \text{sgn}(U_{ij})$, we have $SU^t > 0$.

If so is the case, we call H “optimal” if U is a maximizer of the 1-norm on $O(N)$.

Let J_N be the flat $N \times N$ matrix, having all the entries equal to $1/N$. Also, let us call “Hadamard equivalence”, or just “equivalence”, the equivalence relation on the $N \times N$ matrices coming from permuting the rows and columns, or multiplying them by -1 .

Proposition 1.5. The class of almost Hadamard matrices has the following properties:

- (1) It contains all the Hadamard matrices.
- (2) It contains the matrix $K_N = \sqrt{N}(2J_N - 1_N)$.
- (3) It is stable under equivalence, tensor products, and transposition.

Proof. All the assertions are clear from definitions:

(1) This follows either from Proposition 1.2, or from the fact that $U = H/\sqrt{N}$ is orthogonal, and $SU^t = HU^t = \sqrt{N}1_N$ is positive.

(2) First, the matrix $U = K_N/\sqrt{N}$ is orthogonal, because it is symmetric, and:

$$U^2 = (2J_N - 1_N)^2 = 4J_N^2 - 4J_N + 1_N = 1_N$$

Also, we have $S = NJ_N - 21_N$, so the matrix SU^t is indeed positive:

$$SU^t = (NJ_N - 21_N)(2J_N - 1_N) = (N - 2)J_N + 2(1_N - J_N)$$

(3) For a tensor product of almost Hadamard matrices $H = H' \otimes H''$ we have $U = U' \otimes U''$ and $S = S' \otimes S''$, so that U is unitary and SU^t is positive, as claimed. As for the assertions regarding equivalence and transposition, these are clear from definitions. \square

Regarding now the optimal case, we have the following result, from [5]:

Proposition 1.6. *The optimal almost Hadamard matrices are as follows:*

- (1) *At any N where HC holds, these are the $N \times N$ Hadamard matrices.*
- (2) *At $N = 3$, these are precisely $K_3 = \sqrt{3}(2J_3 - 1_3)$ and its conjugates.*

Proof. The assertion (1) is clear from Proposition 1.2. For (2) we must prove that for $U \in O(3)$ we have $\|U\|_1 \leq 5$, with equality when U is conjugate to $U_3 = 2J_3 - 1_3$. But here we can assume $U \in SO(3)$, and use the Euler-Rodrigues formula:

$$U = \begin{pmatrix} x^2 + y^2 - z^2 - t^2 & 2(yz - xt) & 2(xz + yt) \\ 2(xt + yz) & x^2 + z^2 - y^2 - t^2 & 2(zt - xy) \\ 2(yt - xz) & 2(xy + zt) & x^2 + t^2 - y^2 - z^2 \end{pmatrix}$$

Here $(x, y, z, t) \in S^3$ comes from the standard cover map $S^3 \simeq SU(2) \rightarrow SO(3)$. Now by linearizing, we must prove that for any $(x, y, z, t) \in \mathbb{R}^4$ we have:

$$\|U\|_1 \leq 5(x^2 + y^2 + z^2 + t^2)$$

The proof of this latter inequality is routine, and the equality situation turns to hold indeed exactly for the matrix $U_3 = 2J_3 - 1_3$ and its conjugates. See [5]. \square

Finally, let us mention that a version of Proposition 1.2 above, using the Hölder inequality, shows that the matrices of type $U = H/\sqrt{N}$ with $H \in M_N(\pm 1)$ Hadamard maximize the p -norm on $O(N)$ at $p \in [1, 2)$, and minimize it at $p \in (2, \infty]$. See [5]. Part of the above $p = 1$ results extend to the general setting $p \in [1, \infty] - \{2\}$, and in particular to the exponents $p = 4$ and $p = \infty$, which are of particular interest in connection with several quantum physics questions. This will be discussed in a forthcoming paper.

2. THE CIRCULANT CASE

In this section we study the almost Hadamard matrices which are circulant. We recall that a matrix $H \in M_N(\mathbb{C})$ is called circulant if it is of the form:

$$H = \begin{pmatrix} \gamma_0 & \gamma_1 & \cdots & \gamma_{N-1} \\ \gamma_{N-1} & \gamma_0 & \cdots & \gamma_{N-2} \\ \cdots & \cdots & \cdots & \cdots \\ \gamma_1 & \gamma_2 & \cdots & \gamma_0 \end{pmatrix}$$

Let $F \in U(N)$ be the Fourier matrix, given by $F_{ij} = \omega^{ij}/\sqrt{N}$, where $\omega = e^{2\pi i/N}$. Given a vector $\alpha \in \mathbb{C}^n$, we associate to it the diagonal matrix $\alpha' = \text{diag}(\alpha_0, \dots, \alpha_{N-1})$.

We will make a heavy use of the following well-known result:

Proposition 2.1. *For a matrix $H \in M_N(\mathbb{C})$, the following are equivalent:*

- (1) H is circulant, i.e. $H_{ij} = \gamma_{j-i}$, for a certain vector $\gamma \in \mathbb{C}^N$.
- (2) H is Fourier-diagonal, i.e. $H = FDF^*$, with $D \in M_N(\mathbb{C})$ diagonal.

In addition, if so is the case, then with $D = \sqrt{N}\alpha'$ we have $\gamma = F\alpha$.

Proof. (1) \implies (2) The matrix $D = F^*HF$ is indeed diagonal, given by:

$$D_{ij} = \frac{1}{N} \sum_{kl} \omega^{jl-ik} \gamma_{l-k} = \delta_{ij} \sum_r \omega^{jr} \gamma_r$$

(2) \implies (1) The matrix $H = FDF^*$ is indeed circulant, given by:

$$H_{ij} = \sum_k F_{ik} D_{kk} \bar{F}_{jk} = \frac{1}{N} \sum_k \omega^{(i-j)k} D_{kk}$$

Finally, the last assertion is clear from the above formula of H_{ij} . \square

Let us investigate now the circulant orthogonal matrices. We let the matrix indices i, j vary modulo N . We denote by \mathbb{T} the unit circle in the complex plane.

Lemma 2.2. *For a matrix $U \in M_N(\mathbb{C})$, the following are equivalent:*

- (1) U is orthogonal and circulant.
- (2) $U = F\alpha'F^*$ with $\alpha \in \mathbb{T}^N$ satisfying $\bar{\alpha}_i = \alpha_{-i}$ for any i .

Proof. We will use many times the fact that given $\alpha \in \mathbb{C}^N$, the vector $\gamma = F\alpha$ is real if and only if $\bar{\alpha}_i = \alpha_{-i}$ for any i . This follows indeed from $\overline{F\alpha} = F\tilde{\alpha}$, with $\tilde{\alpha}_i = \bar{\alpha}_{-i}$.

(1) \implies (2) Write $H_{ij} = \gamma_{j-i}$ with $\gamma \in \mathbb{R}^N$. By using Proposition 2.1 we obtain $H = FDF^*$ with $D = \sqrt{N}\alpha'$ and $\gamma = F\alpha$. Now since $U = F\alpha'F^*$ is unitary, so is α' , so we must have $\alpha \in \mathbb{T}^N$. Finally, since γ is real we have $\bar{\alpha}_i = \alpha_{-i}$, and we are done.

(2) \implies (1) We know from Proposition 2.1 that U is circulant. Also, from $\alpha \in \mathbb{T}^N$ we obtain that α' is unitary, and so must be U . Finally, since we have $\bar{\alpha}_i = \alpha_{-i}$, the vector $\gamma = F\alpha$ is real, and hence we have $U \in M_N(\mathbb{R})$, which finishes the proof. \square

Let us discuss now the almost Hadamard case. First, in the usual Hadamard case, the known examples and the corresponding α -vectors are as follows:

Proposition 2.3. *The known circulant Hadamard matrices, namely*

$$\pm \begin{pmatrix} -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \end{pmatrix}, \pm \begin{pmatrix} 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \\ -1 & 1 & 1 & 1 \end{pmatrix}, \pm \begin{pmatrix} 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \\ -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \end{pmatrix}, \pm \begin{pmatrix} 1 & 1 & 1 & -1 \\ -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \end{pmatrix}$$

come from the vectors $\alpha = \pm(1, -1, -1, -1), \pm(1, -i, 1, i), \pm(1, 1, -1, 1), \pm(1, i, 1, -i)$.

Proof. At $N = 4$ the conjugate of the Fourier matrix is given by:

$$F^* = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -i & -1 & i \\ 1 & -1 & 1 & -1 \\ 1 & i & -1 & -i \end{pmatrix}$$

Thus the vectors $\alpha = F^*\gamma$ are indeed those in the statement. \square

We have the following ‘‘almost Hadamard’’ generalization of the above matrices:

Proposition 2.4. *If $q^N = 1$ then the vector $\alpha = \pm(1, -q, -q^2, \dots, -q^{N-1})$ produces an almost Hadamard matrix, which is equivalent to $K_N = \sqrt{N}(2J_N - 1_N)$.*

Proof. Observe first that these matrices generalize those in Proposition 2.3. Indeed, at $N = 4$ the choices for q are $1, i, -1, -i$, and this gives the above α -vectors.

Assume that the \pm sign in the statement is $+$. With $q = \omega^r$, we have:

$$\sqrt{N}\gamma_i = \sum_{k=0}^{N-1} \omega^{ik} \alpha_k = 1 - \sum_{k=1}^{N-1} \omega^{(i+r)k} = 2 - \sum_{k=0}^{N-1} \omega^{(i+r)k} = 2 - \delta_{i,-r}N$$

In terms of the standard long cycle $(C_N)_{ij} = \delta_{i+1,j}$, we obtain:

$$H = \sqrt{N}(2J_N - C_N^{-r})$$

Thus H is equivalent to K_N , and by Proposition 1.5, it is almost Hadamard. \square

In general, the construction of circulant almost Hadamard matrices is quite a tricky problem. At the abstract level, we have the following technical result:

Lemma 2.5. *A circulant matrix $H \in M_N(\mathbb{R}^*)$, written $H_{ij} = \gamma_{j-i}$, is almost Hadamard if and only if the following conditions are satisfied:*

- (1) *The vector $\alpha = F^*\gamma$ satisfies $\alpha \in \mathbb{T}^N$.*
- (2) *With $\varepsilon = \text{sgn}(\gamma)$, $\rho_i = \sum_r \varepsilon_r \gamma_{i+r}$ and $\nu = F^*\rho$, we have $\nu > 0$.*

In addition, if so is the case, then $\bar{\alpha}_i = \alpha_{-i}$, $\rho_i = \rho_{-i}$ and $\nu_i = \nu_{-i}$ for any i .

Proof. According to Definition 1.4 our matrix H is almost Hadamard if and only if the matrix $U = H/\sqrt{N}$ is orthogonal and $SU^t > 0$, where $S_{ij} = \text{sgn}(U_{ij})$. By Lemma 2.2 the orthogonality of U is equivalent to the condition (1). Regarding now the condition $SU^t > 0$, this is equivalent to $S^tU > 0$. But, with $k = i - r$, we have:

$$(S^tH)_{ij} = \sum_k S_{ki}H_{kj} = \sum_k \varepsilon_{i-k}\gamma_{j-k} = \sum_r \varepsilon_r\gamma_{j-i+r} = \rho_{j-i}$$

Thus S^tU is circulant, with ρ/\sqrt{N} as first row. From Proposition 2.1 we get $S^tU = FLF^*$ with $L = \nu'$ and $\nu = F^*\rho$, so $S^tU > 0$ iff $\nu > 0$, which is the condition (2).

Finally, the assertions about α, ν follow from the fact that $F\alpha, F\nu$ are real. As for the assertion about ρ , this follows from the fact that S^tU is symmetric. \square

Theorem 2.6. *For N odd the following matrix is almost Hadamard,*

$$L_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 1 & -\cos^{-1} \frac{\pi}{N} & \cos^{-1} \frac{2\pi}{N} & \dots & \cos^{-1} \frac{(N-1)\pi}{N} \\ \cos^{-1} \frac{(N-1)\pi}{N} & 1 & -\cos^{-1} \frac{\pi}{N} & \dots & -\cos^{-1} \frac{(N-2)\pi}{N} \\ \dots & \dots & \dots & \dots & \dots \\ -\cos^{-1} \frac{\pi}{N} & \cos^{-1} \frac{2\pi}{N} & -\cos^{-1} \frac{3\pi}{N} & \dots & 1 \end{pmatrix}$$

and comes from an α -vector having all entries equal to 1 or -1 .

Proof. Write $N = 2n + 1$, and consider the following vector:

$$\alpha_i = \begin{cases} (-1)^{n+i} & \text{for } i = 0, 1, \dots, n \\ (-1)^{n+i+1} & \text{for } i = n+1, \dots, 2n \end{cases}$$

Let us first prove that $(L_N)_{ij} = \gamma_{j-i}$, where $\gamma = F\alpha$. With $\omega = e^{2\pi i/N}$ we have:

$$\sqrt{N}\gamma_i = \sum_{j=0}^{2n} \omega^{ij} \alpha_j = \sum_{j=0}^n (-1)^{n+j} \omega^{ij} + \sum_{j=1}^n (-1)^{n+(N-j)+1} \omega^{i(N-j)}$$

Now since N is odd, and since $\omega^N = 1$, we obtain:

$$\sqrt{N}\gamma_i = \sum_{j=0}^n (-1)^{n+j} \omega^{ij} + \sum_{j=1}^n (-1)^{n-j} \omega^{-ij} = \sum_{j=-n}^n (-1)^{n+j} \omega^{ij}$$

By computing the sum on the right, with $\xi = e^{\pi i/N}$ we get, as claimed:

$$\sqrt{N}\gamma_i = \frac{2\omega^{-ni}}{1 + \omega^i} = \frac{2\xi^{-2ni}}{1 + \xi^{2i}} = \frac{2\xi^{-Ni}}{\xi^{-i} + \xi^i} = (-1)^i \cos^{-1} \frac{i\pi}{N}$$

In order to prove now that L_N is almost Hadamard, we use Lemma 2.5. Since the sign vector is simply $\varepsilon = (-1)^n \alpha$, the vector $\rho_i = \sum_r \varepsilon_r \gamma_{i+r}$ is given by:

$$\sqrt{N} \rho_i = (-1)^n \sum_{r=0}^{2n} \alpha_r \sum_{j=-n}^n (-1)^{n+j} \omega^{(i+r)j} = \sum_{j=-n}^n (-1)^j \omega^{ij} \sum_{r=0}^{2n} \alpha_r \omega^{rj}$$

Now since the last sum on the right is $(\sqrt{N} F \alpha)_j = \sqrt{N} \gamma_j$, we obtain:

$$\rho_i = \sum_{j=-n}^n (-1)^j \omega^{ij} \gamma_j = \frac{1}{\sqrt{N}} \sum_{j=-n}^n (-1)^j \omega^{ij} \sum_{k=-n}^n (-1)^{n+k} \omega^{jk}$$

Thus we have the following formula:

$$\rho_i = \frac{(-1)^n}{\sqrt{N}} \sum_{j=-n}^n \sum_{k=-n}^n (-1)^{j+k} \omega^{(i+k)j}$$

Let us compute now the vector $\nu = F^* \rho$. We have:

$$\nu_l = \frac{1}{\sqrt{N}} \sum_{i=0}^{2n} \omega^{-il} \rho_i = \frac{(-1)^n}{N} \sum_{j=-n}^n \sum_{k=-n}^n (-1)^{j+k} \omega^{jk} \sum_{i=0}^{2n} \omega^{i(j-l)}$$

The sum on the right is $N \delta_{jl}$, with both j, l taken modulo N , so it is equal to $N \delta_{jL}$, where $L = l$ for $l \leq n$, and $L = l - N$ for $l > n$. We get:

$$\nu_l = (-1)^n \sum_{k=-n}^n (-1)^{L+k} \omega^{Lk} = (-1)^{n+L} \sum_{k=-n}^n (-\omega^L)^k$$

With $\xi = e^{\pi i/N}$, this gives the following formula:

$$\nu_l = (-1)^{n+L} \frac{2(-\omega^L)^{-n}}{1 + \omega^L} = (-1)^L \frac{2\omega^{-nL}}{1 + \omega^L}$$

In terms of the variable $\xi = e^{\pi i/N}$, we obtain:

$$\nu_l = (-1)^L \frac{2\xi^{-2nL}}{1 + \xi^{2L}} = (-1)^L \frac{2\xi^{-NL}}{\xi^{-L} + \xi^L} = \cos^{-1} \frac{L\pi}{N}$$

Now since $L \in [-n, n]$, all the entries of ν are positive, and we are done. \square

At the level of examples now, at $N = 3$ we obtain the matrix $L_3 = -K_3$. At $N = 5$ we obtain a matrix having as entries 1 and $x = -\cos^{-1} \frac{\pi}{5}$, $y = \cos^{-1} \frac{2\pi}{5}$:

$$L_5 = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 & x & y & y & x \\ x & 1 & x & y & y \\ y & x & 1 & x & y \\ y & y & x & 1 & x \\ x & y & y & x & 1 \end{pmatrix}$$

Let us look now more in detail at the vectors $\alpha \in \mathbb{T}^N$ appearing in Proposition 2.4 and in the proof of Theorem 2.6. In both cases we have $\alpha_i^2 = \omega^{ri}$ for a certain $r \in \mathbb{N}$, and this might suggest that any circulant almost Hadamard matrix should come from a vector $\alpha \in \mathbb{T}^N$ having the property that α^2 is formed by roots of unity in a progression.

However, the rescaled adjacency matrix of the Fano plane, to be discussed in the next section, is circulant almost Hadamard, but does not have this property. The problem of finding the correct extension of the circulant Hadamard conjecture to the almost Hadamard matrix case is a quite subtle one, that we would like to raise here.

3. THE TWO-ENTRY CASE

In this section we study the almost Hadamard matrices having only two entries, $H \in M_N(x, y)$, with $x, y \in \mathbb{R}$. As a first remark, the usual Hadamard matrices $H \in M_N(\pm 1)$ are of this form. However, when trying to build a combinatorial hierarchy of the two-entry almost Hadamard matrices, the usual Hadamard matrices stand on top, and there is not so much general theory that can be developed, as to cover them.

We will therefore restrict attention to the following special type of matrices:

Definition 3.1. *An (a, b, c) pattern is a matrix $M \in M_N(x, y)$, with $N = a + 2b + c$, such that, in any two rows, the number of $x/y/x/y$ sitting below $x/x/y/y$ is $a/b/b/c$.*

In other words, given any two rows of our matrix, we are asking for the existence of a permutation of the columns such that these two rows become:

$$\begin{array}{cccc} x \dots x & x \dots x & y \dots y & y \dots y \\ \underbrace{x \dots x}_a & \underbrace{y \dots y}_b & \underbrace{x \dots x}_b & \underbrace{y \dots y}_c \end{array}$$

Observe that the Hadamard matrices do not come in general from patterns. However, there are many interesting examples of patterns coming from block designs [14], [35]:

Definition 3.2. *A (v, k, λ) symmetric balanced incomplete block design is a collection B of subsets of a set X , called blocks, with the following properties:*

- (1) $|X| = |B| = v$.
- (2) *Each block contains exactly k points from X .*
- (3) *Each pair of distinct points is contained in exactly λ blocks of B .*

The incidence matrix of a such block design is the $v \times v$ matrix defined by:

$$M_{bx} = \begin{cases} 1 & \text{if } x \in b \\ 0 & \text{if } x \notin b \end{cases}$$

The connection between designs and patterns comes from:

Proposition 3.3. *If $N = a + 2b + c$ then the adjacency matrix of any $(N, a + b, a)$ symmetric balanced incomplete block design is an (a, b, c) pattern.*

Proof. Indeed, let us replace the 0 – 1 values in the adjacency matrix M by abstract $x - y$ values. Then each row of M contains $a + b$ copies of x and $b + c$ copies of y , and since every pair of distinct blocks intersect in exactly a points, cf. [35], we see that every pair of rows has exactly a variables x in matching positions, so that M is an (a, b, c) pattern. \square




FIGURE 1. The Fano plane

As a first example, consider the Fano plane. The sets X, B of points and lines form a $(7, 3, 1)$ block design, corresponding to the following $(1, 2, 2)$ pattern:

$$I_7 = \begin{pmatrix} x & x & y & y & y & x & y \\ y & x & x & y & y & y & x \\ x & y & x & x & y & y & y \\ y & x & y & x & x & y & y \\ y & y & x & y & x & x & y \\ y & y & y & x & y & x & x \\ x & y & y & y & x & y & x \end{pmatrix}$$

Now remember that the Fano plane is the projective plane over $\mathbb{F}_2 = \{0, 1\}$. The same method works with \mathbb{F}_2 replaced by an arbitrary finite field \mathbb{F}_q , and we get:

Proposition 3.4. *Assume that $q = p^k$ is a prime power. Then the point-line incidence matrix of the projective plane over \mathbb{F}_q is a $(1, q, q^2 - q)$ pattern.*

Proof. The sets X, B of points and lines of the projective plane over \mathbb{F}_q are known to form a $(q^2 + q + 1, q + 1, 1)$ block design, and this gives the result. \square

There are many other interesting examples of symmetric balanced incomplete block designs, all giving rise to patterns, via Proposition 3.3. For instance the famous Paley biplane [11], pictured below, is a $(11, 5, 2)$ block design, and hence gives rise to a $(2, 3, 3)$ pattern. When assigning certain special values to the parameters x, y we obtain a 11×11 almost Hadamard matrix, that we believe to be optimal. See section 4 below.




FIGURE 2. The Paley biplane

We consider now the problem of associating real values to the symbols x, y in an (a, b, c) pattern such that the resulting matrix $U(x, y)$ is orthogonal.

Lemma 3.5. *Given $a, b, c \in \mathbb{N}$, there exists an orthogonal matrix having pattern (a, b, c) iff $b^2 \geq ac$. In this case the solutions are $U(x, y)$ and $-U(x, y)$, where*

$$x = -\frac{t}{\sqrt{b(t+1)}}, \quad y = \frac{1}{\sqrt{b(t+1)}}$$

where $t = (b \pm \sqrt{b^2 - ac})/a$ can be any of the solutions of $at^2 - 2bt + c = 0$.

Proof. First, in order for U to be orthogonal, the following conditions must be satisfied:

$$ax^2 + 2bxy + cy^2 = 0, \quad (a+b)x^2 + (b+c)y^2 = 1$$

The first condition, coming from the orthogonality of rows, tells us that $t = -x/y$ must be the variable in the statement. As for the second condition, this becomes:

$$y^2 = \frac{1}{(a+b)t^2 + (b+c)} = \frac{1}{(at^2 + c) + (bt^2 + b)} = \frac{1}{2bt + bt^2 + b} = \frac{1}{b(t+1)^2}$$

This gives the above formula of y , and hence the formula of $x = -ty$ as well. \square

Lemma 3.6. *Let $U = U(x, y)$ be orthogonal, corresponding to an (a, b, c) pattern. Then $H = \sqrt{N}U$ is almost Hadamard iff $(N(a-b) + 2b)|x| + (N(c-b) + 2b)|y| \geq 0$.*

Proof. We use the criterion in Definition 1.4 (2). So, let $S_{ij} = \text{sgn}(U_{ij})$. Since any row of U consists of $a+b$ copies of x and $b+c$ copies of y , we have:

$$(SU^t)_{ii} = \sum_k \text{sgn}(U_{ik})U_{ik} = (a+b)|x| + (b+c)|y|$$

Regarding now $(SU^t)_{ij}$ with $i \neq j$, we can assume in the computation that the i -th and j -th row of U are exactly those pictured after Definition 3.1 above. Thus:

$$\begin{aligned} (SU^t)_{ij} &= \sum_k \operatorname{sgn}(U_{ik})U_{jk} \\ &= a \operatorname{sgn}(x)x + b \operatorname{sgn}(x)y + b \operatorname{sgn}(y)x + c \operatorname{sgn}(y)y \\ &= a|x| - b|y| - b|x| + c|y| \\ &= (a - b)|x| + (c - b)|y| \end{aligned}$$

We obtain the following formula for the matrix SU^t itself:

$$\begin{aligned} SU^t &= 2b(|x| + |y|)1_N + ((a - b)|x| + (c - b)|y|)NJ_N \\ &= 2b(|x| + |y|)(1_N - J_N) + ((N(a - b) + 2b)|x| + (N(c - b) + 2b)|y|)J_N \end{aligned}$$

Now since the matrices $1_N - J_N$, J_N are orthogonal projections, we have $SU^t > 0$ if and only if the coefficients of these matrices in the above expression are both positive. Since the coefficient of $1_N - J_N$ is clearly positive, the condition left is:

$$(N(a - b) + 2b)|x| + (N(c - b) + 2b)|y| \geq 0$$

So, we have obtained the condition in the statement, and we are done. \square

Proposition 3.7. *Assume that $a, b, c \in \mathbb{N}$ satisfy $c \geq a$ and $b(b - 1) = ac$, and consider the (a, b, c) pattern $U = U(x, y)$, where:*

$$x = \frac{a + (1 - a - b)\sqrt{b}}{Na}, \quad y = \frac{b + (a + b)\sqrt{b}}{Nb}$$

Then $H = \sqrt{N}U$ is an almost Hadamard matrix.

Proof. We have $b^2 - ac = b$, so Lemma 2.5 applies, and shows that with $t = (b - \sqrt{b})/a$ we have an orthogonal matrix $U = U(x, y)$, where:

$$x = -\frac{t}{\sqrt{b}(t + 1)}, \quad y = \frac{1}{\sqrt{b}(t + 1)}$$

In order to compute these variables, we use the following formula:

$$(a + b)^2 - b = a^2 + b^2 + 2ab - b = a^2 + 2ab + ac = Na$$

This gives indeed the formula of y in the statement:

$$y = \frac{a}{(a + b)\sqrt{b} - b} = \frac{(a + b)\sqrt{b} + b}{Nb}$$

As for the formula of x , we can obtain it as follows:

$$x = -ty = \frac{(\sqrt{b} - b)((a + b)\sqrt{b} + b)}{Nab} = \frac{a + (1 - a - b)\sqrt{b}}{Na}$$

Let us compute now the quantity appearing in Lemma 3.6. We have:

$$\begin{aligned}
N(a-b) + 2b &= (a + 2b + c)(a-b) + 2b \\
&= a^2 + ab - 2b^2 + ac - bc + 2b \\
&= a^2 + ab - ac - bc \\
&= (a-c)(a+b)
\end{aligned}$$

Similarly, $N(c-b) + 2b = (c-a)(c+b)$, so the quantity in Lemma 3.6 is Ky , with:

$$\begin{aligned}
K &= (a-c)(a+b)t + (c-a)(c+b) \\
&= (c-a)(c+b - (a+b)t) \\
&= \frac{c-a}{a}(ac + ab - (a+b)(b - \sqrt{b})) \\
&= \frac{c-a}{a}((ac - b^2) + (a+b)\sqrt{b}) \\
&= \frac{c-a}{a}((a+b)\sqrt{b} - b)
\end{aligned}$$

Since this quantity is positive, Lemma 3.6 applies and gives the result. \square

Theorem 3.8. *Assume that $q = p^k$ is a prime power. Then the matrix $I_N \in M_N(x, y)$, where $N = q^2 + q + 1$ and*

$$x = \frac{1 - q\sqrt{q}}{\sqrt{N}}, \quad y = \frac{q + (q+1)\sqrt{q}}{q\sqrt{N}}$$

having $(1, q, q^2 - q)$ pattern coming from the point-line incidence of the projective plane over \mathbb{F}_q is an almost Hadamard matrix.

Proof. Indeed, the conditions $c \geq a$ and $b(b-1) = ac$ needed in Proposition 3.7 are satisfied, and the variables constructed there are $x' = x/\sqrt{N}$ and $y' = y/\sqrt{N}$. \square

There are of course many other interesting examples of two-entry almost Hadamard matrices, all worth investigating in detail, but we will stop here. Indeed, the main purpose of the remainder of this paper is to provide a list of almost Hadamard matrices which are “as optimal as possible”, at $N = 2, 3, \dots, 13$, and in order to establish this list, we will just need the incidence matrices I_N , plus the matrix coming from the Paley biplane.

Let us mention however two more important aspects of the general theory:

- (1) The series I_N is the particular case of a 2-parameter series $I_N^{(d)}$. Indeed, associated to $q = p^k$ and $d \in \mathbb{N}$ is a certain $([d+2]_q, [d+1]_q, [d]_q)$ block design coming from \mathbb{F}_q , where $[e]_q = (q^e - 1)/(q - 1)$, cf. [14], [35]. Thus by Proposition 3.7 we obtain an almost Hadamard matrix $I_N^{(d)}$, having pattern $(\frac{q^d-1}{q-1}, q^d, q^d(q-1))$.
- (2) Trying to find block designs and patterns leads to the following chess problem: consider a $N \times N$ chessboard, take NM rooks with $M \leq N/2$, fix an integer

$K \leq M$, and try to place all the rooks on the board such that: (a) there are exactly M rooks on each row and each column of the board, and (b) for any pair of rows or columns, there are exactly K pairs of mutually attacking rooks.

Finally, let us mention that there are many questions raised by the almost Hadamard matrices, at the quantum algebraic level. For instance the symmetries of Hadamard matrices are known to be described by quantum permutations [3], and it would be interesting to have a similar result for the almost Hadamard matrices. This might probably bring some new ideas on the “homogeneous implies quantum homogeneous” question, raised in [2], and having connections with the finite projective planes. Also, an interesting link between mutually unbiased bases, complex Hadamard matrices and affine planes was emphasized in [9], but its relation with our present investigations is not known yet.

4. LIST OF EXAMPLES

In this section we present a list of examples of almost Hadamard matrices, for small values of N . Since we are mainly interested in the optimal case, our examples will be chosen to be “as optimal as possible”, i.e. will be chosen as to have big 1-norms.

So, let us first compute the 1-norms for the examples that we have. In what follows L_N is the matrix found in Theorem 2.6, and I_N is the matrix found in Theorem 3.8.

Theorem 4.1. *The 1-norms of the basic examples of almost Hadamard matrices are:*

- (1) *Hadamard case:* $\|H/\sqrt{N}\|_1 = N\sqrt{N}$, for any $H \in M_N(\pm 1)$ Hadamard.
- (2) *Basic series case:* $\|K_N/\sqrt{N}\|_1 = 3N - 4$, where $K_N = \sqrt{N}(2J_N - 1_N)$.
- (3) *Circulant series case:* $\|L_N/\sqrt{N}\|_1 = \frac{2}{\pi}N \log N + O(N)$.
- (4) *Incidence series case:* $\|I_N/\sqrt{N}\|_1 = (q^2 - q - 1) + 2q(q + 1)\sqrt{q}$.

Proof. The first two assertions are clear. For the third one, with $N = 2n + 1$ we have:

$$\|L_N/\sqrt{N}\|_1 = 2 \sum_{i=0}^n \cos^{-1} \frac{i\pi}{N} + O(N) = 2 \sum_{i=0}^n \sin^{-1} \frac{(2i+1)\pi}{2N} + O(N)$$

Now by using $\sin x \sim x$ and $\sum_{i=1}^k 1/i = \log k + O(1)$ we obtain, as claimed:

$$\|L_N/\sqrt{N}\|_1 = \frac{4N}{\pi} \sum_{i=0}^n \frac{1}{2i+1} + O(N) = \frac{2N}{\pi} \log N + O(N)$$

As for the last assertion, let first U be the matrix in Lemma 3.6. We have:

$$\|U\|_1 = N((a+b)|x| + (b+c)|y|)$$

In the particular case of the orthogonal matrices in Proposition 3.7, we get:

$$\|U\|_1 = (c-a) + \frac{(a+b)(2a+2b-2)}{a} \sqrt{b}$$

Now with $a = 1$, $b = q$, $c = q^2 - q$, this gives the formula in the statement. \square

Observe that at N big the above matrices K_N, L_N, I_N are far from being optimal. In fact, with $N \rightarrow \infty$, the corresponding orthogonal matrices don't even match the average of the 1-norm on $O(N)$, which, according to [5], is $\sim cN\sqrt{N}$, with $c = 0.797..$

With these ingredients in hand, let us discuss now the various examples:

At $N = 2$ we have the Walsh matrix H_2 , which is of course optimal.

At $N = 3$ we have the almost Hadamard matrix K_3 , which is optimal:

$$K_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}$$

At $N = 4$ we have the Hadamard matrix $H_4 \sim K_4$, once again optimal.

At $N = 5$ we have the matrix K_5 , that we believe to be optimal as well:

$$K_5 = \frac{1}{\sqrt{5}} \begin{pmatrix} -3 & 2 & 2 & 2 & 2 \\ 2 & -3 & 2 & 2 & 2 \\ 2 & 2 & -3 & 2 & 2 \\ 2 & 2 & 2 & -3 & 2 \\ 2 & 2 & 2 & 2 & -3 \end{pmatrix}$$

At $N = 6$ it is plausible that the optimal AHM is simply $K_3 \otimes H_2$:

$$K_3 \otimes H_2 = \frac{1}{\sqrt{3}} \begin{pmatrix} -1 & 2 & 2 & -1 & 2 & 2 \\ 2 & -1 & 2 & 2 & -1 & 2 \\ 2 & 2 & -1 & 2 & 2 & -1 \\ -1 & 2 & 2 & 1 & -2 & -2 \\ 2 & -1 & 2 & -2 & 1 & -2 \\ 2 & 2 & -1 & -2 & -2 & 1 \end{pmatrix}$$

At $N = 7$ we have the incidence matrix of the Fano plane ($x = 2 - 4\sqrt{2}, y = 2 + 3\sqrt{2}$):

$$I_7 = \frac{1}{2\sqrt{7}} \begin{pmatrix} x & x & y & y & y & x & y \\ y & x & x & y & y & y & x \\ x & y & x & x & y & y & y \\ y & x & y & x & x & y & y \\ y & y & x & y & x & x & y \\ y & y & y & x & y & x & x \\ x & y & y & y & x & y & x \end{pmatrix}$$

At $N = 8$ we have the third Walsh matrix $H_8 = H_2 \otimes H_4$, of course optimal.

At $N = 9$ we just have the matrix $K_3 \otimes K_3$, which can be shown not to be optimal.

At $N = 10$ we believe that the matrix $K_5 \otimes H_2$ is optimal:

$$K_5 \otimes H_2 = \frac{1}{\sqrt{5}} \begin{pmatrix} -3 & 2 & 2 & 2 & 2 & -3 & 2 & 2 & 2 & 2 \\ 2 & -3 & 2 & 2 & 2 & 2 & -3 & 2 & 2 & 2 \\ 2 & 2 & -3 & 2 & 2 & 2 & 2 & -3 & 2 & 2 \\ 2 & 2 & 2 & -3 & 2 & 2 & 2 & 2 & -3 & 2 \\ 2 & 2 & 2 & 2 & -3 & 2 & 2 & 2 & 2 & -3 \\ -3 & 2 & 2 & 2 & 2 & 3 & -2 & -2 & -2 & -2 \\ 2 & -3 & 2 & 2 & 2 & -2 & 3 & -2 & -2 & -2 \\ 2 & 2 & -3 & 2 & 2 & -2 & -2 & 3 & -2 & -2 \\ 2 & 2 & 2 & -3 & 2 & -2 & -2 & -2 & 3 & -2 \\ 2 & 2 & 2 & 2 & -3 & -2 & -2 & -2 & -2 & 3 \end{pmatrix}$$

At $N = 11$ we have the matrix of the Paley biplane ($x = 6 - 12\sqrt{3}$, $y = 6 + 10\sqrt{3}$):

$$P_{11} = \frac{1}{6\sqrt{11}} \begin{pmatrix} y & x & y & x & x & x & y & y & y & x & y \\ y & y & x & y & x & x & x & y & y & y & x \\ x & y & y & x & y & x & x & x & y & y & y \\ y & x & y & y & x & y & x & x & x & y & y \\ y & y & x & y & y & x & y & x & x & x & y \\ y & y & y & x & y & y & x & y & x & x & x \\ x & y & y & y & x & y & y & x & y & x & x \\ x & x & y & y & y & x & y & y & x & y & x \\ x & x & x & y & y & y & x & y & y & x & y \\ y & x & x & x & y & y & y & x & y & y & x \\ x & y & x & x & x & y & y & y & x & y & y \end{pmatrix}$$

At $N = 12$ we have the Sylvester Hadamard matrix S_{12} , of course optimal.

At $N = 13$ we have the incidence matrix of $P(\mathbb{F}_3)$ ($x = 3 - 9\sqrt{3}$, $y = 3 + 4\sqrt{3}$):

$$I_{13} = \frac{1}{3\sqrt{13}} \begin{pmatrix} x & x & x & x & y & y & y & y & y & y & y & y & y \\ x & y & y & y & x & x & x & y & y & y & y & y & y \\ x & y & y & y & y & y & y & x & x & x & y & y & y \\ x & y & y & y & y & y & y & y & y & y & x & x & x \\ y & x & y & y & y & y & x & y & x & y & y & y & x \\ y & x & y & y & y & x & y & y & y & x & x & y & y \\ y & x & y & y & x & y & y & x & y & y & y & x & y \\ y & y & x & y & y & y & x & x & y & y & x & y & y \\ y & y & x & y & x & y & y & y & y & x & y & y & x \\ y & y & x & y & y & x & y & y & x & y & y & x & y \\ y & y & y & x & y & x & y & x & y & y & y & y & x \\ y & y & y & x & y & y & x & y & y & x & y & x & y \\ y & y & y & x & x & y & y & y & x & y & x & y & y \end{pmatrix}$$

N	matrix	1-norm (formula)	1-norm (numeric)	$N\sqrt{N}$	remarks
2	H_2	$2\sqrt{2}$	2.828	2.828	Hadamard
3	K_3	5	5.000	5.196	optimal
4	K_4	8	8.000	8.000	Hadamard
5	K_5	11	11.000	11.118	
6	$K_3 \otimes H_2$	$10\sqrt{28}$	14.142	14.697	
7	I_7	$1 + 12\sqrt{2}$	17.971	18.520	
8	H_8	$16\sqrt{2}$	22.627	22.627	Hadamard
9	?	–	> 26.513	27.000	
10	$K_5 \otimes H_2$	$22\sqrt{2}$	31.113	31.623	
11	P_{11}	$1 + 20\sqrt{3}$	35.641	36.483	
12	S_{12}	$24\sqrt{3}$	41.569	41.569	Hadamard
13	I_{13}	$5 + 24\sqrt{3}$	46.569	46.872	

TABLE 1. Almost Hadamard matrices $H \in M_N(\mathbb{R})$, chosen as for the corresponding orthogonal matrices $U = H/\sqrt{N}$ to have big 1-norm. All matrices are believed to be optimal. The lower bound for the maximum of the 1-norm on $O(9)$, which is not optimal, was obtained by numerical simulation.

The norms of the above matrices can be computed by using the various formulae in Theorem 4.1 and its proof, and the results are summarized in Table 1.

5. CONCLUSION

We have seen in this paper that the Hadamard matrices are quite nicely generalized by the almost Hadamard matrices (AHM), which exist at any given order $N \in \mathbb{N}$. Our study of these matrices, which was for the most of algebraic nature, turns to be related to several interesting combinatorial problems, notably to the Circulant Hadamard Conjecture.

We believe that the AHM can be used as well in connection with several problems in quantum physics, in a way somehow similar to the way the complex Hadamard matrices (CHM) are used. Indeed, since the CHM exist as well at any given order $N \in \mathbb{N}$, these matrices proved to be useful in several branches of quantum physics. For instance in quantum optics they are sometimes called the “Zeilinger matrices”, as they can be applied to design symmetric linear multiports, used to split the beam into N parts of the same intensity and to analyze interference effect [33], [22]. In the theory of quantum information one uses quantum Hadamard matrices to construct mutually unbiased bases (MUB) [21], [16] to design teleportation and dense coding schemes. As shown in the seminal work of Werner [38] these two problems are in fact equivalent and also equivalent to construction of unitary depolarisers and maximally entangled bases [39].

Although quantum mechanics in a natural way relays on a complex Hilbert space it is often convenient to study a simplified problem and restrict attention to the subset of real quantum states only. Such an approach can be useful in theoretical investigations of quantum entanglement [13] or also in experimental studies on engineering of quantum states, as creating of a real state by an orthogonal rotation usually requires less effort than construction of an arbitrary complex state.

There exists therefore a natural motivation to ask similar problems concerning e.g. unbiased bases and teleportation schemes in the real setup. For instance, it is known that for any N there exist $\leq N/2 + 1$ real MUB and for most dimensions their actual number is not larger than 3, cf. [10], while for any prime N there exist $N + 1$ complex MUB. Note that the real MUB and Hadamard matrices are closely related to several combinatorial problems [19], [29], [30]. In the case the maximal number of real MUB does not exist one can search for an optimal set of real bases which are approximately unbiased.

Let us now return to the construction of symmetric multiports which relay on the complex Hadamard matrices [33], [22]. In the case of a k -qubit system there exist Hadamard matrices of order $N = 2^k$, so one may use real orthogonal matrices for this purpose. However, already for $N = 3, 5, 6, 7$ real Hadamard matrices do not exist, so in these dimensions, if one restricts the rotations to orthogonal matrices, there are no real symmetric multiports. Therefore, for these dimensions one may always raise the following question: what is the optimal, approximate solution of the problem, if one is allowed to use only real states and orthogonal matrices? The almost Hadamard matrices analyzed in this paper are directly applicable for such a class of problems.

REFERENCES

- [1] A. Álvarez, J.A. Armario, M.D. Frau, E. Martin and A. Osuna, Error correcting codes from quasi-Hadamard matrices, *Lecture Notes in Comput. Sci.* **4547** (2007), 294–302.
- [2] T. Banica, Quantum automorphism groups of homogeneous graphs, *J. Funct. Anal.* **224** (2005), 243–280.
- [3] T. Banica, Quantum permutations, Hadamard matrices, and the search for matrix models, *Banach Center Publ.*, to appear.
- [4] T. Banica, J. Bichon and J.-M. Schlenker, Representations of quantum permutation algebras, *J. Funct. Anal.* **257** (2009), 2864–2910.
- [5] T. Banica, B. Collins and J.-M. Schlenker, On orthogonal matrices maximizing the 1-norm, *Indiana Univ. Math. J.* **59** (2010), 839–856.
- [6] T. Banica, B. Collins and J.-M. Schlenker, On polynomial integrals over the orthogonal group, *J. Combin. Theory Ser. A* **118** (2011), 778–795.
- [7] T. Banica and J.-M. Schlenker, Combinatorial aspects of orthogonal group integrals, *Internat. J. Math.* **22** (2011), 1611–1646.
- [8] I. Bengtsson, W. Bruzda, Å. Ericsson, J.-Å. Larsson, W. Tadej and K. Życzkowski, Mutually unbiased bases and Hadamard matrices of order six, *J. Math. Phys.* **48** (2007), 1–33.
- [9] I. Bengtsson and A. Ericsson, Mutually unbiased bases and the complementarity polytope, *Open Syst. Inf. Dyn.* **12** (2005), 107–120.

- [10] P.O. Boykin, M. Sitharam, M. Tarifi and P. Wocjan, Real mutually unbiased bases, [arxiv:quant-ph/0502024](#).
- [11] E. Brown, The fabulous (11,5,2) biplane, *Math. Mag.* **77** (2004), 87–100.
- [12] A.T. Butson, Generalized Hadamard matrices, *Proc. Amer. Math. Soc.* **13** (1962), 894–898.
- [13] C.M. Caves, C.A. Fuchs and P. Rungta, Entanglement of formation of an arbitrary state of two rebits, *Found. Phys. Lett.* **14** (2001), 199.
- [14] C.J. Colbourn and J.H. Dinitz, Handbook of combinatorial designs, CRC Press (2007).
- [15] B. Collins and P. Śniady, Integration with respect to the Haar measure on unitary, orthogonal and symplectic groups, *Comm. Math. Phys.* **264** (2006), 773–795.
- [16] T. Durt, B.-G. Englert, I. Bengtsson and K. Życzkowski, On mutually unbiased bases, *Int. J. Quantum Information* **8** (2010), 535–640.
- [17] U. Haagerup, Orthogonal maximal abelian *-subalgebras of the $n \times n$ matrices, in “Operator algebras and quantum field theory”, International Press (1997), 296–323.
- [18] J. Hadamard, Résolution d’une question relative aux déterminants, *Bull. Sci. Math.* **2** (1893), 240–246.
- [19] W.H. Holzman, H. Kharaghani and W. Orrick, On real unbiased Hadamard matrices, *Contemp. Math.* **531** (2010), 243–250.
- [20] K.J. Horadam, Hadamard matrices and their applications, Princeton Univ. Press (2007).
- [21] I.D. Ivanović, Geometrical description of quantal state determination, *J. Phys. A* **14** (1981), 3241–3245.
- [22] I. Jex, S. Stenholm and A. Zeilinger, Hamiltonian theory of a symmetric multiphoton, *Opt. Commun.* **117** (1995), 95–101.
- [23] V.F.R. Jones, On knot invariants related to some statistical mechanical models, *Pacific J. Math.* **137** (1989), 311–334.
- [24] H. Kharaghani and B. Tayfeh-Rezaie, A Hadamard matrix of order 428, *J. Combin. Des.* **13** (2005), 435–440.
- [25] J. Konvalina and R.H. Kosloski, Cyclic quasi-Hadamard matrices, *Utilitas Math.* **12** (1977), 273–287.
- [26] T.Y. Lam and K.H. Leung, On vanishing sums of roots of unity, *J. Algebra* **224** (2000), 91–109.
- [27] W. de Launey, On the non-existence of generalized weighing matrices, *Ars Combin.* **17** (1984), 117–132.
- [28] W. de Launey and D.M. Gordon, A comment on the Hadamard conjecture, *J. Combin. Theory Ser. A* **95** (2001), 180–184.
- [29] N. LeCompte, W.J. Martin and W. Owens, On the equivalence between real mutually unbiased bases and certain class of association schemes, *European J. Combin.* **31** (2010), 1499–1512.
- [30] M. Matolcsi, I.Z. Ruzsa and M. Weiner, Real and complex unbiased Hadamard matrices, [arxiv:1201.0631](#).
- [31] K.-H. Park and H.-Y. Song, Quasi-Hadamard matrices, *Proc. ISIT 2010, Austin, TX* (2010).
- [32] S. Popa, Orthogonal pairs of *-subalgebras in finite von Neumann algebras, *J. Operator Theory* **9** (1983), 253–268.
- [33] M. Reck, A. Zeilinger, H.J. Bernstein and P. Bertani, Experimental realization of any discrete unitary operator, *Phys. Rev. Lett.* **73** (1994), 58–61.
- [34] H.J. Ryser, Combinatorial mathematics, Wiley (1963).
- [35] D.R. Stinson, Combinatorial designs: constructions and analysis, Springer-Verlag (2006).
- [36] J.J. Sylvester, Thoughts on inverse orthogonal matrices, simultaneous sign-successions, and tessellated pavements in two or more colours, with applications to Newton’s rule, ornamental tile-work, and the theory of numbers, *Phil. Mag.* **34** (1867), 461–475.

- [37] W. Tadej and K. Życzkowski, A concise guide to complex Hadamard matrices, *Open Syst. Inf. Dyn.* **13** (2006), 133–177.
- [38] R.F. Werner, All teleportation and dense coding schemes, *J. Phys. A* **34** (2001), 7081–7094.
- [39] A. Wójcik, A. Grudka and R.W. Chhajlany, Generation of inequivalent generalized Bell bases, *Quantum Inf. Process.* **2** (2003), 201–206.

T.B.: DEPARTMENT OF MATHEMATICS, CERGY-PONTOISE UNIVERSITY, 95000 CERGY-PONTOISE, FRANCE. teodor.banica@u-cergy.fr

I.N.: CNRS, LABORATOIRE DE PHYSIQUE THÉORIQUE, IRSAMC, UNIVERSITÉ DE TOULOUSE, UPS, 31062 TOULOUSE, FRANCE. nechita@irsamc.ups-tlse.fr

K.Z.: INSTITUTE OF PHYSICS, JAGIELLONIAN UNIVERSITY, CRACOW AND CENTER FOR THEORETICAL PHYSICS, POLISH ACADEMY OF SCIENCES, WARSAW, POLAND. karol@tatrzy.if.uj.edu.pl