

**Table S1.** Biochemical parameters of the participants (n=155)

	OST		Rehabilitation		p-value	Total	
	Mean	SD	Mean	SD		Mean	SD
<b>FBS (mg/dl)</b>	94.05	16.81	85.53	7.54	<b>&lt;0.001</b>	89.95	13.83
<b>HDL-C (mg/dl)</b>	43.06	11.85	43.81	12.89	0.708	43.42	12.33
<b>TG (mg/dl)</b>	119.46	69.67	133.81	156.89	0.459	126.37	119.59
<b>Total Cholesterol (mg/dl)</b>	188.94	53.36	188.07	38.06	0.908	188.52	46.49
<b>LDL-C (mg/dl)</b>	115.48	44.41	114.07	31.42	0.822	114.80	38.60

SD: Standard deviation; OST: Opioid substitution therapy; FBS: Fasting blood sugar; HDL-C: High-density lipoprotein-cholesterol; TG: Triglycerides; LDL-C: Low-density lipoprotein-cholesterol.

**Table S2.** Anthropometric measurements of the participants (n=155)

	OST		Rehabilitation		p-value	Total	
	Mean	SD	Mean	SD		Mean	SD
<b>Waist circumference (cm)</b>	91.16	13.84	94.42	12.56	0.128	92.75	13.29
<b>Systolic blood pressure (mmHg)</b>	123.44	12.21	127.85	15.31	<b>0.048</b>	125.6	13.93
<b>Diastolic blood pressure (mmHg)</b>	75.00	11.47	78.99	15.94	0.075	76.93	13.91
<b>BMI (Kg/m<sup>2</sup>)</b>	26.97	5.37	27.57	4.62	0.453	27.41	4.99

SD: Standard deviation; OST: Opioid substitution therapy; BMI: Body mass index.