



Search for the lepton-flavour violating decays $B^+ \rightarrow K^+ \mu^\pm e^\mp$

LHCb collaboration[†]

Abstract

A search for the lepton-flavour violating decays $B^+ \rightarrow K^+ \mu^\pm e^\mp$ is performed using a sample of proton-proton collision data, collected with the LHCb experiment at centre-of-mass energies of 7 and 8 TeV and corresponding to an integrated luminosity of 3 fb^{-1} . No significant signal is observed, and upper limits on the branching fractions are set as $\mathcal{B}(B^+ \rightarrow K^+ \mu^- e^+) < 7.0 (9.5) \times 10^{-9}$ and $\mathcal{B}(B^+ \rightarrow K^+ \mu^+ e^-) < 6.4 (8.8) \times 10^{-9}$ at 90 (95)% confidence level. The results improve the current best limits on these decays by more than one order of magnitude.

Submitted to Phys. Rev. Lett.

© 2019 CERN for the benefit of the LHCb collaboration. CC-BY-4.0 licence.

[†]Authors are listed at the end of this paper.

The observation of neutrino oscillations has provided the first evidence for lepton-flavour violation (LFV) in neutral leptons. By contrast, LFV in the charged sector is negligible in the Standard Model (SM) [1] and any observation of a charged LFV decay would be indisputable evidence for physics beyond the SM (BSM). In light of recent flavour anomalies in semileptonic $b \rightarrow s \ell^+ \ell^-$ transitions [2–4], many SM extensions have been proposed that link lepton-universality violation to LFV, predicting in particular a significantly enhanced decay rate in $b \rightarrow s \mu^\mp e^\pm$ processes. In this Letter a search for the decays of $B^+ \rightarrow K^+ \mu^\pm e^\mp$ is reported (Inclusion of charge-conjugate processes is implied throughout the letter). Their branching fractions are predicted to be in the range $10^{-8} - 10^{-10}$ in leptoquark models [5, 6], extended gauge boson models [7], or models including CP violation in the neutrino sector [8]. Currently, the best limits of $\mathcal{B}(B^+ \rightarrow K^+ \mu^- e^+) < 9.1 \times 10^{-8}$ and $\mathcal{B}(B^+ \rightarrow K^+ \mu^+ e^-) < 13 \times 10^{-8}$ have been set by the BaBar collaboration at the 90% confidence level [9].

A data set of proton-proton (pp) collisions corresponding to an integrated luminosity of 3 fb^{-1} , recorded with the LHCb detector in 2011 and 2012 at centre-of-mass energies of 7 TeV and 8 TeV, respectively, is used in this analysis. The two final states with different lepton charge combinations are studied independently, since they could be affected differently by BSM dynamics. The yields of the $B^+ \rightarrow K^+ \mu^\pm e^\mp$ decays are normalised to those of the $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ decay, which has a well-known branching fraction [10], the same topology, and similar signatures in the detector. The $B^+ \rightarrow K^+ J/\psi (\rightarrow e^+ e^-)$ decay is also used as a control channel in the analysis.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [11, 12]. The detector includes a silicon-strip vertex detector surrounding the pp interaction region, tracking stations located either side of a dipole magnet, ring-imaging Cherenkov (RICH) detectors, calorimeters and muon chambers.

The online event selection is performed by a trigger [13], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. At the hardware trigger stage, $B^+ \rightarrow K^+ \mu^\pm e^\mp$ and $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ event candidates are required to have a muon with high transverse momentum, p_T . In the subsequent software trigger, at least one charged particle must have a $p_T > 1.7 \text{ GeV}/c$ in the 2011 data set and $p_T > 1.6 \text{ GeV}/c$ in 2012, unless the particle is identified as a muon in which case $p_T > 1.0 \text{ GeV}/c$ is required. This track must be significantly displaced from any primary interaction vertex (PV) in the event. Finally, a two- or three-track secondary vertex with a significant displacement from any PV is required, where a multivariate algorithm [14] is used for the identification of secondary vertices consistent with the weak decay of a b hadron.

Simulated samples are used to evaluate signal efficiencies, to train multivariate classifiers, to model the shape of the invariant mass of the $B^+ \rightarrow K^+ \mu^\pm e^\mp$ signal candidates, and to study physics backgrounds. In the simulation, pp collisions are generated using PYTHIA [15, 16] with a specific LHCb configuration [17]. Decays of unstable particles are described by EVTGEN [18], in which final-state radiation is generated using PHOTOS [19]. A phase-space model is adopted for signal $B^+ \rightarrow K^+ \mu^\pm e^\mp$ decays. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [20] as described in Ref. [21].

The $B^+ \rightarrow K^+ \mu^\pm e^\mp$ candidates passing the trigger selection are reconstructed by combining three charged tracks originating from a good-quality common vertex. The

tracks forming the B^+ candidate are required not to originate from any PV and must have sizeable transverse momentum. Due to the long lifetime of the B^+ meson, this vertex is required to be well separated from any PV. The B^+ direction vector, determined from its production and decay vertex positions, must be aligned with its momentum vector. The mass of the reconstructed B^+ candidate, $m(K^+\mu^\pm e^\mp)$, is restricted to lie within $\pm 1500 \text{ MeV}/c^2$ of the known B^+ meson mass [10]. Furthermore, the B -meson decay products must be well identified as a kaon, an electron and a muon, exploiting information from the Cherenkov detectors, the calorimeters, and the muon stations. The electron candidate kinematics are corrected for bremsstrahlung photon emission if a compatible photon candidate in the calorimeter is found. Kaon and electron candidates that have hits in the muon stations consistent with their trajectories are rejected. The same selection is applied to the normalisation (control) channels, for which the dimuon (dielectron) invariant mass is additionally required to be consistent with the known J/ψ mass [10]. The selection and analysis procedures were developed without inspecting the signal data in the region $m(K^+\mu^\pm e^\mp) \in [4985, 5385] \text{ MeV}/c^2$.

The most significant backgrounds originate from partially reconstructed B^+ decays, *e.g.* from double semileptonic $B^+ \rightarrow \bar{D}^0 X \ell^+ \nu_\ell$ with $\bar{D}^0 \rightarrow K^+ Y \ell'^- \bar{\nu}_{\ell'}$ decays, where X and Y represent hadrons, while ℓ and ℓ' are leptons. They are removed by imposing the requirement $m(K^+\ell^-) > 1885 \text{ MeV}/c^2$. Contributions from decays involving charmonium resonances, where one lepton is misidentified as a kaon or as a lepton of a different flavour, are rejected by mass vetoes.

The combinatorial background, which consists of random tracks that are associated to a common vertex, is reduced using a boosted decision tree (BDT) [22, 23] algorithm. This BDT combines information about the B^+ meson kinematics and information related to its flight distance, decay vertex quality and impact parameter with respect to the primary vertex. In addition, it uses information such as the impact parameters of the electron, muon and kaon candidates, and the isolation of the B^+ candidate from any other charged track in the event [24].

The BDT is trained on simulated $B^+ \rightarrow K^+\mu^\pm e^\mp$ events that have satisfied the previous requirements. The simulated samples are adjusted using $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ and $B^+ \rightarrow K^+ J/\psi (\rightarrow e^+ e^-)$ decays in data to correct data-simulation differences in the B -meson production kinematics, vertex quality, and detector occupancy represented by the number of tracks in the detector. The upper-mass sideband, corresponding to $m(K^+\mu^\pm e^\mp) \in [5385, 6000] \text{ MeV}/c^2$, is used as a proxy for the background. The training is performed using a k -folding approach [25] with ten folds, which allows the whole sample to be used without biasing the output of the classifier. The optimal requirement on the BDT classifier is chosen to give the best expected upper limits on the branching fractions $\mathcal{B}(B^+ \rightarrow K^+\mu^\pm e^\mp)$.

The candidates surviving this multivariate selection are used to train a second BDT, dedicated to reject background from partially reconstructed b -hadron decays. The background sample for the training is taken from the lower-mass sideband in data, corresponding to $m(K^+\mu^\pm e^\mp) \in [4550, 4985] \text{ MeV}/c^2$, where the partially reconstructed background is expected to contribute. The signal proxy is the same as for the first BDT. The training procedure shares the k -folding approach and the same set of discriminating variables used to construct the first multivariate discriminant, with the addition of the ratio between the projections of the electron and the $K^+\mu^\pm$ momenta orthogonal to the B -meson direction of flight. The requirements on the second BDT are optimized in the same manner as the first

BDT. The final stage of the selection, where requirements on the particle identification (PID) variables based on a neural net classifier for the kaon, electron and muon are applied [26], ensures the suppression of candidates from decays with misidentification of at least one particle.

The performance of the PID algorithms is not perfectly simulated, and thus a correction is performed using high-purity calibration data samples of muons from $B \rightarrow XJ/\psi (\rightarrow \mu^+\mu^-)$ decays, electrons from $B^+ \rightarrow K^+J/\psi (\rightarrow e^+e^-)$ decays, and kaons from $D^{*+} \rightarrow D^0 (\rightarrow K^-\pi^+)\pi^+$ decays [27]. The calibration data are binned in the particle's momentum and pseudorapidity, and in the detector occupancy. Particle identification variables for the simulated data sets are sampled from the distributions of calibration data in the corresponding bin. The performance of the PID resampling is validated on both the $B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-)$ and $B^+ \rightarrow K^+J/\psi (\rightarrow e^+e^-)$ control channels.

The potential contamination from b -hadron decays in the signal mass region after selection is analysed using dedicated simulated samples. Two categories are analysed: fully reconstructed B decays, with at least one particle in the final state misidentified, such as the semileptonic decays $B^+ \rightarrow K^+\ell^+\ell^-$ and $B^+ \rightarrow K^+J/\psi (\rightarrow \ell^+\ell^-)$, or fully hadronic B^+ decays as $B^+ \rightarrow K^+\pi^+\pi^-$; partially reconstructed decays in which at least one particle is not reconstructed and one or more particles are misidentified in addition, such as $B^0 \rightarrow K^{*0}\ell^+\ell^-$, $\Lambda_b^0 \rightarrow pK^-\ell^+\ell^-$, $\Lambda_b^0 \rightarrow pK^-J/\psi (\rightarrow \ell^+\ell^-)$ and $B^+ \rightarrow \bar{D}^0\ell^+\nu_\ell$ transitions, where the \bar{D}^0 meson decays further to $K^+\pi^-$ or $K^+\ell^-\bar{\nu}_\ell$. The expected number of candidates from each possible background source after the selection is evaluated from simulation and is found to be negligible.

The branching fraction $\mathcal{B}(B^+ \rightarrow K^+\mu^\pm e^\mp)$ is measured relative to the normalisation channel using

$$\mathcal{B}(B^+ \rightarrow K^+\mu^\pm e^\mp) = N(B^+ \rightarrow K^+\mu^\pm e^\mp) \times \alpha, \quad (1)$$

$$\alpha \equiv \frac{\mathcal{B}(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-)) \varepsilon(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-))}{\varepsilon(B^+ \rightarrow K^+\mu^\pm e^\mp) N(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-))},$$

where the $\varepsilon(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-))$ and $\varepsilon(B^+ \rightarrow K^+\mu^\pm e^\mp)$ denote the efficiencies of the normalisation and signal channels, respectively; $N(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-))$ and $N(B^+ \rightarrow K^+\mu^\pm e^\mp)$ are the observed $B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-)$ and $B^+ \rightarrow K^+\mu^\pm e^\mp$ yields, respectively. The value of the branching fraction of the normalisation mode is $\mathcal{B}(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-)) = (6.02 \pm 0.17) \times 10^{-5}$, taken from Ref. [10]. The yield of the normalisation channel is determined from an unbinned extended maximum-likelihood fit to the invariant mass $m(K^+\mu^+\mu^-)$ of the selected $B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-)$ candidates, performed separately on 2011 and 2012 data. The sum of two Crystal Ball functions [28] is used to parameterise the signal, while an exponential function models the background. The yields resulting from the fits are 26940 ± 170 for 2011 and 59220 ± 250 for 2012 data.

The efficiencies are calculated taking into account all selection requirements. The analysis is performed assuming a phase-space model for the signal decay. Efficiency maps in bins of the invariant masses of the particles in the final state $m_{K^+e^\pm}^2$ and $m_{K^+\mu^\mp}^2$ are provided in Fig 1 to allow for the interpretation of the result in different BSM scenarios.

All efficiencies are determined from calibrated simulation samples and the normalisation factors for the two decay channels are given in Table 1. The two data taking periods are combined into a single normalisation factor taking into account the relative data sizes and efficiencies. The ratio $\alpha/\mathcal{B}(B^+ \rightarrow K^+J/\psi (\rightarrow \mu^+\mu^-))$, which excludes external inputs, is also quoted.




Figure 1: Efficiency of (left) $B^+ \rightarrow K^+ \mu^- e^+$ and (right) $B^+ \rightarrow K^+ \mu^+ e^-$ as function of the squared invariant masses $m_{K^+ e^\pm}^2$ and $m_{K^+ \mu^\mp}^2$. The variation of efficiency across the Dalitz plane is due to applied vetoes. The efficiencies are given in per mille.

Table 1: Normalisation factor α for $B^+ \rightarrow K^+ \mu^- e^+$ and $B^+ \rightarrow K^+ \mu^+ e^-$ final states. The ratio $\alpha/\mathcal{B}(B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-))$ is independent of external inputs.

Decay	α	$\alpha/\mathcal{B}(B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-))$
$B^+ \rightarrow K^+ \mu^- e^+$	$(1.97 \pm 0.14) \times 10^{-9}$	$(3.27 \pm 0.22) \times 10^{-5}$
$B^+ \rightarrow K^+ \mu^+ e^-$	$(2.21 \pm 0.14) \times 10^{-9}$	$(3.68 \pm 0.22) \times 10^{-5}$

The invariant-mass distribution of $B^+ \rightarrow K^+ \mu^\pm e^\mp$ candidates is modeled differently depending on whether bremsstrahlung photons have been included in the momentum calculation for the electrons. The sum of two Crystal Ball functions with common mean value is used in both cases. In the case of bremsstrahlung the tails are on opposite sides of the peak. Otherwise, the two tails share the same parameters. Their values, obtained from the $B^+ \rightarrow K^+ \mu^\pm e^\mp$ simulation, are corrected taking into account the differences between data and simulation in $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ and $B^+ \rightarrow K^+ J/\psi (\rightarrow e^+ e^-)$ decays. Two types of unbinned maximum-likelihood fits are performed on the dataset. The first fit assumes only background is present, with the background modeled with an exponential function. From this fit 3.9 ± 1.1 and 0.9 ± 0.6 background candidates are expected in the signal mass window for the $B^+ \rightarrow K^+ \mu^+ e^-$ and $B^+ \rightarrow K^+ \mu^- e^+$ modes, respectively. The second fit includes the signal component, which is used to determine the signal yields. The $B^+ \rightarrow K^+ \mu^- e^+$ and $B^+ \rightarrow K^+ \mu^+ e^-$ invariant-mass distributions are fitted separately.

After unblinding the data set, there are 1 (2) candidates in the signal mass window $m(K^+ \mu^\pm e^\mp) \in [5100, 5370]$ MeV/ c^2 for the $B^+ \rightarrow K^+ \mu^- e^+$ ($B^+ \rightarrow K^+ \mu^+ e^-$) channels, respectively, in agreement with the background-only hypothesis (*cf.* Fig. 2). The upper limits on the branching fractions are set with the CL_s method [29], using the GammaCombo framework [30, 31] with a one-sided test statistic. The likelihoods are computed from fits to the invariant-mass distributions with the normalisation constant constrained to its nominal value accounting for statistical and systematic uncertainties. Pseudoexperiments, in which the nuisance parameters are input at their best fit value and the background yield is varied according to its systematic uncertainty, are used for the evaluation of the test statistic. The resulting upper limits are shown in Fig. 3 and Table 2.

The dominant sources of systematic uncertainty on the upper limits are due to applied




Figure 2: Invariant-mass distributions of the (left) $B^+ \rightarrow K^+ \mu^- e^+$ and (right) $B^+ \rightarrow K^+ \mu^+ e^-$ candidates obtained on the combined data sets recorded in 2011 and 2012 with background only fit functions (blue continuous line) and the signal model normalised to 10 candidates (red dashed line) superimposed. The signal window is indicated with grey dotted lines. Difference between the two distributions arises from the effect of the $m(K^+ \ell^-)$ requirement.




Figure 3: Upper limits on the branching fractions of (left) $B^+ \rightarrow K^+ \mu^- e^+$ and (right) $B^+ \rightarrow K^+ \mu^+ e^-$ decays obtained on the combined data sets recorded in 2011 and 2012. The red solid line (black solid line with data points) corresponds to the distribution of the expected (observed) upper limits, and the light blue (dark blue) band contains the 1σ (2σ) uncertainties.

simulation corrections. These include the kinematic difference of B -meson production, residual difference between correcting the muon and electron candidates, and PID resampling. Furthermore, the determination of trigger efficiencies and the knowledge of the background invariant-mass distribution are also considered in evaluating the systematic uncertainty.

Table 2: Upper limit on the branching fraction of $B^+ \rightarrow K^+ \mu^- e^+$ and $B^+ \rightarrow K^+ \mu^+ e^-$ decays obtained on the combined data sets recorded in 2011 and 2012 for confidence levels of 90% and 95%.

	90% C. L.	95% C. L.
$\mathcal{B}(B^+ \rightarrow K^+ \mu^- e^+)/10^{-9}$	7.0	9.5
$\mathcal{B}(B^+ \rightarrow K^+ \mu^+ e^-)/10^{-9}$	6.4	8.8

Table 3: Summary of systematic uncertainties.

Effect	$B^+ \rightarrow K^+ \mu^+ e^-$	$B^+ \rightarrow K^+ \mu^- e^+$
Data-simulation corrections	1.0%	1.0%
Electron-muon differences	1.4%	1.4%
Fitting model	2.1%	2.1%
PID resampling	4.5%	5.5%
Trigger	1.0%	1.0%
Normalisation factor	3.5%	3.5%
Total	6.4%	7.1%
Background	0.60	0.43

The systematic uncertainty on the sampling procedure of the PID variables includes two components. The first stems from applying the *sPlot* [32] method to the calibration data, and adds an uncertainty of 0.1% for kaons and muons, and 3% for electrons, the latter being a conservative estimate originating from a comparison of the *sPlot* method with a cut-and-count method. The second component addresses the choice of binning in the sampling procedure. It is evaluated by recalculating the normalisation factor α using a finer and a coarser binning, and taking the largest deviation with respect to the baseline result.

A small difference in the correction procedure is observed depending on the choice of control channel, namely $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ or $B^+ \rightarrow K^+ J/\psi (\rightarrow e^+ e^-)$. This difference, referred to as electron-muon difference, is taken as systematic uncertainty.

The systematic uncertainty from the fitting model is determined to be 2.1% using a bootstrapping approach. The systematic uncertainty on the background model is calculated by repeating the fit using an alternative model, where the exponential function is obtained from a sample enriched in background events. The difference between the alternative and nominal background parametrization is taken as a systematic uncertainty. The uncertainty on the knowledge of the $B^+ \rightarrow K^+ J/\psi (\rightarrow \mu^+ \mu^-)$ branching fraction is combined with the uncertainty due to the limited size of the simulation sample and is propagated to the normalisation constant, corresponding to a systematic uncertainty of 3.5%. A summary of systematic uncertainties is reported in Tab. 3.

In conclusion, a search for the lepton-flavour violating decays $B^+ \rightarrow K^+ \mu^\pm e^\mp$ is performed using data from proton-proton collisions recorded with the LHCb experiment at centre-of-mass energies of 7 TeV and 8 TeV, corresponding to an integrated luminosity of 3 fb^{-1} . A uniform distribution of signal events within the phase space accessible to the final-state particles is assumed. No excess is observed over the background-only hypothesis, and the resulting upper limits on the branching fractions are $\mathcal{B}(B^+ \rightarrow K^+ \mu^- e^+) < 7.0 (9.5) \times 10^{-9}$ and $\mathcal{B}(B^+ \rightarrow K^+ \mu^+ e^-) < 6.4 (8.8) \times 10^{-9}$ at 90 (95)% confidence level. The results improve the current best limits on the decays [9] by more than one order of magnitude. Moreover, the measurements impose strong constraints on the aforementioned extensions to the SM [5–8].

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).

References

- [1] M. Raidal *et al.*, *Flavour physics of leptons and dipole moments*, Eur. Phys. J. **C57** (2008) 13, [arXiv:0801.1826](#).
- [2] LHCb collaboration, R. Aaij *et al.*, *Search for lepton-universality violation in $B^+ \rightarrow K^+ \ell^+ \ell^-$ decays*, Phys. Rev. Lett. **122** (2019) 191801, [arXiv:1903.09252](#).
- [3] LHCb collaboration, R. Aaij *et al.*, *Test of lepton universality with $B^0 \rightarrow K^{*0} \ell^+ \ell^-$ decays*, JHEP **08** (2017) 055, [arXiv:1705.05802](#).
- [4] LHCb collaboration, R. Aaij *et al.*, *Angular analysis of the $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ decay using 3 fb^{-1} of integrated luminosity*, JHEP **02** (2016) 104, [arXiv:1512.04442](#).
- [5] I. de Medeiros Varzielas and G. Hiller, *Clues for flavor from rare lepton and quark decays*, JHEP **06** (2015) 072, [arXiv:1503.01084](#).
- [6] G. Hiller, D. Loose, and K. Schönwald, *Leptoquark flavor patterns & B decay anomalies*, JHEP **12** (2016) 027, [arXiv:1609.08895](#).
- [7] A. Crivellin, L. Hofer, J. Matias, U. Nierste, S. Pokorski, and J. Rosiek, *Lepton-flavour violating B decays in generic Z' models*, Phys. Rev. **D92** (2015) 054013, [arXiv:1504.07928](#).
- [8] S. M. Boucenna, J. W. F. Valle, and A. Vicente, *Are the B decay anomalies related to neutrino oscillations?*, Phys. Lett. **B750** (2015) 367, [arXiv:1503.07099](#).
- [9] BaBar collaboration, B. Aubert *et al.*, *Measurements of branching fractions, rate asymmetries, and angular distributions in the rare decays $B \rightarrow K \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$* , Phys. Rev. **D73** (2006) 092001, [arXiv:hep-ex/0604007](#).
- [10] Particle Data Group, M. Tanabashi *et al.*, *Review of particle physics*, Phys. Rev. **D98** (2018) 030001.
- [11] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [12] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [13] R. Aaij *et al.*, *The LHCb trigger and its performance in 2011*, JINST **8** (2013) P04022, [arXiv:1211.3055](#).
- [14] V. V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, JINST **8** (2013) P02013, [arXiv:1210.6861](#).
- [15] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [16] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).

- [17] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [18] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [19] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [20] Geant4 collaboration, J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. A. Dubois *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [21] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [22] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [23] Y. Freund and R. E. Schapire, *A decision-theoretic generalization of on-line learning and an application to boosting*, J. Comput. Syst. Sci. **55** (1997) 119.
- [24] CDF collaboration, T. Aaltonen *et al.*, *Search for $B_s^0 \rightarrow \mu^+ \mu^-$ and $B_d^0 \rightarrow \mu^+ \mu^-$ decays with $2fb^{-1}$ of $p\bar{p}$ collisions*, Phys. Rev. Lett. **100** (2008) 101802, [arXiv:0712.1708](#).
- [25] A. Bevan, R. G. Goñi, T. Stevenson, and T. Stevenson, *Support vector machines and generalisation in HEP*, J. Phys. Conf. Ser. **898** (2017) 072021, [arXiv:1702.04686](#).
- [26] R. Aaij *et al.*, *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, EPJ Tech. Instrum. **6** (2018) 1.
- [27] L. Anderlini *et al.*, *The PIDCalib package*, LHCb-PUB-2016-021.
- [28] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [29] A. L. Read, *Presentation of search results: The CL_s technique*, J. Phys. **G28** (2002) 2693.
- [30] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ from a combination of LHCb results*, JHEP **12** (2016) 087, [arXiv:1611.03076](#).
- [31] M. Kenzie, M. Karbach, T. Mombächer, M. Schlupp, and K. Schubert, *GammaCombo framework for combinations of measurements and computation of confidence intervals: Public Release v1.1*, Aug., 2019. doi: 10.5281/zenodo.3373613.
- [32] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).

LHCb collaboration

R. Aaij²⁸, C. Abellán Beteta⁴⁶, T. Ackernley⁵⁶, B. Adeva⁴³, M. Adinolfi⁵⁰, C.A. Aidala⁷⁷, Z. Ajaltouni⁶, S. Akar⁶¹, P. Albicocco¹⁹, J. Albrecht¹¹, F. Alessio⁴⁴, M. Alexander⁵⁵, A. Alfonso Alberio⁴², G. Alkhazov³⁴, P. Alvarez Cartelle⁵⁷, A.A. Alves Jr⁴³, S. Amato², Y. Amhis⁸, L. An¹⁸, L. Anderlini¹⁸, G. Andreassi⁴⁵, M. Andreotti¹⁷, J.E. Andrews⁶², F. Archilli¹³, P. d'Argent¹³, J. Arnau Romeu⁷, A. Artamonov⁴¹, M. Artuso⁶³, K. Arzymatov³⁸, E. Aslanides⁷, M. Atzeni⁴⁶, B. Audurier²³, S. Bachmann¹³, J.J. Back⁵², S. Baker⁵⁷, V. Balagura^{8,b}, W. Baldini^{17,44}, A. Baranov³⁸, R.J. Barlow⁵⁸, S. Barsuk⁸, W. Barter⁵⁷, M. Bartolini^{20,h}, F. Baryshnikov⁷⁴, V. Batozskaya³², B. Batsukh⁶³, A. Battig¹¹, V. Battista⁴⁵, A. Bay⁴⁵, M. Becker¹¹, F. Bedeschi²⁵, I. Bediaga¹, A. Beiter⁶³, L.J. Bel²⁸, V. Belavin³⁸, S. Belin²³, N. Bely⁶⁶, V. Bellee⁴⁵, K. Belous⁴¹, I. Belyaev³⁵, E. Ben-Haim⁹, G. Bencivenni¹⁹, S. Benson²⁸, S. Beranek¹⁰, A. Berezhnoy³⁶, R. Bernet⁴⁶, D. Berninghoff¹³, E. Bertholet⁹, A. Bertolin²⁴, C. Betancourt⁴⁶, F. Betti^{16,e}, M.O. Bettler⁵¹, M. van Beuzekom²⁸, I.a. Bezshyiko⁴⁶, S. Bhasin⁵⁰, J. Bhom³⁰, M.S. Bieker¹¹, S. Bifani⁴⁹, P. Billoir⁹, A. Birnkraut¹¹, A. Bizzeti^{18,u}, M. Bjørn⁵⁹, M.P. Blago⁴⁴, T. Blake⁵², F. Blanc⁴⁵, S. Blusk⁶³, D. Bobulska⁵⁵, V. Bocci²⁷, O. Boente Garcia⁴³, T. Boettcher⁶⁰, A. Boldyrev³⁹, A. Bondar^{40,x}, N. Bondar³⁴, S. Borghi^{58,44}, M. Borisyak³⁸, M. Borsato¹³, J.T. Borsuk³⁰, M. Boubdir¹⁰, T.J.V. Bowcock⁵⁶, C. Bozzi^{17,44}, S. Braun¹³, A. Brea Rodriguez⁴³, M. Brodski⁴⁴, J. Brodzicka³⁰, A. Brossa Gonzalo⁵², D. Brundu^{23,44}, E. Buchanan⁵⁰, A. Buonauro⁴⁶, C. Burr⁵⁸, A. Bursche²³, J.S. Butter²⁸, J. Buytaert⁴⁴, W. Byczynski⁴⁴, S. Cadeddu²³, H. Cai⁶⁸, R. Calabrese^{17,g}, S. Cali¹⁹, R. Calladine⁴⁹, M. Calvi^{21,i}, M. Calvo Gomez^{42,m}, A. Camboni^{42,m}, P. Campana¹⁹, D.H. Campora Perez⁴⁴, L. Capriotti^{16,e}, A. Carbone^{16,e}, G. Carboni²⁶, R. Cardinale^{20,h}, A. Cardini²³, P. Carniti^{21,i}, K. Carvalho Akiba²⁸, A. Casais Vidal⁴³, G. Casse⁵⁶, M. Cattaneo⁴⁴, G. Cavallero²⁰, R. Cenci^{25,p}, M.G. Chapman⁵⁰, M. Charles^{9,44}, Ph. Charpentier⁴⁴, G. Chatzikonstantinidis⁴⁹, M. Chefdeville⁵, V. Chekalina³⁸, C. Chen³, S. Chen²³, A. Chernov³⁰, S.-G. Chitic⁴⁴, V. Chobanova⁴³, M. Chruszcz⁴⁴, A. Chubykin³⁴, P. Ciambone¹⁹, M.F. Cicala⁵², X. Cid Vidal⁴³, G. Ciezarek⁴⁴, F. Cindolo¹⁶, P.E.L. Clarke⁵⁴, M. Clemencic⁴⁴, H.V. Cliff⁵¹, J. Closier⁴⁴, J.L. Cobbledick⁵⁸, V. Coco⁴⁴, J.A.B. Coelho⁸, J. Cogan⁷, E. Cogneras⁶, L. Cojocariu³³, P. Collins⁴⁴, T. Colombo⁴⁴, A. Comerma-Montells¹³, A. Contu²³, N. Cooke⁴⁹, G. Coombs⁵⁵, S. Coquereau⁴², G. Corti⁴⁴, C.M. Costa Sobral⁵², B. Couturier⁴⁴, G.A. Cowan⁵⁴, D.C. Craik⁶⁰, A. Crocombe⁵², M. Cruz Torres¹, R. Currie⁵⁴, C. D'Ambrosio⁴⁴, C.L. Da Silva⁷⁸, E. Dall'Occo²⁸, J. Dalseno^{43,50}, A. Danilina³⁵, A. Davis⁵⁸, O. De Aguiar Francisco⁴⁴, K. De Bruyn⁴⁴, S. De Capua⁵⁸, M. De Cian⁴⁵, J.M. De Miranda¹, L. De Paula², M. De Serio^{15,d}, P. De Simone¹⁹, C.T. Dean⁷⁸, W. Dean⁷⁷, D. Decamp⁵, L. Del Buono⁹, B. Delaney⁵¹, H.-P. Dembinski¹², M. Demmer¹¹, A. Dendek³¹, V. Denysenko⁴⁶, D. Derkach³⁹, O. Deschamps⁶, F. Desse⁸, F. Dettori²³, B. Dey⁶⁹, A. Di Canto⁴⁴, P. Di Nezza¹⁹, S. Didenko⁷⁴, H. Dijkstra⁴⁴, F. Dordei²³, M. Dorigo^{25,y}, A. Dosil Suárez⁴³, L. Douglas⁵⁵, A. Dovbnya⁴⁷, K. Dreimanis⁵⁶, M.W. Dudek³⁰, L. Dufour⁴⁴, G. Dujany⁹, P. Durante⁴⁴, J.M. Durham⁷⁸, D. Dutta⁵⁸, R. Dzhelyadin^{41,†}, M. Dziewiecki¹³, A. Dziurda³⁰, A. Dzyuba³⁴, S. Easo⁵³, U. Egede⁵⁷, V. Egorychev³⁵, S. Eidelman^{40,x}, S. Eisenhardt⁵⁴, U. Eitschberger¹¹, S. Ek-In⁴⁵, R. Ekelhof¹¹, L. Eklund⁵⁵, S. Ely⁶³, A. Ene³³, S. Escher¹⁰, S. Esen²⁸, T. Evans⁶¹, A. Falabella¹⁶, N. Farley⁴⁹, S. Farry⁵⁶, D. Fazzini⁸, P. Fernandez Declara⁴⁴, A. Fernandez Prieto⁴³, F. Ferrari^{16,e}, L. Ferreira Lopes⁴⁵, F. Ferreira Rodrigues², S. Ferreres Sole²⁸, M. Ferro-Luzzi⁴⁴, S. Filippov³⁷, R.A. Fini¹⁵, M. Fiorini^{17,g}, M. Firlej³¹, K.M. Fischer⁵⁹, C. Fitzpatrick⁴⁴, T. Fiutowski³¹, F. Fleuret^{8,b}, M. Fontana⁴⁴, F. Fontanelli^{20,h}, R. Forty⁴⁴, V. Franco Lima⁵⁶, M. Franco Sevilla⁶², M. Frank⁴⁴, C. Frei⁴⁴, D.A. Friday⁵⁵, J. Fu^{22,q}, W. Funk⁴⁴, M. Féo⁴⁴, E. Gabriel⁵⁴, A. Gallas Torreira⁴³, D. Galli^{16,e}, S. Gallorini²⁴, S. Gambetta⁵⁴, Y. Gan³, M. Gandelman², P. Gandini²², Y. Gao³, L.M. Garcia Martin⁷⁶, B. Garcia Plana⁴³, F.A. Garcia Rosales⁸, J. García Pardiñas⁴⁶, J. Garra Tico⁵¹, L. Garrido⁴², D. Gascon⁴², C. Gaspar⁴⁴, G. Gazzoni⁶,

D. Gerick¹³, E. Gersabeck⁵⁸, M. Gersabeck⁵⁸, T. Gershon⁵², D. Gerstel⁷, Ph. Ghez⁵,
 V. Gibson⁵¹, A. Gioventù⁴³, O.G. Girard⁴⁵, P. Gironella Gironell⁴², L. Giubega³³, K. Gizdov⁵⁴,
 V.V. Gligorov⁹, D. Golubkov³⁵, A. Golutvin^{57,74}, A. Gomes^{1,a}, I.V. Gorelov³⁶, C. Gotti^{21,i},
 E. Govorkova²⁸, J.P. Grabowski¹³, R. Graciani Diaz⁴², T. Grammatico⁹,
 L.A. Granado Cardoso⁴⁴, E. Graugés⁴², E. Graverini⁴⁵, G. Graziani¹⁸, A. Grecu³³, R. Greim²⁸,
 P. Griffith²³, L. Grillo⁵⁸, L. Gruber⁴⁴, B.R. Gruberg Cazon⁵⁹, C. Gu³, X. Guo⁶⁷, E. Gushchin³⁷,
 A. Guth¹⁰, Yu. Guz^{41,44}, T. Gys⁴⁴, C. Göbel⁶⁵, T. Hadavizadeh⁵⁹, C. Hadjivasiliou⁶,
 G. Haefeli⁴⁵, C. Haen⁴⁴, S.C. Haines⁵¹, P.M. Hamilton⁶², Q. Han⁶⁹, X. Han¹³, T.H. Hancock⁵⁹,
 S. Hansmann-Menzemer¹³, N. Harnew⁵⁹, T. Harrison⁵⁶, C. Hasse⁴⁴, M. Hatch⁴⁴, J. He⁶⁶,
 M. Hecker⁵⁷, K. Heijhoff²⁸, K. Heinicke¹¹, A. Heister¹¹, K. Hennessy⁵⁶, L. Henry⁷⁶,
 E. van Herwijnen⁴⁴, J. Heuel¹⁰, M. Heß⁷¹, A. Hicheur⁶⁴, R. Hidalgo Charman⁵⁸, D. Hill⁵⁹,
 M. Hilton⁵⁸, P.H. Hopchev⁴⁵, J. Hu¹³, W. Hu⁶⁹, W. Huang⁶⁶, Z.C. Huard⁶¹, W. Hulsbergen²⁸,
 T. Humair⁵⁷, R.J. Hunter⁵², M. Hushchyn³⁹, D. Hutchcroft⁵⁶, D. Hynds²⁸, P. Ibis¹¹, M. Idzik³¹,
 P. Ilten⁴⁹, A. Inglessi³⁴, A. Inyakin⁴¹, K. Ivshin³⁴, R. Jacobsson⁴⁴, S. Jakobsen⁴⁴, J. Jalocha⁵⁹,
 E. Jans²⁸, B.K. Jashal⁷⁶, A. Jawahery⁶², F. Jiang³, M. John⁵⁹, D. Johnson⁴⁴, C.R. Jones⁵¹,
 B. Jost⁴⁴, N. Jurik⁵⁹, S. Kandybei⁴⁷, M. Karacson⁴⁴, J.M. Kariuki⁵⁰, S. Karodia⁵⁵, N. Kazeev³⁹,
 M. Kecke¹³, F. Keizer⁵¹, M. Kelsey⁶³, M. Kenzie⁵¹, T. Ketel²⁹, B. Khanji⁴⁴, A. Kharisova⁷⁵,
 C. Khurewathanakul⁴⁵, K.E. Kim⁶³, T. Kirn¹⁰, V.S. Kirsebom⁴⁵, S. Klaver¹⁹,
 K. Klimaszewski³², S. Koliiev⁴⁸, M. Kolpin¹³, A. Kondybayeva⁷⁴, A. Konoplyannikov³⁵,
 P. Kopciewicz³¹, R. Kopečna¹³, P. Koppenburg²⁸, I. Kostiuk^{28,48}, O. Kot⁴⁸, S. Kotriakhova³⁴,
 M. Kozeiha⁶, L. Kravchuk³⁷, M. Kreps⁵², F. Kress⁵⁷, S. Kretzschmar¹⁰, P. Krokovny^{40,x},
 W. Krupa³¹, W. Krzemien³², W. Kucewicz^{30,l}, M. Kucharczyk³⁰, V. Kudryavtsev^{40,x},
 H.S. Kuindersma²⁸, G.J. Kunde⁷⁸, A.K. Kuonen⁴⁵, T. Kvaratskheliya³⁵, D. Lacarrere⁴⁴,
 G. Lafferty⁵⁸, A. Lai²³, D. Lancierini⁴⁶, J.J. Lane⁵⁸, G. Lanfranchi¹⁹, C. Langenbruch¹⁰,
 T. Latham⁵², F. Lazzari^{25,v}, C. Lazzaroni⁴⁹, R. Le Gac⁷, A. Leflat³⁶, R. Lefèvre⁶, F. Lemaitre⁴⁴,
 O. Leroy⁷, T. Lesiak³⁰, B. Leverington¹³, H. Li⁶⁷, P.-R. Li^{66,ab}, X. Li⁷⁸, Y. Li⁴, Z. Li⁶³,
 X. Liang⁶³, T. Likhomanenko⁷³, R. Lindner⁴⁴, P. Ling⁶⁷, F. Lionetto⁴⁶, V. Lisovsky⁸, G. Liu⁶⁷,
 X. Liu³, D. Loh⁵², A. Loi²³, J. Lomba Castro⁴³, I. Longstaff⁵⁵, J.H. Lopes², G. Loustau⁴⁶,
 G.H. Lovell⁵¹, D. Lucchesi^{24,o}, M. Lucio Martinez²⁸, Y. Luo³, A. Lupato²⁴, E. Luppi^{17,g},
 O. Lupton⁵², A. Lusiani²⁵, X. Lyu⁶⁶, R. Ma⁶⁷, F. Machefert⁸, F. Maciuc³³, V. Macko⁴⁵,
 P. Mackowiak¹¹, S. Maddrell-Mander⁵⁰, L.R. Madhan Mohan⁵⁰, O. Maev^{34,44}, A. Maevskiy³⁹,
 K. Maguire⁵⁸, D. Maisuzenko³⁴, M.W. Majewski³¹, S. Malde⁵⁹, B. Malecki⁴⁴, A. Malinin⁷³,
 T. Maltsev^{40,x}, H. Malygina¹³, G. Manca^{23,f}, G. Mancinelli⁷, D. Marangotto^{22,q}, J. Maratas^{6,w},
 J.F. Marchand⁵, U. Marconi¹⁶, C. Marin Benito⁸, M. Marinangeli⁴⁵, P. Marino⁴⁵, J. Marks¹³,
 P.J. Marshall⁵⁶, G. Martellotti²⁷, L. Martinazzoli⁴⁴, M. Martinelli^{44,21}, D. Martinez Santos⁴³,
 F. Martinez Vidal⁷⁶, A. Massafferri¹, M. Materok¹⁰, R. Matev⁴⁴, A. Mathad⁴⁶, Z. Mathe⁴⁴,
 V. Matiunin³⁵, C. Matteuzzi²¹, K.R. Mattioli⁷⁷, A. Mauri⁴⁶, E. Maurice^{8,b}, B. Maurin⁴⁵,
 M. McCann^{57,44}, L. McConnell¹⁴, A. McNab⁵⁸, R. McNulty¹⁴, J.V. Mead⁵⁶, B. Meadows⁶¹,
 C. Meaux⁷, G. Meier¹¹, N. Meinert⁷¹, D. Melnychuk³², M. Merk²⁸, A. Merli^{22,q}, E. Michielin²⁴,
 D.A. Milanes⁷⁰, E. Millard⁵², M.-N. Minard⁵, O. Mineev³⁵, L. Minzoni^{17,g}, S.E. Mitchell⁵⁴,
 B. Mitreska⁵⁸, D.S. Mitzel¹³, A. Mogini⁹, R.D. Moise⁵⁷, T. Mombächer¹¹, I.A. Monroy⁷⁰,
 S. Monteil⁶, M. Morandin²⁴, G. Morello¹⁹, M.J. Morello^{25,t}, J. Moron³¹, A.B. Morris⁷,
 A.G. Morris⁵², R. Mountain⁶³, H. Mu³, F. Muheim⁵⁴, M. Mukherjee⁶⁹, M. Mulder²⁸,
 C.H. Murphy⁵⁹, D. Murray⁵⁸, A. Mödden¹¹, D. Müller⁴⁴, J. Müller¹¹, K. Müller⁴⁶, V. Müller¹¹,
 P. Naik⁵⁰, T. Nakada⁴⁵, R. Nandakumar⁵³, A. Nandi⁵⁹, T. Nanut⁴⁵, I. Nasteva², M. Needham⁵⁴,
 N. Neri^{22,q}, S. Neubert¹³, N. Neufeld⁴⁴, R. Newcombe⁵⁷, T.D. Nguyen⁴⁵, C. Nguyen-Mau^{45,n},
 E.M. Niel⁸, S. Nieswand¹⁰, N. Nikitin³⁶, N.S. Nolte⁴⁴, D.P. O'Hanlon¹⁶,
 A. Oblakowska-Mucha³¹, V. Obraztsov⁴¹, S. Ogilvy⁵⁵, R. Oldeman^{23,f}, C.J.G. Onderwater⁷², J.
 D. Osborn⁷⁷, A. Ossowska³⁰, J.M. Otalora Goicochea², T. Ovsianikova³⁵, P. Owen⁴⁶,
 A. Oyanguren⁷⁶, P.R. Pais⁴⁵, T. Pajero^{25,t}, A. Palano¹⁵, M. Palutan¹⁹, G. Panshin⁷⁵,

A. Papanestis⁵³, M. Pappagallo⁵⁴, L.L. Pappalardo^{17,g}, W. Parker⁶², C. Parkes^{58,44},
 G. Passaleva^{18,44}, A. Pastore¹⁵, M. Patel⁵⁷, C. Patrignani^{16,e}, A. Pearce⁴⁴, A. Pellegrino²⁸,
 G. Penso²⁷, M. Pepe Altarelli⁴⁴, S. Perazzini¹⁶, D. Pereima³⁵, P. Perret⁶, L. Pescatore⁴⁵,
 K. Petridis⁵⁰, A. Petrolini^{20,h}, A. Petrov⁷³, S. Petrucci⁵⁴, M. Petruzzo^{22,q}, B. Pietrzyk⁵,
 G. Pietrzyk⁴⁵, M. Pikiés³⁰, M. Pili⁵⁹, D. Pinci²⁷, J. Pinzino⁴⁴, F. Pisani⁴⁴, A. Piucci¹³,
 V. Placinta³³, S. Playfer⁵⁴, J. Plews⁴⁹, M. Plo Casasus⁴³, F. Polci⁹, M. Poli Lener¹⁹,
 M. Poliakov⁶³, A. Poluektov⁷, N. Polukhina^{74,c}, I. Polyakov⁶³, E. Polycarpo², G.J. Pomery⁵⁰,
 S. Ponce⁴⁴, A. Popov⁴¹, D. Popov⁴⁹, S. Poslavskii⁴¹, K. Prasanth³⁰, C. Prouve⁴³, V. Pugatch⁴⁸,
 A. Puig Navarro⁴⁶, H. Pullen⁵⁹, G. Punzi^{25,p}, W. Qian⁶⁶, J. Qin⁶⁶, R. Quagliani⁹, B. Quintana⁶,
 N.V. Raab¹⁴, B. Rachwal³¹, J.H. Rademacker⁵⁰, M. Rama²⁵, M. Ramos Pernas⁴³, M.S. Rangel²,
 F. Ratnikov^{38,39}, G. Raven²⁹, M. Ravonel Salzgeber⁴⁴, M. Reboud⁵, F. Redi⁴⁵, S. Reichert¹¹,
 A.C. dos Reis¹, F. Reiss⁹, C. Remon Alepuz⁷⁶, Z. Ren³, V. Renaudin⁵⁹, S. Ricciardi⁵³,
 S. Richards⁵⁰, K. Rinnert⁵⁶, P. Robbe⁸, A. Robert⁹, A.B. Rodrigues⁴⁵, E. Rodrigues⁶¹,
 J.A. Rodriguez Lopez⁷⁰, M. Roehrken⁴⁴, S. Roiser⁴⁴, A. Rollings⁵⁹, V. Romanovskiy⁴¹,
 A. Romero Vidal⁴³, J.D. Roth⁷⁷, M. Rotondo¹⁹, M.S. Rudolph⁶³, T. Ruf⁴⁴, J. Ruiz Vidal⁷⁶,
 J. Ryzka³¹, J.J. Saborido Silva⁴³, N. Sagidova³⁴, B. Saitta^{23,f}, C. Sanchez Gras²⁸,
 C. Sanchez Mayordomo⁷⁶, B. Sanmartin Sedes⁴³, R. Santacesaria²⁷, C. Santamarina Rios⁴³,
 M. Santimaria^{19,44}, E. Santovetti^{26,j}, G. Sarpis⁵⁸, A. Sarti^{19,k}, C. Satriano^{27,s}, A. Satta²⁶,
 M. Saur⁶⁶, D. Savrina^{35,36}, L.G. Scantlebury Smead⁵⁹, S. Schael¹⁰, M. Schellenberg¹¹,
 M. Schiller⁵⁵, H. Schindler⁴⁴, M. Schmelling¹², T. Schmelzer¹¹, B. Schmidt⁴⁴, O. Schneider⁴⁵,
 A. Schopper⁴⁴, H.F. Schreiner⁶¹, M. Schubiger²⁸, S. Schulte⁴⁵, M.H. Schune⁸, R. Schwemmer⁴⁴,
 B. Sciascia¹⁹, A. Sciubba^{27,k}, A. Semennikov³⁵, A. Sergi^{49,44}, N. Serra⁴⁶, J. Serrano⁷,
 L. Sestini²⁴, A. Seuthe¹¹, P. Seyfert⁴⁴, M. Shapkin⁴¹, T. Shears⁵⁶, L. Shekhtman^{40,x},
 V. Shevchenko^{73,74}, E. Shmanin⁷⁴, J.D. Shupperd⁶³, B.G. Siddi¹⁷, R. Silva Coutinho⁴⁶,
 L. Silva de Oliveira², G. Simi^{24,o}, S. Simone^{15,d}, I. Skiba¹⁷, N. Skidmore¹³, T. Skwarnicki⁶³,
 M.W. Slater⁴⁹, J.G. Smeaton⁵¹, E. Smith¹⁰, I.T. Smith⁵⁴, M. Smith⁵⁷, M. Soares¹⁶,
 L. Soares Lavra¹, M.D. Sokoloff⁶¹, F.J.P. Soler⁵⁵, B. Souza De Paula², B. Spaan¹¹,
 E. Spadaro Norella^{22,q}, P. Spradlin⁵⁵, F. Stagni⁴⁴, M. Stahl⁶¹, S. Stahl⁴⁴, P. Stefko⁴⁵,
 S. Stefkova⁵⁷, O. Steinkamp⁴⁶, S. Stemmle¹³, O. Stenyakin⁴¹, M. Stepanova³⁴, H. Stevens¹¹,
 A. Stocchi⁸, S. Stone⁶³, S. Stracka²⁵, M.E. Stramaglia⁴⁵, M. Straticiu³³, U. Straumann⁴⁶,
 S. Strovkov⁷⁵, J. Sun³, L. Sun⁶⁸, Y. Sun⁶², K. Swientek³¹, A. Szabelski³², T. Szumlak³¹,
 M. Szymanski⁶⁶, S. T'Jampens⁵, S. Taneja⁵⁸, Z. Tang³, T. Tekampe¹¹, G. Tellarini¹⁷,
 F. Teubert⁴⁴, E. Thomas⁴⁴, K.A. Thomson⁵⁶, J. van Tilburg²⁸, M.J. Tilley⁵⁷, V. Tisserand⁶,
 M. Tobin⁴, S. Tolk⁴⁴, L. Tomassetti^{17,g}, D. Tonelli²⁵, D.Y. Tou⁹, E. Tournefier⁵, M. Traill⁵⁵,
 M.T. Tran⁴⁵, A. Trisovic⁵¹, A. Tsaregorodtsev⁷, G. Tuci^{25,44,p}, A. Tully⁵¹, N. Tuning²⁸,
 A. Ukleja³², A. Usachov⁸, A. Ustyuzhanin^{38,39}, U. Uwer¹³, A. Vagner⁷⁵, V. Vagnoni¹⁶,
 A. Valassi⁴⁴, S. Valat⁴⁴, G. Valenti¹⁶, H. Van Hecke⁷⁸, C.B. Van Hulse¹⁴, R. Vazquez Gomez⁴⁴,
 P. Vazquez Regueiro⁴³, S. Vecchi¹⁷, M. van Veghel²⁸, J.J. Velthuis⁵⁰, M. Veltri^{18,r},
 A. Venkateswaran⁶³, M. Vernet⁶, M. Veronesi²⁸, M. Vesterinen⁵², J.V. Viana Barbosa⁴⁴,
 D. Vieira⁶⁶, M. Vieites Diaz⁴⁵, H. Viemann⁷¹, X. Vilasis-Cardona^{42,m}, A. Vitkovskiy²⁸,
 V. Volkov³⁶, A. Vollhardt⁴⁶, D. Vom Bruch⁹, B. Voneki⁴⁴, A. Vorobyev³⁴, V. Vorobyev^{40,x},
 N. Voropaev³⁴, J.A. de Vries²⁸, C. Vázquez Sierra²⁸, R. Waldi⁷¹, J. Walsh²⁵, J. Wang⁴,
 J. Wang³, M. Wang³, Y. Wang⁶⁹, Z. Wang⁴⁶, D.R. Ward⁵¹, H.M. Wark⁵⁶, N.K. Watson⁴⁹,
 D. Websdale⁵⁷, A. Weiden⁴⁶, C. Weisser⁶⁰, B.D.C. Westhenry⁵⁰, D.J. White⁵⁸, M. Whitehead¹⁰,
 G. Wilkinson⁵⁹, M. Wilkinson⁶³, I. Williams⁵¹, M.R.J. Williams⁵⁸, M. Williams⁶⁰,
 T. Williams⁴⁹, F.F. Wilson⁵³, M. Winn⁸, W. Wislicki³², M. Witek³⁰, G. Wormser⁸,
 S.A. Wotton⁵¹, H. Wu⁶³, K. Wyllie⁴⁴, Z. Xiang⁶⁶, D. Xiao⁶⁹, Y. Xie⁶⁹, H. Xing⁶⁷, A. Xu³,
 L. Xu³, M. Xu⁶⁹, Q. Xu⁶⁶, Z. Xu³, Z. Xu⁵, Z. Yang³, Z. Yang⁶², Y. Yao⁶³, L.E. Yeomans⁵⁶,
 H. Yin⁶⁹, J. Yu^{69,aa}, X. Yuan⁶³, O. Yushchenko⁴¹, K.A. Zarebski⁴⁹, M. Zavertyaev^{12,c},
 M. Zeng³, D. Zhang⁶⁹, L. Zhang³, S. Zhang³, W.C. Zhang^{3,z}, Y. Zhang⁴⁴, A. Zhelezov¹³,

Y. Zheng⁶⁶, X. Zhou⁶⁶, Y. Zhou⁶⁶, X. Zhu³, V. Zhukov^{10,36}, J.B. Zonneveld⁵⁴, S. Zucchelli^{16,e}.

¹*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*

²*Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*

³*Center for High Energy Physics, Tsinghua University, Beijing, China*

⁴*Institute Of High Energy Physics (IHEP), Beijing, China*

⁵*Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*

⁶*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

⁷*Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*

⁸*LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France*

⁹*LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*

¹⁰*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*

¹¹*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*

¹²*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*

¹³*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

¹⁴*School of Physics, University College Dublin, Dublin, Ireland*

¹⁵*INFN Sezione di Bari, Bari, Italy*

¹⁶*INFN Sezione di Bologna, Bologna, Italy*

¹⁷*INFN Sezione di Ferrara, Ferrara, Italy*

¹⁸*INFN Sezione di Firenze, Firenze, Italy*

¹⁹*INFN Laboratori Nazionali di Frascati, Frascati, Italy*

²⁰*INFN Sezione di Genova, Genova, Italy*

²¹*INFN Sezione di Milano-Bicocca, Milano, Italy*

²²*INFN Sezione di Milano, Milano, Italy*

²³*INFN Sezione di Cagliari, Monserrato, Italy*

²⁴*INFN Sezione di Padova, Padova, Italy*

²⁵*INFN Sezione di Pisa, Pisa, Italy*

²⁶*INFN Sezione di Roma Tor Vergata, Roma, Italy*

²⁷*INFN Sezione di Roma La Sapienza, Roma, Italy*

²⁸*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*

²⁹*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*

³⁰*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*

³¹*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*

³²*National Center for Nuclear Research (NCBJ), Warsaw, Poland*

³³*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*

³⁴*Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia*

³⁵*Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia, Moscow, Russia*

³⁶*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*

³⁷*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*

³⁸*Yandex School of Data Analysis, Moscow, Russia*

³⁹*National Research University Higher School of Economics, Moscow, Russia*

⁴⁰*Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*

⁴¹*Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia*

⁴²*ICCUB, Universitat de Barcelona, Barcelona, Spain*

⁴³*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*

⁴⁴*European Organization for Nuclear Research (CERN), Geneva, Switzerland*

⁴⁵*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*

⁴⁶*Physik-Institut, Universität Zürich, Zürich, Switzerland*

⁴⁷*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*

⁴⁸*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*

⁴⁹*University of Birmingham, Birmingham, United Kingdom*

⁵⁰*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*

- ⁵¹ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵² *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵³ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁴ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁵ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁶ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁷ *Imperial College London, London, United Kingdom*
- ⁵⁸ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁹ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶⁰ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶¹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶² *University of Maryland, College Park, MD, United States*
- ⁶³ *Syracuse University, Syracuse, NY, United States*
- ⁶⁴ *Laboratory of Mathematical and Subatomic Physics , Constantine, Algeria, associated to ²*
- ⁶⁵ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶⁶ *University of Chinese Academy of Sciences, Beijing, China, associated to ³*
- ⁶⁷ *South China Normal University, Guangzhou, China, associated to ³*
- ⁶⁸ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³*
- ⁶⁹ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³*
- ⁷⁰ *Departamento de Física , Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁹*
- ⁷¹ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹³*
- ⁷² *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ²⁸*
- ⁷³ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³⁵*
- ⁷⁴ *National University of Science and Technology “MISIS”, Moscow, Russia, associated to ³⁵*
- ⁷⁵ *National Research Tomsk Polytechnic University, Tomsk, Russia, associated to ³⁵*
- ⁷⁶ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ⁴²*
- ⁷⁷ *University of Michigan, Ann Arbor, United States, associated to ⁶³*
- ⁷⁸ *Los Alamos National Laboratory (LANL), Los Alamos, United States, associated to ⁶³*
- ^a *Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil*
- ^b *Laboratoire Leprince-Ringuet, Palaiseau, France*
- ^c *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^d *Università di Bari, Bari, Italy*
- ^e *Università di Bologna, Bologna, Italy*
- ^f *Università di Cagliari, Cagliari, Italy*
- ^g *Università di Ferrara, Ferrara, Italy*
- ^h *Università di Genova, Genova, Italy*
- ⁱ *Università di Milano Bicocca, Milano, Italy*
- ^j *Università di Roma Tor Vergata, Roma, Italy*
- ^k *Università di Roma La Sapienza, Roma, Italy*
- ^l *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- ^m *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁿ *Hanoi University of Science, Hanoi, Vietnam*
- ^o *Università di Padova, Padova, Italy*
- ^p *Università di Pisa, Pisa, Italy*
- ^q *Università degli Studi di Milano, Milano, Italy*
- ^r *Università di Urbino, Urbino, Italy*
- ^s *Università della Basilicata, Potenza, Italy*
- ^t *Scuola Normale Superiore, Pisa, Italy*
- ^u *Università di Modena e Reggio Emilia, Modena, Italy*
- ^v *Università di Siena, Siena, Italy*
- ^w *MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*
- ^x *Novosibirsk State University, Novosibirsk, Russia*
- ^y *Sezione INFN di Trieste, Trieste, Italy*
- ^z *School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China*

^{aa} *Physics and Micro Electronic College, Hunan University, Changsha City, China*

^{ab} *Lanzhou University, Lanzhou, China*

[†] *Deceased*