

## LONGER CYCLES IN ESSENTIALLY 4-CONNECTED PLANAR GRAPHS

IGOR FABRICI<sup>1,2,a</sup>, JOCHEN HARANT<sup>1,b</sup>

SAMUEL MOHR<sup>1,3,b</sup> AND JENS M. SCHMIDT<sup>1,3,b</sup>

<sup>a</sup>*Pavol Jozef Šafárik University  
Institute of Mathematics, Košice, Slovakia*

<sup>b</sup>*Ilmenau University of Technology  
Department of Mathematics, Ilmenau, Germany*

**e-mail:** igor.fabrici@upjs.sk  
jochen.harant@tu-ilmenau.de  
samuel.mohr@tu-ilmenau.de  
jens.schmidt@tu-ilmenau.de

### Abstract

A planar 3-connected graph  $G$  is called *essentially 4-connected* if, for every 3-separator  $S$ , at least one of the two components of  $G - S$  is an isolated vertex. Jackson and Wormald proved that the length  $\text{circ}(G)$  of a longest cycle of any essentially 4-connected planar graph  $G$  on  $n$  vertices is at least  $\frac{2n+4}{5}$  and Fabrici, Harant and Jendroľ improved this result to  $\text{circ}(G) \geq \frac{1}{2}(n+4)$ . In the present paper, we prove that an essentially 4-connected planar graph on  $n$  vertices contains a cycle of length at least  $\frac{3}{5}(n+2)$  and that such a cycle can be found in time  $O(n^2)$ .

**Keywords:** essentially 4-connected planar graph, longest cycle, circumference, shortness coefficient.

**2010 Mathematics Subject Classification:** 05C38, 05C10.

### REFERENCES

<sup>1</sup>Partially supported by DAAD, Germany (as part of BMBF) and by the Ministry of Education, Science, Research and Sport of the Slovak Republic within the project 57320575.

<sup>2</sup>Partially supported by Science and Technology Assistance Agency under the contract No. APVV-15-0116 and by the Slovak VEGA Grant 1/0368/16.

<sup>3</sup> Gefördert durch die Deutsche Forschungsgemeinschaft (DFG) – 327533333 und 270450205; partially supported by the grants 327533333 and SCHM 3186/1-1 (270450205) from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), respectively.

- [1] M.B. Dillencourt, *Polyhedra of small order and their Hamiltonian properties*, J. Combin. Theory Ser. B **66** (1996) 87–122.  
doi:10.1006/jctb.1996.0008
- [2] I. Fabrici, J. Harant and S. Jendroř, *On longest cycles in essentially 4-connected planar graphs*, Discuss. Math. Graph Theory **36** (2016) 565–575.  
doi:10.7151/dmgt.1875
- [3] B. Grünbaum and J. Malkevitch, *Pairs of edge-disjoint Hamilton circuits*, Aequationes Math. **14** (1976) 191–196.  
doi:10.1007/BF01836218
- [4] B. Jackson and N.C. Wormald, *Longest cycles in 3-connected planar graphs*, J. Combin. Theory Ser. B **54** (1992) 291–321.  
doi:10.1016/0095-8956(92)90058-6
- [5] D.P. Sanders, *On paths in planar graphs*, J. Graph Theory **24** (1997) 341–345.  
doi:10.1002/(SICI)1097-0118(199704)24:4<341::AID-JGT6>3.0.CO;2-O
- [6] A. Schmid and J.M. Schmidt, *Computing Tutte paths* (2017), arXiv-preprint.  
<https://arxiv.org/abs/1707.05994>
- [7] C.-Q. Zhang, *Longest cycles and their chords*, J. Graph Theory **11** (1987) 521–529.  
doi:10.1002/jgt.3190110409

Received 16 October 2017  
Revised 13 March 2018  
Accepted 13 March 2018