

Table 1.10
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2001-2010

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.0*	-1.9*	-0.1	-2.1*	-2.6*	-1.7*
Oral Cavity & Pharynx:	-2.5*	-3.1*	-1.4	-3.9*	-4.1*	-3.2*
Lip	-	-	-	-	-	-
Tongue	-2.1*	-2.5*	-1.1	-3.9*	-4.3*	-2.4
Salivary gland	-1.0	-1.7	-0.5	0.2	1.0	-1.3
Floor of mouth	-7.4*	-8.2*	-5.9*	-10.5*	-	-
Gum & other oral cavity	-3.8*	-5.6*	-1.5	-5.7*	-6.0*	-5.0*
Nasopharynx	-1.6	-1.0	-2.2	-2.1	-2.1	-2.9
Tonsil	-0.2	-0.2	-0.2	-2.2	-2.8*	-0.4
Oropharynx	0.0	0.3	-	-2.5*	-2.5*	-2.3
Hypopharynx	-3.9*	-4.4*	-3.1	-3.1	-4.4	-
Other oral cavity & pharynx	-7.4*	-11.8*	-	-5.2*	-5.0*	-5.1*
Digestive System:	-1.2*	-1.2*	-1.3*	-1.7*	-1.5*	-2.1*
Esophagus	-4.5*	-5.0*	-3.1*	-4.8*	-4.6*	-5.2*
Stomach	-1.8*	-2.1*	-1.2	-3.2*	-3.1*	-3.5*
Small intestine	1.1	1.0	1.2	-0.1	-0.8	0.1
Colon & Rectum:	-2.3*	-2.4*	-2.4*	-2.8*	-2.4*	-3.3*
Colon	-2.7*	-3.0*	-2.6*	-	-	-
Rectum	-1.0*	-0.8	-1.4*	-	-	-
Anus, anal canal & anorectum	1.9*	2.7	0.9	4.8*	6.3*	3.1
Liver & intrahepatic bile duct	3.7*	3.7*	3.2*	2.4*	3.0*	0.8
Gallbladder	2.2*	4.8	1.8	0.4	4.3*	-0.8
Other biliary	2.4	0.7	3.5*	-1.8	-2.4	-1.9
Pancreas	0.5	1.3*	0.0	-0.1	0.0	-0.1
Retroperitoneum	0.3	-	-1.0	-5.5	-	-
Peritoneum, omentum & mesentery	-0.8	-	0.2	-0.7	-	-1.3
Other digestive system	3.5	8.4*	-0.5	-1.9	-3.0	-1.3
Respiratory System:	-1.7*	-2.9*	-0.2	-2.5*	-3.3*	-1.3*
Nose, nasal cavity & middle ear	-1.8	-2.5	-0.8	1.3	-0.7	4.2
Larynx	-3.1*	-3.7*	-1.1	-3.8*	-4.0*	-3.0*
Lung & bronchus	-1.7*	-2.8*	-0.2	-2.4*	-3.3*	-1.2*
Pleura	-	-	-	-	-	-
Trachea & other respiratory organs	-	-	-	-4.6	-	-
Bones & joints	0.8	0.2	1.7	-0.7	0.0	-1.8
Soft tissue (including heart)	0.2	-0.4	0.8	0.1	-0.8	0.7
Skin (excl. basal & squamous):	0.1	-0.8	0.6	-1.5*	-1.4	-1.5
Melanoma of the skin	-0.2	-1.5	0.6	0.1	1.9	-1.2
Other non-epithelial skin	0.4	-0.1	0.7	-3.0*	-3.3*	-1.9
Breast	0.3	1.5	0.4	-1.7*	-1.5	-1.6*
Breast (<i>in situ</i>)	2.1*	-	2.3*	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.10 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2001-2010

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-0.2	-	0.0	-1.2*	-	-1.0*
Cervix uteri	-3.2*	-	-3.0*	-2.6*	-	-2.4*
Corpus uteri	1.7*	-	1.9*	-1.1	-	-0.7
Uterus, NOS	2.0	-	2.2	1.5*	-	1.7*
Ovary ^c	-1.1	-	-0.9	-1.9*	-	-1.7*
Vagina	-2.2	-	-2.0	-3.4*	-	-3.1*
Vulva	-0.9	-	-0.8	-1.6	-	-1.5
Other female genital system	2.6	-	2.3	-0.4	-	-0.5
Male Genital System:	-2.1*	-2.6*	-	-3.4*	-3.7*	-
Prostate	-2.1*	-2.6*	-	-3.4*	-3.8*	-
Testis	-0.7	-0.7	-	-4.7	-5.2	-
Penis	-2.2	-2.3	-	1.8	1.8	-
Other male genital system	-	-	-	-	-	-
Urinary System:	1.5*	1.5*	1.4*	-1.0*	-0.7	-1.5*
Urinary bladder	-0.2	0.4	-1.2	-0.8	0.0	-1.9*
Kidney & renal pelvis	2.9*	2.5*	3.2*	-1.1*	-1.3*	-1.1*
Ureter	-	-	-	-	-	-
Other urinary system	0.6	-	-	-3.1	-	-3.4
Eye & Orbit	-0.4	-	-	-	-	-
Brain & Nervous System: ^d	-0.3	-0.2	-0.2	-0.5	-0.8	-0.3
Brain	0.2	0.0	0.4	-	-	-
Cranial nerves & other nervous system	-4.1	-3.2	-4.3	-	-	-
Endocrine System:	5.6*	4.8*	6.0*	1.5	2.5	1.0
Thyroid	5.7*	4.9*	6.1*	1.3	1.7	1.2
Other endocrine & thymus	4.6*	4.4*	4.7	1.8	3.3	0.7
Lymphoma:	0.0	0.0	0.0	-2.4*	-2.1*	-2.9*
Hodgkin lymphoma	1.3*	1.0	1.3	-2.3*	-3.1	-1.8
Non-Hodgkin lymphoma	-0.2	-0.2	-0.3	-2.5*	-2.0*	-3.0*
Myeloma	0.3	-0.4	0.7	-1.7*	-1.3*	-2.2*
Leukemia:	-1.7*	-1.9*	-1.6*	-1.6*	-1.3*	-1.9*
Lymphocytic:	-2.7*	-2.4	-3.1*	-2.2*	-1.7	-3.3*
Acute lymphocytic	1.2	3.4	-1.5	-2.8*	-2.7	-3.4*
Chronic lymphocytic	-3.8*	-3.8*	-3.6*	-2.1*	-1.4	-3.4*
Other lymphocytic	-2.3	-	-	-2.0	-3.4	-
Myeloid & Monocytic:	-0.3	-0.9	0.4	-1.6*	-1.4	-1.6*
Acute myeloid	0.3	-0.3	1.0	-0.3	-0.3	-0.3
Chronic myeloid	-0.9	-1.7	0.1	-7.5*	-7.0*	-7.6*
Acute monocytic	-	-	-	-	-	-
Other myeloid & monocytic	-	-	-	-1.1	-0.3	-3.3
Other leukemia:	-5.1*	-4.3	-5.7	-1.0	-0.8	-1.0
Other acute leukemia	-9.3*	-	-	-3.2*	-2.8*	-3.6
Aleukemic, subleukemic & NOS	-2.5	-1.8	-1.7	0.1	0.3	0.2
Kaposi Sarcoma ^e	-2.5*	-3.0*	-	-	-	-
Mesothelioma ^e	-0.3	-0.8	-	-	-	-
Ill-defined & unspecified	-4.0*	-4.5*	-3.5*	-3.3*	-3.3*	-3.3*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.